



Contribution ID: 206

Type: Talk

A Deep Learning Based Estimator for Light Flavour Elliptic Flow in Heavy Ion Collisions at RHIC and LHC Energies

Tuesday, June 4, 2024 9:50 AM (20 minutes)

Recent developments on a deep learning feed-forward network for estimating elliptic flow (v_2) coefficients in heavy-ion collisions have shown us the prediction power of this technique. The success of the model is mainly the estimation of v_2 from final state particle kinematic information and learning the centrality and the transverse momentum (p_T) dependence of v_2 . The deep learning model is trained with Pb-Pb collisions at 5.02 TeV minimum bias events simulated with a multiphase transport model (AMPT). We extend this work to estimate v_2 for light-flavor identified particles such as π^\pm , K^\pm , and $p + \bar{p}$ in heavy-ion collisions at RHIC and LHC energies. The number of constituent quark (NCQ) scaling is also shown. The evolution of pT-crossing point of $v_2(p_T)$, depicting a change in meson- baryon elliptic flow at intermediate-pT, is studied for various collision systems and energies. The model is further evaluated by training it for different p_T regions. These results are compared with the available experimental data wherever possible for light hadrons.

See:

[1] Physical Review D 105, 114022 (2022)

[2] Phys. Rev. D 107, 094001 (2023)

Authors: Dr MISHRA, Aditya Nath (Jawaharlal Nehru University); BARNAFOLDI, Gergely (HUN-REN Wigner RCP); Mr MALLICK, Neelkamal (IIT Indore); SAHOO, Raghunath (IIT Indore, India); Mr PRASAD, Suraj (IIT Indore)

Presenter: BARNAFOLDI, Gergely (HUN-REN Wigner RCP)

Session Classification: Track1-LF

Track Classification: Light-flavours and Strangeness