

A potential approach to the $X(3872)$ thermal behavior

Miguel Ángel Escobedo

Universitat de Barcelona

Strangeness in Quark Matter

Grant PID2022-136224NB-C21 funded by:



Institut de Ciències del Cosmos
UNIVERSITAT DE BARCELONA



Work done in collaboration with N. Armesto, E. Ferreiro and V. López-Pardo. Accepted in PLB.

Outline

1 Introduction


2 Theoretical framework

3 Results

4 Conclusions

Quarkonia exotics


- There are Quarkonium-like particles that can not be explained by the simple quark-antiquark model.



Picture taken from Physics Reports 873 (2020)

Quarkonia exotics


- There are Quarkonium-like particles that can not be explained by the simple quark-antiquark model.
- Among them, we focus on the $X(3872)$, whose internal structure is still a matter of debate.



Picture taken from Physics Reports 873 (2020)


Quarkonia exotics

- There are Quarkonium-like particles that can not be explained by the simple quark-antiquark model.
- Among them, we focus on the $X(3872)$, whose internal structure is still a matter of debate.
- There are two competing models. The tetraquark and the hadronic molecule.



Picture taken from Physics Reports 873 (2020)

Quarkonia exotics



Picture taken from https://www.fz-juelich.de/en/ias/ias-4/research/exotic-hadrons/exotics_pad.jpg.

The internal structure of the $X(3872)$

How can we get information about the internal structure of the $X(3872)$?

- Quantum numbers

The internal structure of the $X(3872)$

How can we get information about the internal structure of the $X(3872)$?

- Quantum numbers
- Spectroscopy

The internal structure of the $X(3872)$

How can we get information about the internal structure of the $X(3872)$?

- Quantum numbers
- Spectroscopy
- Decays

The internal structure of the $X(3872)$

How can we get information about the internal structure of the $X(3872)$?


- Quantum numbers
- Spectroscopy
- Decays
- Studying how a thermal medium modifies its properties

The internal structure of the $X(3872)$

How can we get information about the internal structure of the $X(3872)$?

- Quantum numbers
- Spectroscopy
- Decays
- Studying how a thermal medium modifies its properties


The $X(3872)$ in heavy-ion collisions



Picture taken from
Phys.Rev.Lett. 128 (2022) 3,
032001

Conventional quarkonium in heavy-ion collisions


- Improved theoretical understanding in recent years.



Picture taken from Phys.Rev.D
108 (2023) 1, L011502

Conventional quarkonium in heavy-ion collisions


- Improved theoretical understanding in recent years.
- The heavy quarkonium potential has both a real and an imaginary part.



Picture taken from Phys.Rev.D
108 (2023) 1, L011502

Conventional quarkonium in heavy-ion collisions


- Improved theoretical understanding in recent years.
- The heavy quarkonium potential has both a real and an imaginary part.
- The origin of the imaginary part is the collision of quarkonium with medium particles.



Picture taken from Phys.Rev.D
108 (2023) 1, L011502

Conventional quarkonium in heavy-ion collisions

- Improved theoretical understanding in recent years.
- The heavy quarkonium potential has both a real and an imaginary part.
- The origin of the imaginary part is the collision of quarkonium with medium particles.
- In some limits, it is a good approximation to model quarkonium using a Schrödinger equation with a complex potential.



Picture taken from Phys.Rev.D
108 (2023) 1, L011502

Outline

- Our aim is to study how the potential of the $X(3872)$ is modified in a thermal medium.

Outline

- Our aim is to study how the potential of the $X(3872)$ is modified in a thermal medium.
- We assume that it is a **tetraquark**. In the future, we plan to do a similar study assuming that it is an hadronic molecule.

Outline

- Our aim is to study how the potential of the $X(3872)$ is modified in a thermal medium.
- We assume that it is a **tetraquark**. In the future, we plan to do a similar study assuming that it is an hadronic molecule.
- It is challenging because non-perturbative physics has a prominent role in exotic quarkonia.

Outline

- Our aim is to study how the potential of the $X(3872)$ is modified in a thermal medium.
- We assume that it is a **tetraquark**. In the future, we plan to do a similar study assuming that it is an hadronic molecule.
- It is challenging because non-perturbative physics has a prominent role in exotic quarkonia.
- Due to this, our aim is to obtain **qualitative** results. We use as insights results from perturbative computations, lattice QCD and the large N_c limit.

Outline

1 Introduction

2 Theoretical framework

3 Results

4 Conclusions

The Born-Oppenheimer approximation

- We assume that heavy quarks move non-relativistically around the center-of-mass, with velocity v .

The Born-Oppenheimer approximation

- We assume that heavy quarks move non-relativistically around the center-of-mass, with velocity v .
- $\Lambda_{QCD} \gg E \sim m_Q v^2$. From the point of view of the heavy quarks, the light particles move very fast.


The Born-Oppenheimer approximation

- We assume that heavy quarks move non-relativistically around the center-of-mass, with velocity v .
- $\Lambda_{QCD} \gg E \sim m_Q v^2$. From the point of view of the heavy quarks, the light particles move very fast.
- The effect of light particles and gluons can be encoded in a potential computed assuming that the heavy quarks are frozen and separated a given distance r .

The Born-Oppenheimer approximation

- We assume that heavy quarks move non-relativistically around the center-of-mass, with velocity v .
- $\Lambda_{QCD} \gg E \sim m_Q v^2$. From the point of view of the heavy quarks, the light particles move very fast.
- The effect of light particles and gluons can be encoded in a potential computed assuming that the heavy quarks are frozen and separated a given distance r .
- Two step-approximation:
 - ▶ Compute the potential taking the heavy quarks as static color sources.
 - ▶ Solve the Schrödinger equation.

The Born-Oppenheimer approximation



The $T = 0$ potential

- Ideally, we would like to use a lattice QCD potential for the tetraquark as starting point.

The $T = 0$ potential

- Ideally, we would like to use a lattice QCD potential for the tetraquark as starting point.
- However, it is not yet available. Computations with dynamical light quarks are expensive.

The $T = 0$ potential

- Ideally, we would like to use a lattice QCD potential for the tetraquark as starting point.
- However, it is not yet available. Computations with dynamical light quarks are expensive.
- Instead, we use hybrid data. We then make the approximation that the tetraquark potential would behave qualitatively similar to the hybrid potential.


The $T = 0$ potential

- Ideally, we would like to use a lattice QCD potential for the tetraquark as starting point.
- However, it is not yet available. Computations with dynamical light quarks are expensive.
- Instead, we use hybrid data. We then make the approximation that the tetraquark potential would behave qualitatively similar to the hybrid potential.
- Single channel approximation. We assume that the heavy quarks do not contribute to the spin of the tetraquark.

The $T = 0$ potential

- Ideally, we would like to use a lattice QCD potential for the tetraquark as starting point.
- However, it is not yet available. Computations with dynamical light quarks are expensive.
- Instead, we use hybrid data. We then make the approximation that the tetraquark potential would behave qualitatively similar to the hybrid potential.
- Single channel approximation. We assume that the heavy quarks do not contribute to the spin of the tetraquark.
- We use data taken from Phys. Rev. D 99(3), 034502 (2019).

The $T = 0$ potential



This potential is well fitted by the formula

$$V(r, 0) = \frac{A_{-1}}{r} + A_0 + A_2 r^2.$$

Lattice data is not sensitive to large r behavior. But, using Effective String Theory results, we know that at very large r it only grows linearly.

The finite T potential

The real part

We use the following assumption

$$V(\mathbf{p}) = \frac{V_{\text{vac}}(\mathbf{p})}{\epsilon(\mathbf{p}, m_D)},$$

where ϵ is the medium permittivity in the HTL approximation and m_D is the Debye mass.

$$\begin{aligned} \Re[V(r, m_D)] &= A_{-1} \left(m_D + \frac{e^{-m_D r}}{r} \right) + A_0 \\ &+ A_2 \left[\frac{6}{m_D^2} (1 - e^{-m_D r}) - \left(2r^2 + \frac{6r}{m_D} \right) e^{-m_D r} \right] \end{aligned}$$

The finite T potential

The real part

We use the following assumption

$$V(\mathbf{p}) = \frac{V_{\text{vac}}(\mathbf{p})}{\epsilon(\mathbf{p}, m_D)},$$

where ϵ is the medium permittivity in the HTL approximation and m_D is the Debye mass.


$$\Re[V(r, m_D)] = A_{-1} \left(m_D + \frac{e^{-m_D r}}{r} \right) + A_0 \\ + A_2 \left[\frac{6}{m_D^2} (1 - e^{-m_D r}) - \left(2r^2 + \frac{6r}{m_D} \right) e^{-m_D r} \right]$$

Rationale

This model was able to describe lattice quarkonium potential at finite T using m_D as a fitting parameter (Phys. Rev. D 101(5), 056010 (2020)).

The finite T potential


The real part



Plot taken from Phys. Rev. D 101(5), 056010 (2020)

The finite T potential

The real part



The finite T potential

The imaginary part

The imaginary part of the potential of *conventional* quarkonium has the following properties:

The finite T potential

The imaginary part

The imaginary part of the potential of *conventional* quarkonium has the following properties:

- At short distances it goes like r^2 because the medium sees quarkonium as a small dipole.

The finite T potential

The imaginary part

The imaginary part of the potential of *conventional* quarkonium has the following properties:

- At short distances it goes like r^2 because the medium sees quarkonium as a small dipole.
- At long distances, the heavy quarks are not correlated. Therefore, the imaginary part of the potential is equal to $-i$ times the decay width of a single heavy quark.

The finite T potential

The imaginary part

The imaginary part of the potential of *conventional* quarkonium has the following properties:


- At short distances it goes like r^2 because the medium sees quarkonium as a small dipole.
- At long distances, the heavy quarks are not correlated. Therefore, the imaginary part of the potential is equal to $-i$ times the decay width of a single heavy quark.
- Between these two limits it is a smoothly increasing function.

The imaginary part

The imaginary part

For example, in the HTL approximation

$$\Im V(r) = -\alpha_s C_F T \phi(m_D r).$$



The finite T potential

The imaginary part

In a tetraquark state treated in the BO approximation, the heavy quarks are in an octet state.

- When $r \rightarrow 0$ the medium sees the heavy quark pair as a non-relativistic heavy gluon. The imaginary part of the potential will be $-i/2$ times the decay width of a heavy gluon.

The finite T potential

The imaginary part

In a tetraquark state treated in the BO approximation, the heavy quarks are in an octet state.

- When $r \rightarrow 0$ the medium sees the heavy quark pair as a non-relativistic heavy gluon. The imaginary part of the potential will be $-i/2$ times the decay width of a heavy gluon.
- At large distances the two quarks are uncorrelated. Same as in the color singlet case.

The finite T potential

The imaginary part

In a tetraquark state treated in the BO approximation, the heavy quarks are in an octet state.

- When $r \rightarrow 0$ the medium sees the heavy quark pair as a non-relativistic heavy gluon. The imaginary part of the potential will be $-i/2$ times the decay width of a heavy gluon.
- At large distances the two quarks are uncorrelated. Same as in the color singlet case.
- At intermediate distances we expect that the imaginary part of the potential is a smooth function that interpolates between the two regimes.

The finite T potential

The imaginary part

In a tetraquark state treated in the BO approximation, the heavy quarks are in an octet state.

- When $r \rightarrow 0$ the medium sees the heavy quark pair as a non-relativistic heavy gluon. The imaginary part of the potential will be $-i/2$ times the decay width of a heavy gluon.
- At large distances the two quarks are uncorrelated. Same as in the color singlet case.
- At intermediate distances we expect that the imaginary part of the potential is a smooth function that interpolates between the two regimes.
- In the large N_c limit the decay width of a heavy gluon is equal to that of two heavy quarks.

The finite T potential

The imaginary part

In a tetraquark state treated in the BO approximation, the heavy quarks are in an octet state.

- When $r \rightarrow 0$ the medium sees the heavy quark pair as a non-relativistic heavy gluon. The imaginary part of the potential will be $-i/2$ times the decay width of a heavy gluon.
- At large distances the two quarks are uncorrelated. Same as in the color singlet case.
- At intermediate distances we expect that the imaginary part of the potential is a smooth function that interpolates between the two regimes.
- In the large N_c limit the decay width of a heavy gluon is equal to that of two heavy quarks.
- Therefore, we can take the imaginary part of the potential to be a constant.

The Decay Width

$$\Gamma = A_{-1} T + A_2 \frac{T}{m_D^3} .$$

The Decay Width

$$\Gamma = A_{-1} T + A_2 \frac{T}{m_D^3} .$$

- It is a function of the same constants that appear in the real part of the potential.

The Decay Width

$$\Gamma = A_{-1} T + A_2 \frac{T}{m_D^3} .$$

- It is a function of the same constants that appear in the real part of the potential.
- Dimensional analysis.

The Decay Width

$$\Gamma = A_{-1} T + A_2 \frac{T}{m_D^3} .$$

- It is a function of the same constants that appear in the real part of the potential.
- Dimensional analysis.
- It is a educated guess. However, note that our aim is to get a qualitative understanding.

Outline

1 Introduction

2 Theoretical framework

3 Results

4 Conclusions

The dissociation temperature

- Obtained by solving the Schrödinger equation using the complex potential.


The dissociation temperature

- Obtained by solving the Schrödinger equation using the complex potential.
- Since the imaginary part is a constant, it factors out.

The dissociation temperature

- Obtained by solving the Schrödinger equation using the complex potential.
- Since the imaginary part is a constant, it factors out.
- The dissociation temperature is the one in which we can no longer find bound state solutions. In our case, we obtain $T_d \sim 250 \text{ MeV}$.

Mean radius vs Debye mass



Survival probability

We consider:

Survival probability

We consider:

- A Bjorken expansion starting at $t_0 = 0.6$ fm.

Survival probability

We consider:

- A Bjorken expansion starting at $t_0 = 0.6$ fm.
- A linear relation between T and m_D .

Survival probability

We consider:

- A Bjorken expansion starting at $t_0 = 0.6$ fm.
- A linear relation between T and m_D .
- We stop the evolution at the time in which $T = 175$ MeV. Around the phase transition.

Survival probability

We consider:


- A Bjorken expansion starting at $t_0 = 0.6$ fm.
- A linear relation between T and m_D .
- We stop the evolution at the time in which $T = 175$ MeV. Around the phase transition.

Survival probability


$$S(t) = \exp\left[-\int_{t_0}^t d\tau \Gamma(T(\tau), \tau)\right],$$

Survival probability

Note that:




Survival probability



Note that:


- The initial temperature depends on the collision centrality and the point in which the bound state is produced.


Survival probability




Note that:

- The initial temperature depends on the collision centrality and the point in which the bound state is produced.
- If $T \gtrsim 250$ MeV the state melts.







- We did not include recombination.



- We did not include recombination.
- Average for p_{\perp} and y values accessible at LHCb.



- We did not include recombination.
- Average for p_{\perp} and y values accessible at LHCb.
- Preliminary result.



- We did not include recombination.
- Average for p_{\perp} and y values accessible at LHCb.
- Preliminary result.
- Cold Nuclear Matter model effects taken from Phys.Rev.D 105 (2022) 1, 014019.

Outline

1 Introduction

2 Theoretical framework

3 Results

4 Conclusions

Conclusions

- Studying exotic quarkonia in heavy-ion collisions can shed light on its internal structure.

Conclusions

- Studying exotic quarkonia in heavy-ion collisions can shed light on its internal structure.
- We have developed a qualitative model for the potential of the $X(3872)$.

Conclusions

- Studying exotic quarkonia in heavy-ion collisions can shed light on its internal structure.
- We have developed a qualitative model for the potential of the $X(3872)$.
- Our results indicates that it melts around 250 MeV. The effect of the decay width is mild in heavy-ion collisions.