

ID de Contribution: 226

Type: **Talk**

Detailed study of the production of Υ mesons in PbPb collisions with CMS

mardi 4 juin 2024 15:20 (20 minutes)

One of the fundamental aspects of the formation and interaction of heavy quark bound states in the quark-gluon plasma is the amount of their suppression, expected to be stronger for quarkonia with smaller binding energies. Indeed, past results show a significant suppression of $\Upsilon(1S)$ mesons in heavy ion collisions, with $\Upsilon(2S)$ mesons being even more suppressed. However, for the $\Upsilon(3S)$ meson, only upper limits have been reported so far. At the same time, features like polarization of quarkonia or feed-down contributions from excited states remain to be assessed. In this talk, we report the observation of the $\Upsilon(3S)$ meson and a detailed study of the production of Υ mesons in lead-lead collisions recorded by CMS. For the first time, the nuclear modification factor has been extended to the strongly suppressed $\Upsilon(3S)$ state. The results are compared with model calculations describing the dynamics of quarkonia in a hot environment, whereas future prospects are shown too.

Auteur principal: RANJAN PUJAHARI, Prabhat

Orateur: RANJAN PUJAHARI, Prabhat

Classification de Session: Track2-HF&Q

Classification de thématique: Heavy-Flavours & Quarkonia