

# Mass degeneracy of chiral partners at finite density and its discovery potential at present facilities

mardi 4 juin 2024 19:19 (1 minute)

The spectral functions of chiral partners should become degenerate when the QCD chiral symmetry is restored. The axial-vector spectra are experimentally more challenging to construct than those of vector mesons that directly couple to virtual photons and then to dileptons. Chiral mixing of the vector with axial-vector mesons is thus a key phenomenon to probe in-medium modifications of vector spectrum due to the onset of chiral symmetry restoration carried by the axial-vector counterpart. The mixing effect is expected to be stronger at higher density due to a mechanism driven by chiral anomalies [1], in striking contrast to the vanishing mixing at chiral crossover driven by thermal pions [2]. This feature encourages us to perform the experimental search in the high-density regime, where the recent experimental trend has begun to shift toward.

We propose that experiments at medium energies with paying attention to the new mixing mechanism, may provide a direct evidence of the chiral symmetry restoration. In this presentation, we focus on the density-induced mixing and the spectral functions of  $\phi$  and its chiral partner  $f_1(1420)$ . We present the invariant mass distribution of dileptons using a transport approach developed in [3, 4] under the conditions of the J-PARC E16 experiment as a prime example. A potential to discover the mass degeneracy of chiral partners at finite density as a signature of the partial chiral symmetry restoration is advocated.

## References

- [1] C. Sasaki, Anomaly-induced chiral mixing in cold and dense matter, *Phys. Rev. D* 106, 054034 (2022).
- [2] A. Sakai, M. Harada, C. Nonaka, C. Sasaki, K. Shigaki, and S. Yano, Probing the QCD phase transition with chiral mixing in dilepton production, 2308.03305, *arXiv*, nucl-th(2023).
- [3] W. Cassing, and E. L. Bratkovskaya, Parton transport and hadronization from the dynamical quasiparticle point of view, *Phys. Rev. C* 78, 034919 (2008).
- [4] P. Gubler, E. L. Bratkovskaya, and T. Song,  $\phi$  meson properties in nuclear matter from dilepton spectra in a transport approach, *EPJ Web Conf.* 274, 07015 (2022).

**Author:** EJIMA, Ren (Hiroshima University)

**Co-auteurs:** SASAKI, Chihiro (University of Wroclaw); SHIGAKI, Kenta (Hiroshima University); GUBLER, Philipp (JAEA)

**Orateur:** EJIMA, Ren (Hiroshima University)

**Classification de Session:** Posters

**Classification de thématique:** Bulk matter phenomena, QCD phase diagram and Critical point