

Tsallis-Thermometer as a QGP Indicator for Large & Small **Collisional Systems**

<u>Gergely Gábor Barnaföldi¹</u>, Gábor Bíró^{1,2}, Tamás Sándor Biró¹, Aditya Nath Mishra³, Guy Paić⁴


¹Wigner Research Centre for Physics, 29-33 Konkoly-Thege Miklós street, 1121, Budapest, Hungary ² Institute of Physics, Eötvös Loránd University, 1/A Pázmány Péter Sétány, H-1117 Budapest, Hungary. ³ University Centre For Research and Development (UCRD), Chandigarh University, Mohali-140413, Punjab, India ⁴ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Apartado Postal 70-543, México City 04510, México




Introduction

Recent experimental results present collectivity also in small systems with high-multiplicity. Today these phenomena are not completely understood: it is an important question whether the presence of the QGP is necessary for the observed collectivity or not. Moreover, the connection between the experimental observables and theories is not trivial. In our phenomenological study we introduce the 'Tsallisthermometer' as an indicator of QGP, that result in the description of the the smooth transition from small to large collisional systems. The method also works well with geometry-selection of an event and correlates well with spherocity classified events. Results enable us to qualitatively define the underlying event definition beyond the CDF definition.

Tsallis distributions	Hadron spectra and multiplicity scaling
Thermodynamically consistent fit function:	Scaling with multiplicity & center of mass energy
$\frac{d^2 N}{2\pi p_T dp_T dy}\Big _{y\approx 0} = Am_T \left[1 + \frac{q-1}{T}(m_T - m)\right]^{-\frac{q}{q-1}}$	$T(\sqrt{s_{NN}}, \langle N_{ch}/\eta \rangle, m) = T_0 + T_1 \ln \frac{\sqrt{s_{NN}}}{m} + T_2 \ln \ln \langle N_{ch}/\eta \rangle,$





Investigated experimental data: $\pi^{\pm}, K^{\pm}, K^{0}_{s}, K^{*0}, p(\bar{p}), \Phi, \Lambda^{0}, \Xi^{\pm}, \Sigma^{\pm}, \Xi^{0}, \Omega^{\pm}$ $\sqrt{s_{NN}} \in 130 \text{ GeV}, 13 \text{ TeV}$ pp, p-A and AA collisions Various multiplicity classes **Consistency:** $P=Ts+\mu n-\varepsilon$




 $q(\sqrt{s_{NN}}, \langle N_{ch}/\eta \rangle, m) = q_0 + q_1 \ln \frac{\sqrt{s_{NN}}}{m} + q_2 \ln \ln \langle N_{ch}/\eta \rangle,$

Strong correlation between the parameters has been recognized as function of multiplicity:









Scaling with UE activity



Underlying event activity can be quantitatively seen by the Tsallis thermometer. Geometrical scanning is correlates with spherocity classified events.





Equation of State

EoS can be obtained, due the thermodynamical



- → Hadron mass order
- \rightarrow Present evolution
- \rightarrow Branches compatible with hadron d.o.f.

