

Constraining NSIs through CEvNS measurements at the European Spallation Source

Sabya Sachi Chatterjee




IRN Neutrino Meeting
Subatech Nantes June 19-20, 2023

Based on **PRD 107 (2023) 5, 055019** by
S. S. Chatterjee, S. Lavignac, O. G. Miranda, and G. S. Garcia

Outline

- Brief introduction to Coherent Elastic Neutrino-Nucleus Scattering (CEvNS)
- Introduction to Non-Standard Interaction
- Exploring the sensitivities of European Spallation Source (ESS) to NSI through CEvNS measurements.

Coherent Elastic Neutrino-Nucleus Scattering (CEvNS)




A neutrino of any flavor scatters-off a nucleus as a whole, transferring some energy in the form of nuclear recoil.

Condition: $QR \leq 1$

Q → three momentum transfer

R → radius of the nucleus



The nucleon wave functions are in phase with each other. The cross-sections of the individual nucleons are added coherently.

The differential cross-section for this process:

$$\frac{d\sigma}{dT}(E_\nu, T) = \frac{G_F^2 M}{\pi} \left(1 - \frac{MT}{2E_\nu^2}\right) F(|\vec{q}|^2) (Q_W^V)^2$$

$$(Q_{W,\text{SM}}^V)^2 = [(Z g_V^p + N g_V^n)]^2 \longrightarrow \text{Weak vector charge in SM.}$$

$$g_V^p = \frac{1}{2} - 2 \sin^2 \theta_W \text{ and } g_V^n = -\frac{1}{2} \longrightarrow \text{Neutral current vector couplings}$$

T \longrightarrow Nuclear recoil energy


$F(|\vec{q}|^2)$ \longrightarrow Nuclear form factor, we use Helm parametrization.

$$\text{For, } \sin^2 \theta_W \sim \frac{1}{4},$$

$$\sigma \propto N^2$$

So, the cross-section is large in compared to other neutrino interactions !

Large cross-section



A trade-off between the large cross-section and the small-ness of the recoil energy!

Why do we study CE ν NS ?

1. As a probe of SM physics:

- Weak mixing angle
- Neutron *rms* radius

2. Physics beyond the SM:

- Non-standard interactions (NSIs) of neutrinos
- Generalized neutrino interactions
- Neutrino electromagnetic properties
- Light neutral-current (NC) mediators
- New tool to probe sterile neutrinos and non-unitarity of the PMNS matrix

A lot more !

Where do we study CE ν NS ?

Tremendous progress has been made for the measurement of CE ν NS !

Spallation Neutron Sources:

- COHERENT
- European Spallation Sources (ESS)
- Captain Mills Experiment

Reactor Sources:

- CONUS
- CONNIE
- NUCLEUS
- nuGeN
- RED-100
- RICOCHET

And many more!





COHERENT SNS

In 2017 only, the COHERENT collaboration reported the first measurement of CE ν NS, 43 yrs after its proposal !

This talk will cover NSI at ESS

Flux at the Spallation Neutron Sources

Pion decay at rest:




The differential fluxes are given as:

$$\frac{dN_{\nu_\mu}}{dE_\nu} = \xi \delta \left(E_\nu - \frac{m_\pi^2 - m_\mu^2}{2m_\pi} \right),$$

$$\frac{dN_{\bar{\nu}_\mu}}{dE_\nu} = \xi \frac{64 E_\nu^2}{m_\mu^3} \left(\frac{3}{4} - \frac{E_\nu}{m_\mu} \right),$$

$$\frac{dN_{\nu_e}}{dE_\nu} = \xi \frac{192 E_\nu^2}{m_\mu^3} \left(\frac{1}{2} - \frac{E_\nu}{m_\mu} \right).$$



→ Prompt component

→ Delayed components

$$\xi = r N_{\text{POT}} / 4\pi L^2.$$

Non-Standard Interactions (NSIs)

The presence of the effective 4-Fermi neutral current non-standard interactions (NSI) can be realized through the dimension-six operators as,

$$-\mathcal{L}_{NSI} = \frac{G_F}{\sqrt{2}} \sum_{\alpha, \beta, f} \varepsilon_{\alpha\beta}^{fC} [\bar{\nu}_\alpha \gamma^\mu (1 - \gamma^5) \nu_\beta] [\bar{f} \gamma_\mu P_C f]$$


C=L, R

$\alpha, \beta = e, \mu, \tau$ and $f = e, u, d$

For CEvNS :

$$\varepsilon_{\alpha\beta}^{qV} \equiv (\varepsilon_{\alpha\beta}^{qL} + \varepsilon_{\alpha\beta}^{qR}), \quad q = u, d$$

L. Wolfenstein PRD 17, 2369 (1978)



NC-interaction

CEvNS cross-section changes in presence of NSI

$$\frac{d\sigma}{dT}(E_\nu, T) = \frac{G_F^2 M}{\pi} \left(1 - \frac{MT}{2E_\nu^2}\right) F(|\vec{q}|^2) (Q_{W,NSI}^V)^2$$

$$(Q_{W,NSI}^V)^2 = \left[Z (g_V^p + 2\varepsilon_{\alpha\alpha}^{uV} + \varepsilon_{\alpha\alpha}^{dV}) + N (g_V^n + \varepsilon_{\alpha\alpha}^{uV} + 2\varepsilon_{\alpha\alpha}^{dV}) \right]^2 + \sum_{\beta \neq \alpha} \left| Z (2\varepsilon_{\alpha\beta}^{uV} + \varepsilon_{\alpha\beta}^{dV}) + N (\varepsilon_{\alpha\beta}^{uV} + 2\varepsilon_{\alpha\beta}^{dV}) \right|^2.$$

Possibilities to explore:

- Bound on the NSI parameters using the available data
- Expected sensitivities of the NSI parameters using the future data

Brief description of the European Spallation Source (ESS) setup

The European Spallation source (ESS) is a highly ambitious and multi-disciplinary research facility. Apart from producing the World's most intense pulsed neutron beams, it will also generate neutrino fluxes from low energy to high energy suitable for exploring neutrino oscillation and Coherent Elastic Neutrino Nucleus Scattering (CEvNS).

ESS (Sourced at Lund, Sweden)

Baseline	20 m
Proton Energy	2 GeV
Average Beam Power	5 MW
Protons on target (POT) Per year	2.8×10^{23}

One order of
magnitude increase in
flux w.r.t. the
COHERENT SNS
experiment

A number of detectors have been proposed !

For more details, see JHEP 02 (2020) 123, arXiv: 2211.10396

Properties of the detector materials

Detectors proposed for the study of CEvNS: CsI, Si, Ge, Xe, Ar, C₃F₈

Target nucleus	Z	N	Z/N	M (a.m.u)	R (fm)
Cs	55	78	0.71	132.91	4.83
I	53	74	0.72	126.90	4.83
Xe	54	78	0.69	131.29	4.79
Si	14	14	1	27.98	3.12
Ge	32	40	0.8	72.00	4.06
Ar	18	22	0.81	39.95	3.43
C	6	6	1	12.01	2.47
F	9	10	0.9	19.00	2.90





Suitable combination of different detectors (different n/p ratio) can help breaking the degeneracies arise naturally when considering the single or multiple NSI Couplings with up or down type of quarks !

Proposed detectors properties

Detector technology	Target nucleus	Detector mass (kg)	Steady-state background	E_{th}^{ee} (keV _{ee})	QF (%)	T_{th} (keV _{nr})	$\frac{\Delta E}{E}$ (%) at T_{th}	σ_0 (keV)
Cryogenic scintillator	CsI	22.5	10 ckkd	0.1	10	1	30	0.3
High-pressure gaseous TPC	Xe	20	10 ckkd	0.18	20	0.9	40	0.36
Charge-coupled device	Si	1	1 ckkd	0.007	20	0.16	60	0.096
p-type point contact HPGe	Ge	7	15 ckkd	0.12	20	0.6	15	0.09
Scintillating bubble chamber	Ar	10	0.1 c/kg-day	–	–	0.1	40	0.04
Standard bubble chamber	C ₃ F ₈	10	15 c/kg-day	–	–	0.2	40	0.08


See also, JHEP 02 (2020) 123

Run time : 3 yrs



Single detector analysis

One NSI at a time




degeneracies

Constraints on the NSI parameters

Target nucleus	Quark type	ε_{ee}^{qV}		$\varepsilon_{\mu\mu}^{qV}$	
		90% C.L.	2σ C.L.	90% C.L.	2σ C.L.
CsI	u	$[-0.055, 0.051]$ $\cup [0.35, 0.45]$	$[-0.067, 0.064]$ $\cup [0.34, 0.47]$	$[-0.027, 0.023]$ $\cup [0.38, 0.43]$	$[-0.034, 0.028]$ $\cup [0.37, 0.43]$
	d	$[-0.040, 0.047]$ $\cup [0.31, 0.41]$	$[-0.061, 0.057]$ $\cup [0.30, 0.42]$	$[-0.024, 0.020]$ $\cup [0.34, 0.38]$	$[-0.031, 0.024]$ $\cup [0.33, 0.39]$

Target nucleus	Quark type	$\varepsilon_{e\mu}^{qV}$		$\varepsilon_{e\tau}^{qV}$		$\varepsilon_{\mu\tau}^{qV}$	
		90% C.L.	2σ C.L.	90% C.L.	2σ C.L.	90% C.L.	2σ C.L.
CsI	u	$[-0.089, 0.089]$	$[-0.1, 0.1]$	$[-0.16, 0.16]$	$[-0.18, 0.18]$	$[-0.11, 0.11]$	$[-0.12, 0.12]$
	d	$[-0.08, 0.08]$	$[-0.09, 0.09]$	$[-0.14, 0.14]$	$[-0.16, 0.16]$	$[-0.096, 0.096]$	$[-0.11, 0.11]$

Combination of two detectors



Disallowed at 3sigma C.L.

(CsI+Si)

		$\varepsilon_{\mu\mu}^{qV}$		$\varepsilon_{e\mu}^{qV}$	
		90% C.L.	2σ C.L.	90% C.L.	2σ C.L.
u		$[-0.02, 0.016]$	$[-0.025, 0.022]$	$[-0.074, 0.074]$	$[-0.081, 0.081]$
		–	–		
d		$[-0.018, 0.015]$	$[-0.023, 0.02]$	$[-0.068, 0.068]$	$[-0.075, 0.075]$
		–	$\cup [0.34, 0.36]$		

Not only constrain the parameter space but also break the degeneracy!

Current constraints using CsI + LAr data (1 sigma)

$$\epsilon_{ee}^{dV} = [-0.027, 0.048] \cup [0.30, 0.39],$$

$$\epsilon_{ee}^{uV} = [-0.024, 0.045] \cup [0.34, 0.43],$$

$$\epsilon_{\mu\mu}^{dV} = [-0.012, 0.016] \cup [0.33, 0.37],$$

$$\epsilon_{\mu\mu}^{uV} = [-0.002, 0.001] \cup [0.37, 0.41].$$

$$\epsilon_{e\mu}^{dV} = [-0.071, 0.071], \quad \epsilon_{e\mu}^{uV} = [-0.081, 0.081],$$



$$\epsilon_{e\tau}^{dV} = [-0.12, 0.12], \quad \epsilon_{e\tau}^{uV} = [-0.13, 0.13],$$

$$\epsilon_{\mu\tau}^{dV} = [-0.087, 0.087], \quad \epsilon_{\mu\tau}^{uV} = [-0.098, 0.098].$$

JHEP 04 (2023) 035 by V. D. Romeri, O. G. Miranda,
D. K. Papoulias, G. S. Garcia, M. Tortola, and J.W.F. Valle



ESS will further improve the bounds !

Single detector analysis of two non-zero NSI parameters



$$N_{th} = C_e (2Z + N)^2 \left(\epsilon_{ee}^{uV} + \frac{Zg_V^p + Ng_V^n}{2Z + N} \right)^2 + C_\mu (2Z + N)^2 \left(\epsilon_{\mu\mu}^{uV} + \frac{Zg_V^p + Ng_V^n}{2Z + N} \right)^2$$



Characterises elliptical shape



$$N_{th} = \left[Zg_V^p + Ng_V^n + (2Z + N) \varepsilon_{ee}^{uV} + (Z + 2N) \varepsilon_{ee}^{dV} \right]^2 C_e + \left[Zg_V^p + Ng_V^n \right]^2 C_\mu$$


Characterises two linear bands

Combined analysis of two non-zero NSI parameters



Breaking some part of the degeneracies



Combined analysis of two non-zero NSI parameters




Degeneracy broken in a large parameter spaces with suitable combinations of the different target materials of the detectors.


Combined analysis of two non-zero NSI parameters

A few more examples:





Same gas TPC detector filled with Ar or Xe



Analysis with two different detectors

Suitable combination of two different detectors gives better sensitivity!

Summary


- The measurement of CEvNS has opened a new window to test SM as well as beyond SM physics opportunities.
- We have explored the physics potential of European Spallation Source in constraining the NC-NSI.
- The combined analysis of two ESS detectors significantly enhance the sensitivities and partially break degeneracies of the NSI parameters taken one at a time. CsI+Si provides one of the best sensitivities.
- The combined analysis of two detectors becomes even more powerful reducing the degeneracies arising naturally when considered two at a time. We identify CsI+Si, Xe+Si, Csi+C3F8, and Xe+C3F8 as the best possible combinations in this regard.
- Combining two different type of detectors provide better sensnitivity than filling the same gas TPC detector with Xe or Ar.
- Hope the sensitivities will further improve with more runtime and with better systematic uncertainties.

Thank You !

We use the following chi-squared function for the analysis at ESS

$$\chi^2(\kappa) = \min_{\xi} \left[\sum_i 2 \left\{ N_i(\kappa, \xi) - \tilde{N}_i + \tilde{N}_i \ln \left(\frac{\tilde{N}_i}{N_i(\kappa, \xi)} \right) \right\} + \left(\frac{\xi_{sig}}{\sigma_{sig}} \right)^2 + \left(\frac{\xi_{bg}}{\sigma_{bg}} \right)^2 \right]$$

Current status of 3ν parameters (3σ bound)



ArXiv: 2006.11237 by P. Salas et al., arXiv: 2007.14792 by Esteban et al., and arXiv: 2107.00532 by F. Capozzi et al.

Neutrino flavor eigenstates are related to the mass eigenstates as

$$|\nu_\alpha\rangle = \sum_{i=1}^3 U_{\alpha i}^* |\nu_i\rangle,$$

Where,

$$U = R(\theta_{23}) R(\theta_{13}, \delta_{\text{CP}}) R(\theta_{12})$$

$$= \begin{pmatrix} c_{12} c_{13} & s_{12} c_{13} & s_{13} e^{-i\delta_{\text{CP}}} \\ -s_{12} c_{23} - c_{12} s_{23} s_{13} e^{i\delta_{\text{CP}}} & c_{12} c_{23} - s_{12} s_{23} s_{13} e^{i\delta_{\text{CP}}} & s_{23} c_{13} \\ s_{12} s_{23} - c_{12} c_{23} s_{13} e^{i\delta_{\text{CP}}} & -c_{12} s_{23} - s_{12} c_{23} s_{13} e^{i\delta_{\text{CP}}} & c_{23} c_{13} \end{pmatrix}$$