

# ACTAR TPC: Performances



Active target: (Gaseous) detector in which the atoms of the gas are used as a target

## ✓ Drift region: principle

- Transparent to particles on 4 sides
- Wire field cage
- Homogeneous vertical drift electric field
- Double wire field cage: 2 mm/1 mm pitch

## ✓ Amplification region: principle

- Bulk Micromegas (CERN PCB workshop)
- Local gain reduction via pad polarization



## ✓ Segmented pad plane

- Micromegas
- transverse multiplicity  $\approx$  electron straggling:  $2 \times 2 \text{ mm}^2$  pads
- 16384 pads with very high density: challenge!
- Two solutions investigated

## ✓ Electronics: GET

GET electronics:

- 512 samples ADC readout depth x 16384 pads
- volume sampling in 8 Mega voxel
- adjustable gain, peaking time, individual trigger: pad per pad



Active target: (Gaseous) detector in which the atoms of the gas are used as a target

Physics cases addressed:

- Time Projection Chamber mode
  - Ideal for implantation/decay studies: proton(s) radioactivity, beta-delayed charged particle emission, ...
  
- Study of excitation functions (resonant scattering, etc...)
  - thick target, need to differentiate the reactions channels
  
- Reactions with (very) low intensity beams
  - thick target, mixed target, possibly no  $^{12}\text{C}$  contamination

Active target: (Gaseous) detector in which the atoms of the gas are used as a target

Physics cases addressed:

□ Time Projection Chamber mode

→ Ideal for implantation/decay studies: proton(s) radioactivity, beta-delayed charged particle emission, ...

Study of excitation functions (resonant scattering, etc...)

→ thick target, need to differentiate the reactions channels



Reactions with (very) low intensity beams

→ thick target, mixed target, possibly no  $^{12}\text{C}$  contamination

✓ Proton-decay branches from the  $10^+$  isomer in  $^{54}\text{Ni}$  (May 2019)

$^{54}\text{Ni}$  implantation – proton decay:  $\rightarrow$  10A MeV  $^{54}\text{Ni}$  beam in 900 mbar Ar(95%) + CF<sub>4</sub>(5%)

J. Giovinazzo et al. “4D imaging of proton radioactivity”  
Nature communications 2021



J. Giovinazzo (2020)




- ✓ Simultaneous observation of Ni track (6 MeV/pad) and proton tracks (60 keV/pad)
- ✓ Decay of  $T_{1/2} = 155$  ns isomer : OK!

✓ Proton-decay branches from the  $10^+$  isomer in  $^{54}\text{Ni}$  (May 2019)

$^{54}\text{Ni}$  implantation – proton decay:  $\rightarrow$  10A MeV  $^{54}\text{Ni}$  beam in 900 mbar Ar(95%) + CF<sub>4</sub>(5%)

J. Giovinazzo et al.: 4D imaging of proton radioactivity, Nature communications 2021

J. Giovinazzo (2020)



- Proton decay branches carry unusually high angular momentum


- Branching ratio approximated theoretically with potential model for barrier penetration & Shell Model calculation for the initial and final WF overlap  
 $\rightarrow$   $C^2S$  of the order of  $10^{-6}$

- the “high  $l$ ” orbitals that mediate proton radioactivity in this region are also active in super-heavy nuclei and responsible for magic numbers in these nuclei



✓ Proton-decay branches from the  $19/2^-$  isomer in  $^{53}\text{Co}$  (May 2019)




$^{53}\text{Co}$  implantation – proton decay:  $\rightarrow$  10A MeV  $^{53}\text{Co}$  beam in 400 mbar Ar(95%) + CF<sub>4</sub>(5%)








✓ Decay of  $T_{1/2} = 239 \text{ ms}$  isomer : OK  
 ✓ Measurement of 0.025 % BR : OK  
 (not affected by  $\beta$  background)

50 years after the discovery of proton radioactivity ( $^{53m}\text{Co}$ ), we reach a complete comprehension of this state

## L. Sarmiento et al.: Nature Communications (2023)



✓ Exotic decays in the region of  $^{48}\text{Ni}$  (May 2021): PhD thesis **A. Ortega Moral (LP2IB)**



- tracking of low energy protons
- multiple particle tracking in 3D, half-life, ...



Active target: (Gaseous) detector in which the atoms of the gas are used as a target

Physics cases addressed:

Time Projection Chamber mode

→ Ideal for implantation/decay studies: proton(s) radioactivity, beta-delayed charged particle emission, ...

□ Study of excitation functions (resonant scattering, etc...)


→ thick target, need to differentiate the reactions channels

Reactions with (very) low intensity beams


→ thick target, mixed target, possibly no  $^{12}\text{C}$  contamination

- ✓ “Classic” TTIK method (thick solid target, beam stopped inside):
  - 3 unknown:  $E_{CM}$ ,  $\theta_{CM}$ ,  $E^*$  but only 2 observables ( $\theta_{light}$ ,  $E_{light}$ )
  - unable to disentangle elastic and inelastic channels (no info on  $E^*$ )
- ✓ Active Target: one more kinematic parameter (stopping point of the beam-like particle)
  - full identification of the reaction
  - + reconstruction of double differential cross section ( $d^2\sigma/d\Omega dE$ )



B. Mauss, PhD thesis (GANIL)




Particle identification



Channel selection




Double differential cross section



✓ Commissioning of the 128x128 pad full detector

$^{18}\text{O}(p,p)$  and  $^{18}\text{O}(p,\alpha)$  excitation functions:  $\rightarrow$  3.2A MeV  $^{18}\text{O}$  beam in 100 mbar  $\text{iC}_4\text{H}_{10}$

B. Mauss, et al., NIM A 940, 498 (2019)



- ✓ Absolute cross section measurement
- ✓  $^{18}\text{O}(p,p)$ : c.m. energy resolution: 38(3) keV FWHM
- ✓  $^{18}\text{O}(p,\alpha)$ : c.m. energy resolution: 54(9) keV FWHM

$\rightarrow$  Resolution limited by the angular straggling of the ions in the gas



✓ Resonant scattering : Ecm resolution dominated by the angular straggling in the gas

→ Use pure H<sub>2</sub> gas instead of isobutane :



example with <sup>20</sup>O+p excitation function, initial energy = 5A MeV

	iC <sub>4</sub> H <sub>10</sub>	H <sub>2</sub>	CH <sub>2</sub>
Pressure for stopping the beam	206 mbar	2.87 bar	165 μm
dN <sub>proton</sub> /dE @ 3A MeV (protons/MeV)	2.4.10 <sup>20</sup>	7.2.10 <sup>20</sup>	2.2.10 <sup>20</sup>
Angular straggling for 5 MeV proton on 10 cm gas	11 mrad	6.5 mrad	12.6 mrad




- requires high pressure (partially accomplished)
- requires amplification system for pure mono/diatomic gas (GEM): ongoing

✓ Search for  $\alpha$ -cluster states in  $^{10}\text{B}$  (B. Mauss, PhD thesis – to be published)

$^6\text{Li}(\alpha,\alpha)$  elastic and inelastic excitation functions @ LNS, Catania



3+ state @ 7.9 MeV,  
large alpha decay width



Preliminary work, B. Mauss


✓ Beam line modifications required for  ${}^8\text{Li}(\alpha, n)$  reaction @ 1.2A MeV (Expt. M.G. Pellegriti – 2024)

→ CIME cyclotron has (very) low transport efficiency below 2A MeV

→ Energy degradation in the physics room (G2)

→ Large acceptance Qpoles to reach 100% transmission (for 1 charge state)

→ Energy dispersion due to straggling & degrader inhomogeneities: reconstruction of the incident beam energy with ToF (event by event): 0.3% resolution possible up to  $10^5$  pps






- ✓ New development for high intensity / heavy beams: electrostatic beam mask

High intensity / heavy beams will create important space charge in ACTAR

- Space charge will distort the drift electric field and create deformed tracks
- Beam region must be screened to minimize the distortions
- Construction of an electrostatic beam mask (ongoing)




*C. Rodriguez et al., NIM A768, 179 (2014)*

### mask with double wire planes

space charge density 140 pC/cm<sup>3</sup>

→ Equivalent: 10<sup>6</sup> Hz of <sup>136</sup>Xe @ 7A MeV in 100 mbar iC<sub>4</sub>H<sub>10</sub>

Simulations: R. Revenko (GANIL)



Active target: (Gaseous) detector in which the atoms of the gas are used as a target

Physics cases addressed:

Time Projection Chamber mode

→ Ideal for implantation/decay studies: proton(s) radioactivity, beta-delayed charged particle emission, ...

Study of excitation functions (resonant scattering, etc...)

→ thick target, need to differentiate the reactions channels




□ Reactions with (very) low intensity beams

→ thick target, mixed target, possibly no  $^{12}\text{C}$  contamination

✓ Study of the  $^{19}\text{N}(d, ^3\text{He})$  reaction (2020-2021-2022)

$^{19}\text{N}$  at 30A MeV in 1 bar  $\text{D}_2(90\%) + \text{iC}_4\text{H}_{10}(10\%) \rightarrow$  Equivalent **11 mg/cm<sup>2</sup> CD<sub>2</sub> target + 5.4 mg/cm<sup>2</sup> CH<sub>2</sub> target**

**J. Lois Fuentes PhD thesis (U. Santiago de Compostella)**



**Solid state detectors can be placed on 4 sides (according to the reaction kinematics)**








**PID can be performed for a wide range of energy**



✓ Study of the  $^{19}\text{N}(d, ^3\text{He})$  reaction (2020-2021-2022)

$^{19}\text{N}$  at 30A MeV in 1 bar  $\text{D}_2(90\%) + \text{iC}_4\text{H}_{10}(10\%) \rightarrow$  Equivalent **11 mg/cm<sup>2</sup> CD<sub>2</sub> target + 5.4 mg/cm<sup>2</sup> CH<sub>2</sub> target**

**J. Lois Fuentes PhD thesis (U. Santiago de Compostella)**




**Thick target, E\* resolution limited by the Silicon energy resolution...**

✓ Study of the  $^{19}\text{N}(d, ^3\text{He})$  reaction (2020-2021-2022)



$^{19}\text{N}$  at 30A MeV in 1 bar  $\text{D}_2(90\%) + \text{iC}_4\text{H}_{10}(10\%) \rightarrow$  Equivalent **11 mg/cm<sup>2</sup> CD<sub>2</sub> target + 5.4 mg/cm<sup>2</sup> CH<sub>2</sub> target**

**J. Lois Fuentes PhD thesis (U. Santiago de Compostella)**



Center of mass angular coverage limited by:  
→ Backward cm angles: Si thickness


→ Add CsI to back Si layer !  
(+  $\gamma$  ray detection!)



✓ Study of the  $^{19}\text{N}(d, ^3\text{He})$  reaction (2020-2021-2022)

$^{19}\text{N}$  at 30A MeV in 1 bar  $\text{D}_2(90\%) + \text{iC}_4\text{H}_{10}(10\%) \rightarrow$  Equivalent **11 mg/cm<sup>2</sup> CD<sub>2</sub> target + 5.4 mg/cm<sup>2</sup> CH<sub>2</sub> target**


**J. Lois Fuentes PhD thesis (U. Santiago de Compostella)**



Center of mass angular coverage limited by:

- Backward cm angles: Si thickness
- Forward cm angles: Si trigger: delta electrons (inherent to the use of “high energy” beams) systematically trigger lots of pads (the gain is set high enough to see “high energy” light particles)

→ Use « mixed gain » pad plane (GET electronics feature) : 2/3rd of the pad plane in “low gain” configuration (i.e. transparent to deltas) + last 3rd with high gain (tracking of “high energy” light particles)



- ✓ Resonant scattering : Ecm resolution dominated by the angular straggling in the gas
  - Optimized with pure H<sub>2</sub> gas (proton scattering)
  - New setup for “low energy” beams ongoing
  - Technical development for high intensity/heavy beams
  
- ✓ Implantation / decay : Tested for 3 different experiments:
  - OK for lifetimes > ~ 100 ns.
  - OK for small branching ratio (no beta background)
  - OK for multiparticle tracking
  
- ✓ Transfer reactions : thick target without loss of resolution
  - (d,<sup>3</sup>He) reaction study with few 10<sup>4</sup> pps
  - Plans to increase the c.m. angular coverage



# ACTAR TPC Collaboration

