

The Sun as a cornerstone for (stellar) foundation science

European Nuclear Physics Conference 2025
Caen, France – September 22nd-26th - 2025

A. Serenelli



This project has received funding
from the European Union's
Horizon 2020 research and
innovation programme under
grant agreement No 101008324
(ChETEC-INFRA).

➤ Fundamental science with the Sun

solar atmosphere and theoretical spectroscopy

a view from helioseismology

a view from solar neutrinos – CN-vs, a unique opportunity

Sun as a calibrator for stellar ages


➤ Solar-like stars

convective cores in $1.1 - 1.5 M_{\odot}$ stars in the context of asteroseismic missions (PLATO)



➤ The need for accurate & precise cross section



Why the Sun? It is “foundation” science



- ~ 10^9 individual stars with measurements
colors, temperature, luminosity, (composition)
- ~ 10^3 with accurate, precise, (model) independent
mass determinations
selective club: eclipsing binaries
- **1 star with accurate, precise, (model) independent
age determination**
meteoritic dating
+ highly accurate radius & mass
- **Solar composition is the yardstick for all
astronomical abundance determinations**





Solar envelope is convective
→ hydrodynamic models
→ 3D atmosphere model

Model atmosphere
→ detailed radiative transfer
→ synthetic spectrum to compare with observed one
→ determination of abundances

Foundation science: Solar spectrum & abundances

Only star that allows detailed tests, e.g. center-to-limb variations




Pietrow, Hoppe, Bergemann et al. 2023

Bergemann, Hoppe, et al. 2021

(plots from M. Bergemann)


Which solar composition?



AAG21: Asplund et al. 2021
MB22: Magg et al. 2022

**Chemical abundances are a boundary condition
imposed to, not a prediction of,
(non-) standard solar models**

A unique view into the Sun: helioseismology




Helioseismology

- $>10^5$ eigenmodes → inversion of internal structure:
sound speed, density, adiabatic index (EoS)
- global quantities:
surface helium, depth of convective envelope
- beyond standard solar models:
internal rotation profile (depth and latitude)

Allows testing theory of stellar evolution by looking at internal structure


What helioseismology tells us



Model	$R_{CZ}/R_{\odot}$	Y_S	$\langle \delta c/c \rangle$	Y_{ini}	Z_{ini}
MB22	0.7123	0.2439	0.0010	0.2734	0.0176
AAG21	0.7197	0.2343	0.0027	0.2638	0.0155
GS98	0.7122	0.2425	0.0010	0.2718	0.0187
Solar	0.713 ± 0.001	0.2485 ± 0.0035			



What helioseismology tells us



Model	$R_{CZ}/R_{\odot}$	Y_S	$\langle \delta c/c \rangle$	Y_{ini}	Z_{ini}
MB22-phot	0.7123	0.2439	0.0010	0.2734	0.0176
AAG21	0.7197	0.2343	0.0027	0.2638	0.0155
GS98	0.7122	0.2425	0.0010	0.2718	0.0187
Solar	0.713 ± 0.001	0.2485 ± 0.0035			

Results sensitive to thermal structure because
sound speed scales with $T^{1/2}$

$$\nabla T^4 \propto \kappa \qquad c^2 \propto \frac{T}{\mu}$$

thermal gradient →
“composition + radiative opacities”

Another look from helioseismology


Propagation of sound waves carry information about composition through adiabatic index:

$$\Gamma_1 = \left(\frac{\partial \ln P}{\partial \ln \rho} \right)_{\text{ad}} = 5/3 \text{ (for fully ionized gas)} < 5/3 \text{ in partial ionization regions}$$

- It can be determined through inversion of solar oscillations and compared to solar models.
- Sensitive to total Z (very difficult to extract individual elements)

Composition + Equation of State


Low-Z favoured over high-Z



Buldgen et al. 2023

A unique view into the Sun: solar neutrinos


Solar neutrinos → information on solar core, nuclear physics



Model uncertainties >> experimental uncertainties!

What solar neutrinos from pp-chain tell us

Mostly a temperature sequence with slope determined by nuclear reaction rates



pp-chain neutrinos provide a measure of core temperature

cm ⁻² s ⁻¹	AAG21	MB22	Sun
⁷ Be (10 ⁹)	4.52 (7.4%)	4.90 (7.6%)	4.93 (2%)
⁸ B (10 ⁶)	4.31 (12.6%)	5.13 (13.1%)	5.20 (1.9%)

"Sun": experimental results from Gonzalez-García et al. 2024

Model uncertainties >> experimental ones

x-sections (S_{17} , S_{34} , S_{11})

6% for ⁷Be

8% for ⁸B



core thermal structure (comp. + opacities)



Dating the Sun “as a star”

Cancellation effects limit modes to $l=0, 1, 2, (3)$ for other stars (e.g. Kepler, TESS, PLATO)

$$\nu_{n,\ell} - \nu_{n-1,\ell+2} \propto \frac{1}{4\pi\nu_{n,\ell}} \int_0^R \boxed{\frac{dc}{dr} \frac{dr}{r}} \longrightarrow \text{sound speed gradient at solar/stellar core as age diagnostics}$$




$$c^2 \propto \frac{T}{\mu}$$

The Sun as a star – only independent age calibrator

No independent age for other stars

$$\nu_{n,\ell} - \nu_{n-1,\ell+2} \propto \frac{1}{4\pi\nu_{n,\ell}} \int_0^R \frac{dc}{dr} \frac{dr}{r}$$




	Solar age (Gyr)
Sun	4.568 ± 0.020
AAG21	4.790 ± 0.034
MB22	4.611 ± 0.032

Composition+opacities introduces a systematic effect on age determination of about to 250Myr (5%)



CN-cycle is a trace contribution to solar structure

CN operates against a “fixed” structure determined by pp-chains




Changes in physics affecting CN do not change structure, **i.e. core temperature**,
→ retain explicit dependences:

- e.g. linear response to bottleneck nuclear reaction
 $^{14}\text{N}(p, \gamma)^{15}\text{O}$
- **linear dependence on abundance of catalyzers in solar core: C+N**
- **one-to-one relation between neutrino fluxes and CN abundance**

What CN solar neutrinos tell us

^8B as a thermometer



Updated from Serenelli et al. 2013

Neutrino fluxes depend on:

➤ **solar core temperature – environmental quantities**

opacity
heavy elements (Si, Mg, Fe)
luminosity, age

uncertainties in these quantities affect n-fluxes in a fully correlated way

➤ **nuclear reaction rates**

specific dependence for specific fluxes
(e.g. $^{14}\text{N}(p,g)^{15}\text{O}$ does not affect pp-chain)


➤ **catalyzing effect of abundances**

C & N abundance in the solar core → CN-cycle



What CN solar neutrinos tell us

^8B as a thermometer



Neutrino fluxes as power-laws expansions:

$$\frac{\phi(^{15}\text{O})}{\phi(^{15}\text{O})_{\text{SSM}}} = [L_{\odot}^{5.942} O^{2.034} A^{1.364} D^{0.382}]$$

$$\times [S_{11}^{-2.912} S_{33}^{0.024} S_{34}^{-0.052} S_{17}^{0.0} S_{e7}^{0.0} S_{114}^{1.00}]$$

$$[x_C^{0.815} x_N^{0.217} x_O^{0.112} x_{\text{Ne}}^{0.081} x_{\text{Mg}}^{0.069} x_{\text{Si}}^{0.150} x_S^{0.109} x_{\text{Ar}}^{0.028} x_{\text{Fe}}^{0.397}]$$

$$\frac{\phi(^8\text{B})}{\phi(^8\text{B})_{\text{SSM}}} = [L_{\odot}^{6.966} O^{2.734} A^{1.319} D^{0.278}]$$


$$\times [S_{11}^{-2.665} S_{33}^{-0.419} S_{34}^{0.831} S_{17}^{1.028} S_{e7}^{-1} S_{114}^{0.00}]$$

$$[x_C^{0.022} x_N^{0.007} x_O^{0.128} x_{\text{Ne}}^{0.102} x_{\text{Mg}}^{0.092} x_{\text{Si}}^{0.198} x_S^{0.138} x_{\text{Ar}}^{0.034} x_{\text{Fe}}^{0.498}]$$



What CN solar neutrinos tell us

^8B as a thermometer




Thermal uncertainties are cancelled out, absorbed by a ^8B experimental measurement, down to 0.3%

$$\frac{\phi(^{15}\text{O})}{\phi(^{15}\text{O})_{\text{SSM}}} / \left[\frac{\phi(^8\text{B})}{\phi_{\text{SSM}}(^8\text{B})} \right]^{0.785} = x_C^{0.794} x_N^{0.212} D^{0.172} \\ \times [L_\odot^{0.515} O^{-0.016} A^{0.308}] \\ \times [S_{11}^{-0.831} S_{33}^{0.342} S_{34}^{-0.685} S_{17}^{-0.785} S_{e7}^{0.785} S_{114}^{0.995}] \\ \times [x_O^{0.003} x_{\text{Ne}}^{-0.005} x_{\text{Mg}}^{-0.003} x_{\text{Si}}^{-0.001} x_{\text{S}}^{-0.001} x_{\text{Ar}}^{0.001} x_{\text{Fe}}^{0.003}]$$

0.3% uncertainty

What CN solar neutrinos tell us

^8B as a thermometer



**Linear dependence
on C+N**


$$\frac{\phi(^{15}\text{O})}{\phi(^{15}\text{O})_{\text{SSM}}} / \left[\frac{\phi(^8\text{B})}{\phi_{\text{SSM}}(^8\text{B})} \right]^{0.785} = x_C^{0.794} x_N^{0.212} D^{0.172} \times [L_{\odot}^{0.515} O^{-0.016} A^{0.308}] \times [S_{11}^{-0.831} S_{33}^{0.342} S_{34}^{-0.685} S_{17}^{-0.785} S_{e7}^{0.785} S_{114}^{0.995}] \times [x_O^{0.003} x_{\text{Ne}}^{-0.001} x_{\text{Fe}}^{0.003}]$$

Nuclear reaction rates

$$\frac{\phi(^{15}\text{O})}{\phi(^{15}\text{O})_{\text{SSM}}} \simeq \left(\frac{\phi(^8\text{B})}{\phi(^8\text{B})_{\text{SSM}}} \right)^{0.785} (x_{\text{C+N}}) [1 \pm 0.10(\text{nuc}) + 0.03(\text{D})]$$

What CN solar neutrinos tell us

^8B as a thermometer



SuperKamiokande, SNO, others $\rightarrow$ 2% measurement of ^8B flux


$$\frac{\phi(^{15}\text{O})}{\phi(^{15}\text{O})_{\text{SSM}}} \simeq \left(\frac{\phi(^8\text{B})}{\phi(^8\text{B})_{\text{SSM}}} \right)^{0.785} (x_{\text{C+N}}) [1 \pm 0.10(\text{nuc}) + 0.03(\text{D})]$$

C+N abundance if ^{15}O flux is measured
(or any combination of ^{13}N & ^{15}O fluxes)

10% uncertainty (nuclear rates)
experimental uncertainty in CN-n fluxes dominated by



- $^{14}\text{N}(p,\gamma)^{15}\text{O}$ (8.5%)
- $^3\text{He}(^4\text{He},\gamma)^7\text{Be}$ (5%)

Borexino measurement of $^{13}\text{N}+^{15}\text{O}$ fluxes (Borexino coll. 2022, 2023)





What CN solar neutrinos tell us

$$\frac{\phi(^{15}\text{O})}{\phi(^{15}\text{O})_{\text{SSM}}} \simeq \left(\frac{\phi(^8\text{B})}{\phi(^8\text{B})_{\text{SSM}}} \right)^{0.785} (x_{\text{C+N}})[1 \pm 0.10(\text{nuc}) + 0.03(\text{D})]$$



What CN solar neutrinos tell us

$$\frac{\phi(^{15}\text{O})}{\phi(^{15}\text{O})_{\text{SSM}}} \simeq \left(\frac{\phi(^8\text{B})}{\phi(^8\text{B})_{\text{SSM}}} \right)^{0.785} (x_{\text{C+N}})[1 \pm 0.10(\text{nuc}) + 0.03(\text{D})]$$



CN neutrinos break the degeneracy between composition and opacity

Nuclear rates largest source of uncertainty, but one we can control



SFIII - Consensus (as of end of 2022) H-burning x-sections

INT WORKSHOP INT-22-82W

Solar Fusion Cross Sections III

July 26, 2022 - July 29, 2022


ORGANIZERS

Daniel Bemmerer
Helmholtz-Zentrum Dresden-Rossendorf
d.bemmerer@hzdr.de

Alessandra Guglielmetti
Università degli Studi di Milano - INFN
Milano
alessandra.guglielmetti@mi.infn.it

Wick Haxton
UC Berkeley
haxton@berkeley.edu

Aldo Serenelli
Institute of Space Sciences (ICE, CSIC)
aldos@ice.csic.es



Note to applicants: This workshop will be held in the David Brower Center near the UC Berkeley campus, Berkeley, CA.

WORKING GROUP AND
PRESENTATIONS WEBSITE



REVIEWS OF MODERN PHYSICS, VOLUME 97, JULY–SEPTEMBER 2025

Solar fusion III: New data and theory for hydrogen-burning stars

B. Acharya

Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

M. Aliotta

SUPA, School of Physics and Astronomy, University of Edinburgh, Edinburgh EH9 3FD, United Kingdom

A. B. Balantekin

Department of Physics, University of Wisconsin–Madison, Madison, Wisconsin 53706, USA

D. Bemmerer

Helmholtz-Zentrum Dresden-Rossendorf, 01328 Dresden, Germany

C. A. Bertulani

Department of Physics and Astronomy, Texas A&M University–Commerce, Commerce, Texas 75429-3011, USA

A. Best

*University of Napoli Federico II, 80126 Napoli, Italy
and Istituto Nazionale di Fisica Nucleare–Napoli, 80126 Napoli, Italy*

C. R. Brune

Edwards Accelerator Laboratory, Department of Physics and Astronomy, Ohio University, Athens, Ohio 45701, USA

R. Buompane and L. Gialanella

*Dipartimento di Matematica e Fisica, Università degli Studi della Campania “Luigi Vanvitelli,” 81100 Caserta, Italy
and Istituto Nazionale di Fisica Nucleare–Napoli, 80126 Napoli, Italy*

F. Cavanna

Istituto Nazionale di Fisica Nucleare–Torino, 10125 Torino, Italy

arXiv: 2405.06470
RMP, 97, 035002

H-burning x-sections.- uncertainties

Reaction	1 σ error (SFIII)	1 σ error (SFII, 2010)	
p+p	1.5%	0.9%	GA/GV
$^3\text{He}+^3\text{He}$	6.6%	5.2%	uncert. in protons spectral shape
$^3\text{He}+^4\text{He}$	5.1%	5%	
$^7\text{Be}+p$	3.6%	7.5%	halo EFT & Fitting
$^{14}\text{N}+p$	8.4%	7.2%	R-matrix and data tension for ground state transition


But see Chen et al.
arXiv: 2410.16086
15% offset

Not much improvement in terms of precision over the last (long) decade in key reactions


A concerted effort is needed to push down uncertainties to 1%-level



Impact for Solar-like stars




Mass distribution in the
Plato Input Catalogue




Impact for Solar-like stars: convective cores – to be or not to be

Stellar mass = $1.2 M_{\odot}$ - Sub-solar metallicity




Stellar mass = $1.2 M_{\odot}$ - Supra-solar metallicity



Nuclear cross section vs hydrodynamics (convection & overshooting)



Why 1% level precision x-sections




Constraints to compute solar models – composition is the big uncertainty:

- High solar metallicity is favored by (degenerate with opacity):
sound speed, surface helium, depth of convective envelope
pp-chain solar neutrinos
- but lower solar metallicity is favored by (degenerate with equation of state):
adiabatic index
- CNO neutrino break degeneracy with opacity → nuclear reactions main (and controllable) uncertainty
- Uncertainty in composition can be tamed by better CN neutrinos (models and experiments)

For solar-like oscillators:


- main target of PLATO missions – finding an Earth analog around solar-like stars
- $14\text{N}+p$ is fundamental to model evolution of stars in range 1.1 to 1.5 $M_{\odot}$
- empirical determinations of overshooting – the largest uncertainty in stellar modeling – are contingent to our knowledge of $14\text{N}+p$ rate




Orebi-Gann et al. 2021

Size of mixed core: convective boundary + overshooting

Size of chemically homogenous (mixed) core is formed by
+ truly convective core
+ **overshooting region** (parametrized, α_{ov} , no 1st principles model)
It can be measured with astereoseismology




Calibrating OV in stars
Only if N14+p well known




Deheuvels et al. 2016

Size of mixed core: convective boundary + overshooting

Size of chemically homogenous (mixed) core is formed by
+ truly convective core
+ **overshooting region** (parametrized, α_{ov} , no 1st principles model)
It can be measured with astereoseismology



Calibrating OV in M67
Only if N14+p well known



Size of mixed core: convective boundary + overshooting

Parametrization of overshooting for stars in the $1.1\text{--}1.4\ M_{\odot}$ range requires improved $14\text{N}+\text{p}$

Typical OV values are 0.2-0.25 $\rightarrow$ at low masses, uncertainty is very large

