

# Taking aim at the wino-higgsino plane with the LHC

*mercredi 25 octobre 2023 17:04 (20 minutes)*

In this work we explore multiple search strategies for higgsinos and mixed higgsino-wino states in the MSSM and project the results onto the  $(\mu, M_2)$  plane. Assuming associated production of higgsino-like pairs with a  $W/Z$  boson, we develop a search in a channel characterized by a hadronically tagged vector boson accompanied by missing energy. We use as our template an ATLAS search for dark matter produced in association with a hadronically decaying vector boson, but upgrade the search by implementing a joint likelihood analysis, binning the missing transverse energy distribution, which greatly improves the search sensitivity. For higgsino-like states (more than 96% admixture) we find sensitivity to masses up to 550 GeV. For well-mixed higgsino-wino states (70-30% higgsino) we still find sensitivities above 300 GeV. Using this newly proposed search, we draw a phenomenological map of the wino-higgsino parameter space, recasting several complementary searches for disappearing tracks, soft leptons, trileptons, and hadronic diboson events in order to predict LHC coverage of the  $(\mu, M_2)$  mass plane at integrated luminosities of up to  $3 \text{ ab}^{-1}$ . Altogether, the full run of the HL-LHC can exclude much of the “natural”  $(\mu, M_2 < 500 \text{ GeV})$  wino-higgsino parameter space.

**Auteur principal:** MURPHY, Taylor (LPTHE (Sorbonne U./CNRS))

**Co-auteurs:** GILMER, Humberto (Brown University); KAWAMURA, Junichiro (Keio University); CARPENTER, Linda (The Ohio State University)

**Orateur:** MURPHY, Taylor (LPTHE (Sorbonne U./CNRS))

**Classification de Session:** Beyond the Standard Model

**Classification de thématique:** BSM