
String theory, from the landscape to the swampland

Mariana Graña
Institut de Physique Théorique
CEA / Université Paris-Saclay
France

String theory


- A remarkable theory of quantum gravity, calculable
- Second superstring revolution ('95): unique theory!



Unique theory 10d

6d compact space

 $M_{10} = M_4 \times M_6$




Geometry of M₆ determines physics in 4d

Unique theory 10d

6d compact space

 $M_{10} = M_4 \times M_6$




Geometry of M₆ determines physics in 4d

Multiple solutions

Unique theory 10d


6d compact space

 $M_{10} = M_4 \times M_6$



Geometry of M₆ determines physics in 4d


Multiple solutions



Unique theory 10d


6d compact space

 $M_{10} = M_4 \times M_6$



Geometry of M₆ determines physics in 4d


Multiple solutions : 10500



Unique theory 10d

6d compact space


 $M_{10} = M_4 \times M_6$




Geometry of M₆ determines physics in 4d

Multiple solutions : 10500


Landscape of possible 4d "universes"



Simplest exemple




$$M_{10} = M_4 \times M_6$$
$$T_6$$




Observable world

Simplest exemple



$$M_{10} = M_4 \times \underline{M_6}$$
$$T_6$$



Observable world

Supersymmetric

→

supersymmetric spectrum

BUT


4D space = Minkowski ——


no cosmological constant

Radii of circles not fixed _____

Massless scalars (moduli)

Simplest exemple






Observable world

supersymmetric spectrum


no cosmological constant

Radii of circles not fixed _____

Massless scalars (moduli)



$$M_{10} = M_4 \times M_6$$




$$M_{10} = M_4 \times M_6$$

Eq. of motion: $E \ll M_{s} \lesssim M_{P}$

 $M_6:R_{mn}=0$ Ricci-flat

Calabi-Yau manifolds




 $M_{10} = M_4 \times M_6$

Still supersymmetric (less than for tori, $\mathcal{N}=1$)

Eq. of motion: $E \ll M_s \lesssim M_P$

 $M_6:R_{mn}=0$ Ricci-flat

Calabi-Yau manifolds



 $M_{10} = M_4 \times M_6$

Eq. of motion: $E \ll M_s \lesssim M_P$


 $M_6:R_{mn}=0$ Ricci-flat

Calabi-Yau manifolds

4D:

Still supersymmetric (less than for tori, $\mathcal{N}=1$)

Massless particles (e.g. the volume)



 $M_{10} = M_4 \times M_6$

Eq. of motion: $E \ll M_s \lesssim M_P$

M₆: R_{mn}=0 Ricci-flat


Calabi-Yau manifolds

4D:

Still supersymmetric (less than for tori, $\mathcal{N}=1$)

Massless particles (e.g. the volume)

4D space-time= Minkowski (no cosmological constant)



$$M_{10} = M_4 \times M_6$$

Richer than tori, can overcome problems

Eq. of motion: $E \ll M_s \lesssim M_P$

M₆: R_{mn}=0 Ricci-flat


Calabi-Yau manifolds

4D:

Still supersymmetric (less than for tori, $\mathcal{N}=1$)

Massless particles (e.g. the volume)

4D space-time= Minkowski (no cosmological constant)



$$M_{10} = M_4 \times M_6$$

Richer than tori, can overcome problems

Eq. of motion:
$$E \ll M_S \lesssim M_P$$

M₆: R_{mn}=0 Ricci-flat

Calabi-Yau manifolds


4D:

Still supersymmetric (less than for tori, $\mathcal{N}=1$)

Massless particles (e.g. the volume)

4D space-time= Minkowski (no cosmological constant)

Large number (finite?) of 6d CY and (for each of them) 10^{500} possibilities give masses to scalars and break supersymmetry



$$M_{10} = M_4 \times M_6$$

Richer than tori, can overcome problems

Eq. of motion:
$$E \ll M_S \lesssim M_P$$

M₆: R_{mn}=0 Ricci-flat

Calabi-Yau manifolds


Still supersymmetric (less than for tori, $\mathcal{N}=1$)

Massless particles (e.g. the volume)

4D space-time= Minkowski (no cosmological constant)

Large number (finite?) of 6d CY and (for each of them) 10^{500} possibilities give masses to scalars and break supersymmetry

Landscape of possibles 4d universes



Standard Model of particles

Positive cosmological constant (4d de Sitter space)

Standard Model of particles

2020: 10⁷⁰⁰ constructions with a Standard Model spectrum! (minimal supersymmetric), found using machine learning

Positive cosmological constant (4d de Sitter space)

Standard Model of particles

2020: 10⁷⁰⁰ constructions with a Standard Model spectrum! (minimal supersymmetric), found using machine learning Next step ~2025?: compute Yukawa couplings (need metric of CY, itself found using machine learning)

Positive cosmological constant (4d de Sitter space)

Standard Model of particles

2020: 10⁷⁰⁰ constructions with a Standard Model spectrum! (minimal supersymmetric), found using machine learning Next step ~2025?: compute Yukawa couplings (need metric of CY, itself found using machine learning)

Positive cosmological constant (4d de Sitter space)
 Much more difficult!!!

10D

Standard Model of particles

requiring in 4D:

2020: 10⁷⁰⁰ constructions with a Standard Model spectrum! (minimal supersymmetric), found using machine learning Next step ~2025?: compute Yukawa couplings (need metric of CY, itself found using machine learning)

Positive cosmological constant (4d de Sitter space)

Much more difficult!!!

2003: 10⁵⁰⁰ possibilities Kachru, Kallosh, Linde, Trivedi 2003 (KKLT)

Obtained using Calabi-Yau spaces, and the rest bottom-up, requiring in 4D 10D

Standard Model of particles

2020: 10⁷⁰⁰ constructions with a Standard Model spectrum! (minimal supersymmetric), found using machine learning Next step ~2025?: compute Yukawa couplings (need metric of CY, itself found using machine learning)

Positive cosmological constant (4d de Sitter space)

Much more difficult!!!

2003: 10⁵⁰⁰ possibilities Kachru, Kallosh, Linde, Trivedi 2003 (KKLT)

Many issues found @ IPhT (e.g. instabilities, singularities) Bena, M.G., students & post-docs 2009-on

Obtained using Calabi-Yau spaces, and the rest bottom-up, requiring in 4D 10D

Standard Model of particles

2020: 10⁷⁰⁰ constructions with a Standard Model spectrum! (minimal supersymmetric), found using machine learning Next step ~2025?: compute Yukawa couplings (need metric of CY, itself found using machine learning)

Positive cosmological constant (4d de Sitter space)

Much more difficult!!!

2003: 10⁵⁰⁰ possibilities Kachru, Kallosh, Linde, Trivedi 2003 (KKLT)

Many issues found @ IPhT (e.g. instabilities, singularities) Bena, M.G., students & post-docs 2009-on Ways around?

Obtained using Calabi-Yau spaces, and the rest bottom-up, requiring in 4D 10D

Standard Model of particles

2020: 10⁷⁰⁰ constructions with a Standard Model spectrum! (minimal supersymmetric), found using machine learning Next step ~2025?: compute Yukawa couplings (need metric of CY, itself found using machine learning)

Positive cosmological constant (4d de Sitter space)

Much more difficult!!!

2003: 10⁵⁰⁰ possibilities Kachru, Kallosh, Linde, Trivedi 2003 (KKLT)

Many issues found @ IPhT (e.g. instabilities, singularities) Bena, M.G., students & post-docs 2009-on Ways around?

Since 2003: Other solutions found, but far less solid

Obtained using Calabi-Yau spaces, and the rest bottom-up, requiring in 4D:

Standard Model of particles

2020: 10⁷⁰⁰ constructions with a Standard Model spectrum! (minimal supersymmetric), found using machine learning

Next step ~2025?: compute Yukawa couplings (need metric of CY, itself found using machine learning)

Positive cosmological constant (4d de Sitter space)

Much more difficult!!!

2003: 10⁵⁰⁰ possibilities Kachru, Kallosh, Linde, Trivedi 2003 (KKLT)

Many issues found @ IPhT (e.g. instabilities, singularities) Bena, M.G., students & post-docs 2009-on Ways around?

Since 2003: Other solutions found, but far less solid

The problem of bottom-up: are we really in the landscape?

Obtained using Calabi-Yau spaces, and the rest bottom-up, requiring in 4D:

Standard Model of particles

2020: 10⁷⁰⁰ constructions with a Standard Model spectrum! (minimal supersymmetric), found using machine learning

Next step ~2025?: compute Yukawa couplings (need metric of CY, itself found using machine learning)

Positive cosmological constant (4d de Sitter space)


Much more difficult!!!


2003: 10⁵⁰⁰ possibilities Kachru, Kallosh, Linde, Trivedi 2003 (KKLT)

Many issues found @ IPhT (e.g. instabilities, singularities) Bena, M.G., students & post-docs 2009-on Ways around?


Since 2003: Other solutions found, but far less solid

The problem of bottom-up: are we really in the landscape? Or did we fall into the swampland?






• Series of conjectures on properties of low energy effective Vafa + O(200) people, 2006-on field theories (EFT) to be in the landscape



• Series of conjectures on properties of low energy effective Vafa + O(200) people, 2006-on field theories (EFT) to be in the landscape This october school on swampland in Munich, 120 students!



• Series of conjectures on properties of low energy effective Vafa + O(200) people, 2006-on field theories (EFT) to be in the landscape This october school on swampland in Munich, 120 students!

Conjectures based on physics of black holes and/or inspired by string theory

An EFT theory is not in the swampland if

(i.e. it is consistent with quantum gravity if it satisfies)

An EFT theory is not in the swampland if


(i.e. it is consistent with quantum gravity if it satisfies)

No exact global symmetries (either gauged or broken)

An EFT theory is not in the swampland if

(i.e. it is consistent with quantum gravity if it satisfies)

• No exact global symmetries (either gauged or broken)



(i.e. it is consistent with quantum gravity if it satisfies)

• Weak gravity conjecture $:\exists$ a particle for which gravity is weaker than the gauge force m < q

(i.e. it is consistent with quantum gravity if it satisfies)

• Weak gravity conjecture $:\exists$ a particle for which gravity is weaker than the gauge force m < q



(i.e. it is consistent with quantum gravity if it satisfies)

• Weak gravity conjecture $:\exists$ a particle for which gravity is weaker than the gauge force m < q

extremal black hole


$$M = Q$$



(i.e. it is consistent with quantum gravity if it satisfies)

• Weak gravity conjecture $:\exists$ a particle for which gravity is weaker than the gauge force m < q


Black holes $M \ge Q$



extremal black hole


(i.e. it is consistent with quantum gravity if it satisfies)

• Weak gravity conjecture $:\exists$ a particle for which gravity is weaker than the gauge force m < q




(i.e. it is consistent with quantum gravity if it satisfies)

• Weak gravity conjecture $:\exists$ a particle for which gravity is weaker than the gauge force m < q



(i.e. it is consistent with quantum gravity if it satisfies)


• Weak gravity conjecture $:\exists$ a particle for which gravity is weaker than the gauge force m < q



(i.e. it is consistent with quantum gravity if it satisfies)

• Weak gravity conjecture $:\exists$ a particle for which gravity is weaker than the gauge force m < q

Black holes $M \ge Q$




superextremal black hole: naked singularity

(i.e. it is consistent with quantum gravity if it satisfies)

• Weak gravity conjecture $:\exists$ a particle for which gravity is weaker than the gauge force m < q

Black holes $M \ge Q$



⇒ extremal black holes cannot decay: remnants

• Conjectures based on physics of black holes and/or inspired by string theory

• Conjectures based on physics of black holes and/or inspired by string theory

• Some on very firm footing (like no global symmetries and weak gravity)

• Conjectures based on physics of black holes and/or inspired by string theory

• Some on very firm footing (like no global symmetries and weak gravity)

Some more speculative

• Conjectures based on physics of black holes and/or inspired by string theory

• Some on very firm footing (like no global symmetries and weak gravity)

Some more speculative


• The most speculative one: de Sitter swampland conjecture


- Conjectures based on physics of black holes and/or inspired by string theory
- Some on very firm footing (like no global symmetries and weak gravity)
- Some more speculative
- The most speculative one: de Sitter swampland conjecture


de Sitter space is in the swampland!!


• de Sitter swampland conjecture


Obied, Ooguri, Spodyneiko, Vafa 18 (OOSV)












In an EFT in the landscape

(inspired largely by our work)

The potential satisfies $|\nabla V| \ge cV$

or with c, c' > 0 (O(I) in Planck units)

$$\min(\nabla_i \nabla_j V) \le -c'V$$



$$\Phi^i$$
 (e.g. Φ = vol of M_6)

In an EFT in the landscape


(inspired largely by our work)

The potential satisfies $|\nabla V| \ge cV$

or

with c, c' > 0 (O(I) in Planck units)

$$\min(\nabla_i \nabla_j V) \le -c'V$$



$$\Phi^i$$
 (e.g. Φ = vol of M_6)


(inspired largely by our work)

In an EFT in the landscape

The potential satisfies $|\nabla V| \ge cV$

with c, c' > 0 (O(I) in Planck units) or

$$\min(\nabla_i \nabla_j V) \le -c'V$$



$$\Phi^i$$
 (e.g. Φ = vol of M_6)

(inspired largely by our work)


In an EFT in the landscape

The potential satisfies $|\nabla V| \ge cV$

or

with c, c' > 0 (O(I) in Planck units)

$$\min(\nabla_i \nabla_j V) \le -c' V$$



Metastable dS vacua are in the swampland

$$\Phi^i$$
 (e.g. Φ = vol of M_6)

(inspired largely by our work)


In an EFT in the landscape

The potential satisfies $|\nabla V| \ge cV$

or

with c, c' > 0 (O(I) in Planck units)

$$\min(\nabla_i \nabla_j V) \le -c' V$$



Metastable dS vacua are in the swampland

(e.g. Φ = vol of M_6)

If so, dark energy comes from quintessence field(s)

KKLT 03 10⁵⁰⁰ dS vacua





OOSV 18

No dS vacua

Complicated construction

uses non-perturbative ingredients

Demonstrated to have instabilities & other problems (with a way out?)




Complicated construction

uses non-perturbative ingredients

Demonstrated to have instabilities & other problems (with a way out?)




Speculative conjecture



(with a way out?)

Speculative conjecture



KKLT 03

10⁵⁰⁰ dS vacua

Complicated construction

uses non-perturbative ingredients

Demonstrated to have instabilities & other problems (with a way out?)



OOSV 18

No dS vacua

Speculative conjecture

KKLT 03

10⁵⁰⁰ dS vacua

Complicated construction

uses non-perturbative ingredients

Demonstrated to have instabilities & other problems (with a way out?)



OOSV 18

No dS vacua

Speculative conjecture

KKLT 03

10500

dS vacua

Complicated construction

uses non-perturbative ingredients

Demonstrated to have instabilities & other problems (with a way out?)



OOSV 18

No dS vacua


Speculative conjecture

Happy 60th birthday IPhT !! For many more to come

Summary

Summary


String theory is a remarkable and unique theory of quantum gravity




Summary

String theory is a remarkable and unique theory of quantum gravity


• Unique in 10d, but leads to many 4d effective theories




- Unique in 10d, but leads to many 4d effective theories
- Landscape of string theory 4d universes




- Unique in 10d, but leads to many 4d effective theories
- Landscape of string theory 4d universes
- Mostly explored: landscape CY compactifications




- Unique in 10d, but leads to many 4d effective theories
- Landscape of string theory 4d universes
- Mostly explored: landscape CY compactifications
 - Spectrum of standard model, Yukawa couplings to be computed



- Unique in 10d, but leads to many 4d effective theories
- Landscape of string theory 4d universes
- Mostly explored: landscape CY compactifications
 - Spectrum of standard model, Yukawa couplings to be computed
 - But no cosmological constant




- Unique in 10d, but leads to many 4d effective theories
- Landscape of string theory 4d universes
- Mostly explored: landscape CY compactifications
 - Spectrum of standard model, Yukawa couplings to be computed
 - But no cosmological constant
- Positive cosmological constant?



String theory is a remarkable and unique theory of quantum gravity

- Unique in 10d, but leads to many 4d effective theories
- Landscape of string theory 4d universes
- Mostly explored: landscape CY compactifications
 - Spectrum of standard model, Yukawa couplings to be computed
 - But no cosmological constant
- Positive cosmological constant?

All seems to indicate that if dS vacua in string theory



String theory is a remarkable and unique theory of quantum gravity


• Unique in 10d, but leads to many 4d effective theories

• Landscape of string theory 4d universes

- Mostly explored: landscape CY compactifications
 - Spectrum of standard model, Yukawa couplings to be computed
 - But no cosmological constant
- Positive cosmological constant?

All seems to indicate that if dS vacua in string theory

Actually the string landscape is mostly unexplored



String theory is a remarkable and unique theory of quantum gravity


• Unique in 10d, but leads to many 4d effective theories

• Landscape of string theory 4d universes

- Mostly explored: landscape CY compactifications
 - Spectrum of standard model, Yukawa couplings to be computed
 - But no cosmological constant
- Positive cosmological constant?

All seems to indicate that if dS vacua in string theory

- Actually the string landscape is mostly unexplored
 - Go beyond Calabi-Yau manifolds




String theory is a remarkable and unique theory of quantum gravity

- Unique in 10d, but leads to many 4d effective theories
- Landscape of string theory 4d universes
- Mostly explored: landscape CY compactifications
 - Spectrum of standard model, Yukawa couplings to be computed
 - But no cosmological constant
- Positive cosmological constant?

All seems to indicate that if dS vacua in string theory

- Actually the string landscape is mostly unexplored
 - Go beyond Calabi-Yau manifolds
 - Go beyond geometry!



String theory is a remarkable and unique theory of quantum gravity

• Unique in 10d, but leads to many 4d effective theories


• Landscape of string theory 4d universes

- Mostly explored: landscape CY compactifications
 - Spectrum of standard model, Yukawa couplings to be computed
 - But no cosmological constant
- Positive cosmological constant?

All seems to indicate that if dS vacua in string theory



- Go beyond Calabi-Yau manifolds
- Go beyond geometry!



11D supergravity

Type IIA

Type IIB-


String theory is a remarkable and unique theory of quantum gravity

- Unique in 10d, but leads to many 4d effective theories
- Landscape of string theory 4d universes
- Mostly explored: landscape CY compactifications
 - Spectrum of standard model, Yukawa couplings to be computed
 - But no cosmological constant
- Positive cosmological constant?

All seems to indicate that if dS vacua in string theory



- Go beyond Calabi-Yau manifolds
- Go beyond geometry!



Type I

11D supergravity

Type IIA

Type IIB-



Transplanckian Censorship Conjecture

Bedroya, Vafa 2019

• Sub-Planckian quantum fluctuations should remain quantum (cannot become larger than Hubble radius)

Or equivalently: The expansion of the universe must slow down before all sub-Planckian modes are stretched beyond the Hubble size

Two implications:

- no dS close to boundaries of moduli space, where

$$\frac{|\nabla V|}{V} \ge \frac{2}{\sqrt{(d-1)(d-2)}}$$

- dS can exist inside the bulk, if short lived

$$\tau \le \frac{1}{H} \log \frac{M_P}{H}$$