



DEEP UNDERGROUND
NEUTRINO EXPERIMENT

Sterile Neutrino search in DUNE




Camille Sironneau

DUNE France Workshop


16/11/23



Looking for
this little guy




Sterile neutrinos



While the 3 neutrino model is a good fit to most measurements, multiple anomalies have been detected by different experiments

If we add a new neutrino to the model, it has to be sterile i.e. interacting only through gravitational interaction and neutrino oscillation

Sterile neutrinos



RAA: anomalies measured by multiple experiments
in the total flux and energy spectrum of reactor
anti neutrinos

Also :

- excess of anti (electron neutrinos)
in anti (muon neutrinos) beam
was shown in LSND
- anomaly confirmed by MiniBooNE
- neutrino rate deficit in calibration
runs in the Gallium solar nu
experiments

→ Anomalies studied (and confirmed)
by multiple experiments but the best fit
point for 4nu model is usually ruled out
(i.e. STEREO)

Recent sterile studies

- **Reactor experiments :**
 - **Double Chooz**, explores $\Delta m^2 \sim 0.1 \text{ eV}^2$
 - **PROSPECT** and **STEREO** study the Reactor Antineutrino Anomaly (RAA) and have both excluded the best fit of RAA sterile nu with $\Delta m^2 \sim 1 \text{ eV}^2$ (papers from 2021 and 2023)
 - Future : **JUNO** with **RENO 50** sensitive to “super light sterile neutrino” ($\Delta m^2 \sim 10^{-5} \text{ eV}^2$)
- **Accelerator experiments :**
 - **LSND**, low energy beam of anti ν_{μ} , look for excess in ν_{e} events
 - **MiniBooNE**, higher energies but same L/E
 - **T2K**, no evidence of sterile mixing in “3+1” model (2019 paper)
- **Atmospherics / cosmic rays :**
 - **IceCube**, sensitive to high energy events
 - **KM3NeT**, no analysis yet but expected sensitivity
 - **SuperK**, set limit on sterile mixing to tau and mu (no delta sensitivity)
- **Gallium based solar nu experiments :**
 - **GALLEX**, **SAGE**, **BEST** : anomalies could be explained with $\Delta m^2 \sim 1 \text{ eV}^2$



Sterile neutrinos in DUNE

- We can use **beam neutrinos and higher L/E** to constrain more the sterile mixing parameters → ongoing work
- **Atmospheric neutrinos** can nicely complement the studies with beam neutrinos:
 - **wider range of L/E**
 - extra probe for sterile neutrinos with a **different source**
 - easier **comparison** with other ongoing or future experiments detecting atmospheric (KM3NeT, IceCube, Super-K, Hyper-K)
 - explore whether DUNE **energy and angular resolution** can be of advantage



(myself included)

Preliminary study

- Focus on **atmospheric neutrinos**
- The addition of a neutrino in the model changes oscillation probabilities
→ produce **appearance or disappearance** of neutrino flavours depending on energy/angle
- Compare expected number of events assuming Standard Model and “3+1” model with one sterile neutrino
- Example of calculation done for **numu** (everything is **also done for nue**)
- Focus on **CC interactions and NO** at first

$$N_{exp,ev} = [(\phi_{\nu_\mu} P_{\mu\mu} + \phi_{\nu_e} P_{e\mu}) \sigma_{\nu_\mu} + (\phi_{\bar{\nu}_\mu} P_{\bar{\mu}\mu} + \phi_{\bar{\nu}_e} P_{\bar{e}\mu}) \sigma_{\bar{\nu}_\mu}] \cdot N_{Ar} \cdot \Delta E \cdot \Delta\theta_z \cdot \Delta t$$

**Atmospheric neutrino flux
for different flavors → Honda flux for
Homestake**

Preliminary study

- Focus on **atmospheric neutrinos**
- The addition of a neutrino in the model changes oscillation probabilities
→ produce **appearance or disappearance** of neutrino flavours depending on energy/angle
- Compare expected number of events assuming Standard Model and “3+1” model with one sterile neutrino
- Example of calculation done for **numu** (everything is **also done for nue**)
- Focus on **CC interactions and NO** at first

$$N_{exp,ev} = [(\phi_{\nu_\mu} P_{\mu\mu} + \phi_{\nu_e} P_{e\mu}) \sigma_{\nu_\mu} + (\phi_{\bar{\nu}_\mu} P_{\mu\mu} + \phi_{\bar{\nu}_e} P_{e\mu}) \sigma_{\bar{\nu}_\mu}] \cdot N_{Ar} \cdot \Delta E \cdot \Delta\theta_z \cdot \Delta t$$

Oscillation probabilities to
numu or numu_bar

Preliminary study

- Focus on **atmospheric neutrinos**
- The addition of a neutrino in the model changes oscillation probabilities
→ produce **appearance or disappearance** of neutrino flavours depending on energy/angle
- Compare expected number of events assuming Standard Model and “3+1” model with one sterile neutrino
- Example of calculation done for **numu** (everything is **also done for nue**)
- Focus on **CC interactions and NO** at first

$$N_{exp,ev} = [(\phi_{\nu_\mu} P_{\mu\mu} + \phi_{\nu_e} P_{e\mu}) \sigma_{\nu_\mu} + (\phi_{\bar{\nu}_\mu} P_{\mu\mu} + \phi_{\bar{\nu}_e} P_{e\mu}) \sigma_{\bar{\nu}_\mu}] \cdot N_{Ar} \cdot \Delta E \cdot \Delta\theta_z \cdot \Delta t$$

Interaction cross section
with Ar40

Preliminary study

- Focus on **atmospheric neutrinos**
- The addition of a neutrino in the model changes oscillation probabilities
→ produce **appearance or disappearance** of neutrino flavours depending on energy/angle
- Compare expected number of events assuming Standard Model and “3+1” model with one sterile neutrino
- Example of calculation done for **numu** (everything is **also done for nue**)
- Focus on **CC interactions and NO** at first

$$N_{exp, ev} = [(\phi_{\nu_\mu} P_{\mu\mu} + \phi_{\nu_e} P_{e\mu})\sigma_{\nu_\mu} + (\phi_{\bar{\nu}_\mu} P_{\bar{\mu}\mu} + \phi_{\bar{\nu}_e} P_{\bar{e}\mu})\sigma_{\bar{\nu}_\mu}] \boxed{N_{Ar}} \boxed{\Delta E \cdot \Delta\theta_z} \boxed{\Delta t}$$


$$\boxed{N_{Ar} = \frac{m_{det}}{m_{Ar40}}}$$

Number of target atoms

$\boxed{\Delta t}$: time of data
taking in seconds


$\boxed{\Delta E, \Delta\theta_z}$: width of the bins used for the
histograms (binning is based on
the flux histograms)

Probability computation



- The **oscillation probabilities** are calculated using the OscProb code available on GitHub: <https://github.com/joaoabcoelho/OscProb>
- Here show survival probability of ν_μ for SM and “3+1” model assuming NO
- Put sterile CP violating phases to 0
- Probability averaged on each energy bin
- Active mixing angles from **nuFit v5.2** with Kamiokande atmospheric data (<http://arxiv.org/abs/2007.14792>, <http://www.nu-fit.org>)

Sensitivity estimator



Use a quick estimator of the sensitivity as a **preliminary probe** to compare with the results from a KM3NeT/ORCA paper (<https://arxiv.org/pdf/2107.00344.pdf>)

→ **the results seem to be coherent, encouraging**


Reminder : detector effects not included yet so we can compare without too many issues

$$S_\sigma = \frac{(N_{sterile} - N_{standard}) | N_{sterile} - N_{standard} |}{N_{sterile}}$$

similar to gaussian signed chi2

Sensitivity estimator


DUNE Preliminary



Tarak Thakore: "Sensitivity to Neutrino Mass Ordering and sterile neutrino model parameters with atmospheric neutrinos measurements at DUNE"

- Taking S_{σ} as an intermediate step towards a sensitivity estimation
- Goal: compare with previous simple DUNE study
→ <https://indico.kps.or.kr/event/30/contributions/503/>
- Need to fix values of other mixing parameters: $\sin^2\theta_{14}$ and $\sin^2\theta_{34}$ → chose minimum value for simplicity
→ **10^{-3} for both $\sin^2\theta_{14}$ and $\sin^2\theta_{34}$**
- Effect of variations under study
- See if we get something somewhat close to this distribution with our S_{σ} estimator summed over all the bins
- Also change exposure to **400kt . year** → more realistic value


Sensitivity estimator



See semblance of the same shape so it's encouraging → **contour plots in the future** to confirm, as this is a rough estimator

EarthProbe

- Code co-written by Joao : **EarthProbe** (<https://gitlab.in2p3.fr/apc-tomography/earthprobe>)
- Initially designed to use atmospheric neutrino oscillations to study Earth's composition
- (My) **current work**: extend the code to do neutrino oscillation studies and **sterile neutrino search**
- EarthProbe includes **OscProb** and manages :
 - **flux and xsec** histograms
 - **detector** specificities
 - number of events computation
 - reconstruction **efficiency**
 - oscillation probabilities
 - analysis → likelihood computations and parameter fitting using **Minuit2**



Next steps

- Finalize code adaptations of EarthProbe in order to obtain **sensitivity plots**
- Produce the same preliminary results as before as sanity check
- Understand **parameter fitting** with EarthProbe
- Produce sensitivity plots
- Add detector **reconstruction effects** and DUNE **systematics**
- After this, move on to the study with **atmospherics simulation and compatibility with Mach3**
 - will need to adapt the code




Thanks a lot for your attention !

Standard mixing parameters

NuFIT 5.2 (2022)


	Normal Ordering (best fit)		Inverted Ordering ($\Delta\chi^2 = 2.3$)		
	bfp $\pm 1\sigma$	3σ range	bfp $\pm 1\sigma$	3σ range	
	without SK atmospheric data				
$\sin^2 \theta_{12}$	$0.303^{+0.012}_{-0.011}$	0.270 \rightarrow 0.341	$0.303^{+0.012}_{-0.011}$	0.270 \rightarrow 0.341	
$\theta_{12}/^\circ$	$33.41^{+0.75}_{-0.72}$	31.31 \rightarrow 35.74	$33.41^{+0.75}_{-0.72}$	31.31 \rightarrow 35.74	
$\sin^2 \theta_{23}$	$0.572^{+0.018}_{-0.023}$	0.406 \rightarrow 0.620	$0.578^{+0.016}_{-0.021}$	0.412 \rightarrow 0.623	
$\theta_{23}/^\circ$	$49.1^{+1.0}_{-1.3}$	39.6 \rightarrow 51.9	$49.5^{+0.9}_{-1.2}$	39.9 \rightarrow 52.1	
$\sin^2 \theta_{13}$	$0.02203^{+0.00056}_{-0.00059}$	0.02029 \rightarrow 0.02391	$0.02219^{+0.00060}_{-0.00057}$	0.02047 \rightarrow 0.02396	
$\theta_{13}/^\circ$	$8.54^{+0.11}_{-0.12}$	8.19 \rightarrow 8.89	$8.57^{+0.12}_{-0.11}$	8.23 \rightarrow 8.90	
$\delta_{CP}/^\circ$	197^{+42}_{-25}	108 \rightarrow 404	286^{+27}_{-32}	192 \rightarrow 360	
$\frac{\Delta m_{21}^2}{10^{-5} \text{ eV}^2}$	$7.41^{+0.21}_{-0.20}$	6.82 \rightarrow 8.03	$7.41^{+0.21}_{-0.20}$	6.82 \rightarrow 8.03	
$\frac{\Delta m_{3\ell}^2}{10^{-3} \text{ eV}^2}$	$+2.511^{+0.028}_{-0.027}$	$+2.428 \rightarrow +2.597$	$-2.498^{+0.032}_{-0.025}$	$-2.581 \rightarrow -2.408$	
with SK atmospheric data	Normal Ordering (best fit)		Inverted Ordering ($\Delta\chi^2 = 6.4$)		
	bfp $\pm 1\sigma$	3σ range	bfp $\pm 1\sigma$	3σ range	
	$\sin^2 \theta_{12}$	$0.303^{+0.012}_{-0.012}$	0.270 \rightarrow 0.341	$0.303^{+0.012}_{-0.011}$	0.270 \rightarrow 0.341
	$\theta_{12}/^\circ$	$33.41^{+0.75}_{-0.72}$	31.31 \rightarrow 35.74	$33.41^{+0.75}_{-0.72}$	31.31 \rightarrow 35.74
	$\sin^2 \theta_{23}$	$0.451^{+0.019}_{-0.016}$	0.408 \rightarrow 0.603	$0.569^{+0.016}_{-0.021}$	0.412 \rightarrow 0.613
	$\theta_{23}/^\circ$	$42.2^{+1.1}_{-0.9}$	39.7 \rightarrow 51.0	$49.0^{+1.0}_{-1.2}$	39.9 \rightarrow 51.5
	$\sin^2 \theta_{13}$	$0.02225^{+0.00056}_{-0.00059}$	0.02052 \rightarrow 0.02398	$0.02223^{+0.00058}_{-0.00058}$	0.02048 \rightarrow 0.02416
	$\theta_{13}/^\circ$	$8.58^{+0.11}_{-0.11}$	8.23 \rightarrow 8.91	$8.57^{+0.11}_{-0.11}$	8.23 \rightarrow 8.94
	$\delta_{CP}/^\circ$	232^{+36}_{-26}	144 \rightarrow 350	276^{+22}_{-29}	194 \rightarrow 344
	$\frac{\Delta m_{21}^2}{10^{-5} \text{ eV}^2}$	$7.41^{+0.21}_{-0.20}$	6.82 \rightarrow 8.03	$7.41^{+0.21}_{-0.20}$	6.82 \rightarrow 8.03
$\frac{\Delta m_{3\ell}^2}{10^{-3} \text{ eV}^2}$	$+2.507^{+0.026}_{-0.027}$	$+2.427 \rightarrow +2.590$	$-2.486^{+0.025}_{-0.028}$	$-2.570 \rightarrow -2.406$	

Sensitivity estimator




- We obtain a similar behaviour → excess around 20 GeV and deficit above and below but unclear due to low statistics
- Possibility to explore lower energy ranges
- The S parameter is intended for higher statistics → **will be updated to log-likelihood**

Sensitivity estimator



- Increasing the size of the detector to 1Mt to allow a better cross-check of result
→ shows **similar behaviour** as KM3NeT/ORCA plot
- Higher frequency effect has been investigated and will likely be diffused when detector effects are included

Expected number of events



Can calculate **number of expected events** and do the difference between 4-neutrino and 3-neutrino models for ν_μ and $\bar{\nu}_e$ (also anti- ν) depending on **true angle and true energy** of incoming neutrinos