



ID de Contribution: 63

Type: Non spécifié

## The recoil distance Doppler-shift technique: a valuable method for state-of-the-art nuclear structure studies

*vendredi 8 novembre 2024 09:35 (20 minutes)*

Absolute transition strengths between excited states yield fundamental information on nuclear structure. These observables can be determined from level lifetimes. The recoil distance Doppler-shift (RDDS) technique employing so-called plunger devices provides a valuable method for the determination of lifetimes in the picosecond range and has been in the focus of our Cologne group since many years.

In this talk, new results from RDDS measurements of our group especially in tellurium isotopes close to neutron midshell and neutron-deficient nuclei in the  $A=170$  region are discussed. Special emphasis is paid to the evolution of collectivity including hints for shape coexistence. In the tellurium isotopes, among other results, also lifetimes of excited  $0^+$  states were measured allowing the calculation of  $\rho^2(E0)$  strengths that are indicators of the mixing of coexisting structures. In the  $A=170$  region, signatures for a structural change with decreasing neutron number were found that cannot be reproduced with nuclear models so far. Further, we will discuss recent experiments with the RDDS technique on neutron-rich nuclei in the  $A=40-50$  region with respect to the evolution and a possible weakening of the  $N=28$  shell closure.

We will also present our latest developments of the RDDS technique with respect to the application with very different kinematic conditions and reactions techniques, e.g., multinucleon transfer, fusion-evaporation reactions and direct reactions, also employing radioactive beams, and the required detection techniques to identify the reaction products. The related recoil velocities of the reaction products range from few percent of the speed of light up to the relativistic regime with  $v/c \approx 60\%$ . For this purpose, our group developed sophisticated plunger devices adopted to the respective spectrometers and beam geometries.

Funded by the German Research Foundation (DFG), grant No. FR 3276/3-1

**Auteur principal:** FRANSEN, Christoph (IKP, Köln, Germany)

**Co-auteurs:** ESMAYLZADEH, Arwin (IKP, University of Cologne, Germany); LAKENBRINK, Casper-David (IKP, University of Cologne, Germany); MÜLLER-GATERMANN, Claus (Argonne National Laboratory, Illinois, USA); DUNKEL, Felix (IKP, University of Cologne, Germany); VON SPEE, Franziskus (IKP, University of Cologne, Germany); JOLIE, Jan (IKP, University of Cologne, Germany); NOVAK, Richard (IKP, University of Cologne, Germany)

**Orateur:** FRANSEN, Christoph (IKP, Köln, Germany)

**Classification de Session:** Session 15