

The 2018 Fani Maoré submarine eruption offshore Mayotte (Indian Ocean): Insights on fine-scale morphologies from ROV high-resolution bathymetry and optical observations

Valentine Puzenat¹, Nathalie Feuillet¹, Javier Escartín², Jean-Christophe Komorowski¹, Christine Deplus¹, Cécile Cathalot³, Benoit Loubrieu³, Delphine Pierre³, Mathilde Pitel-Roudaut³, Emmanuel Rinnert³ and Carla Scalabrin³

¹ Université Paris Cité, Institut de Physique du Globe de Paris, CNRS, F-75005 Paris, France, ² Laboratoire de Géologie - CNRS, UMR 8538, École Normale Supérieure, PSL University, Paris, France, ³ Geo-Ocean, Univ Brest, CNRS, Ifremer, UMR6538, F-29280 Plouzane, France

What?

Using lava flow morphology as a proxy to understand the dynamics of submarine eruptions

Where?

- Fani Maoré submarine volcano
- Offshore Mayotte Island (Indian Ocean)
- Largest submarine eruption ever recorded

How?

- High-resolution bathymetry data (~2 m)
- Optical imagery from submarines
- 3D reconstructions

Why?

- Document the fine-scale morphologies and eruptive processes associated with the eruption
- Constrain the precise chronology and spatial variability of lava emplacement
- Ultimate goal: better understanding of deep-sea volcanic processes

So what?

See you in the entrance hall 😊

