

*Insights from*  
*Galaxies at long distances*

# Questions...

$$\mathcal{P}(* | \text{map}) = ?$$


\*: "(new) physics"

$$\mathcal{P}(* | \delta) = ?$$

$$\delta(\mathbf{x}) \equiv \frac{\rho(\mathbf{x})}{\bar{\rho}} - 1$$

# Cosmic maps have the answer?

Coarse-graining



$$\delta(\mathbf{k}) = \delta_\ell(\mathbf{k}) + \delta_s(\mathbf{k})$$

$$\delta_\ell(\mathbf{k}) = \delta(|\mathbf{k}| < \Lambda_s^{-1})$$

$$\delta_\ell(\mathbf{k}) \lesssim \mathcal{O}(1) \text{ for } k \lesssim k_{\text{nl}}$$

$$(k < \Lambda_s^{-1} < k_{\text{nl}})$$

In this talk


$$\mathcal{P}(* | \delta) \rightarrow \mathcal{P}(* | \delta_\ell) = ?$$

*What can we learn*

*looking at galaxies from afar?*

# Cosmic maps have the answer?

Coarse-graining



$$\delta(\mathbf{k}) = \delta_\ell(\mathbf{k}) + \delta_s(\mathbf{k})$$

$$\delta_\ell(\mathbf{k}) = \delta(|\mathbf{k}| < \Lambda_s^{-1})$$

$$\delta_\ell(\mathbf{k}) \lesssim \mathcal{O}(1) \text{ for } k \lesssim k_{\text{nl}}$$

$$(k < \Lambda_s^{-1} < k_{\text{nl}})$$

In this talk

$$\mathcal{P}(* | \delta) \rightarrow \mathcal{P}(* | \delta_\ell) = ?$$

*What can we learn*

*looking at galaxies from afar?*

# Plan

1. *Galaxies from afar*: an effective description at long distances
2. Cosmology with galaxies from afar: a *tour d'horizon*

# *Looking at galaxies from afar*

An effective description at long wavelength

# Effective Field Theory of Large-Scale Structure

Zel'dovich 70's, Peebles 80's, Bouchet, Bernardeau, Scoccimarro, ... 90's-00's, Baumann, Carrasco, Hertzberg, Nicolis, Pajer, Senatore, Zaldarriaga, ... 10-13

Looking from afar, we want to know fields describing matter, baryons, galaxies, etc., e.g.  $\delta$ ,  $\delta_b$ ,  $\delta_g$ ,  $v$ , ...

## Ingredients

- Dark matter: *Continuity and Euler equations* (coarse-grained)
- Gravity: *Poisson equation*  $\partial^2 \Phi \sim \delta$
- Symmetries: *Galilean invariance*  $\mathbf{x} \rightarrow \mathbf{x} + \mathbf{n}$ ,  $\mathbf{v} \rightarrow \mathbf{v} + \partial_t \mathbf{n}$

Weinberg 03, Kehagias, Riotto, Peloso, Pietroni, Creminelli, Gleyzes, Noreña, Simonović, Vernizzi 13

## Receipe

- Solve dark matter equations perturbatively
  - $\delta = \delta_1 + \delta_2 + \dots$
- For unknowns, write down all terms allowed by the symmetries with free Wilson coefficients
  - $\delta_g = b_1 \delta_1 + b_2 \delta_2 + \dots$
  - *Euler*  $\sim \partial_j \tau^{ij} / k_{nl}^2$ ,  $\tau^{ij} = \delta^{ij} c_1 \delta_1 + \dots$
- For UV-sensitive operators, add counterterms

# Galaxies fluid expansion

McDonald 06-09, Angulo, Assassi, Baumann, Fasiello, Fujita, Green, Mirbabayi, Schmidt, Senatore, Vlah, Zaldarriaga, ... 14-16

$$\delta_g(\mathbf{x}, t) = \int^t dt' f\left(\partial_i \partial_j \Phi(\mathbf{x}, t'), \partial_i v_j(\mathbf{x}, t'), \partial_i / k_M, \epsilon_{ij}(\mathbf{x}, t'), \kappa_\star(t, t')\right)$$

- fluctuations (equivalence principle)  $\partial_i \partial_j \Phi, \partial_i v_j$
- gradients (spatial extension)  $\partial_i / k_M$
- stochasticity  $\epsilon, \dots$
- fluid expansion  $\mathbf{x} \rightarrow \mathbf{x}_{\text{fl}}(\mathbf{x}, t, t')$
- time responses  $\int^t dt' \kappa_\star(t, t') D(t'), \dots$

w/ D'Amico, Donath, Lewandowski, Senatore 22a

Up to 4<sup>th</sup> order, equivalent to *local-in-time* expansion

$$\delta_g(\mathbf{x}, t) = f\left(\partial_i \partial_j \Phi(\mathbf{x}, t), \partial_i v_j(\mathbf{x}, t), \partial_i / k_M, \epsilon_{ij}(\mathbf{x}, t), \kappa_\star(t)\right)$$

End of the day:  $\delta_g(\mathbf{x}, t) = \sum_\alpha b_\alpha(t) \mathcal{O}^\alpha[\delta_1(\mathbf{x}, t)]$



# Redshift space

Lewandowski, Perko, Senatore, Zaldarriaga, ... 14-16, ...

Change of coordinates  $\mathbf{x} \rightarrow \mathbf{x} + (\mathbf{v}_{\mathcal{H}} \cdot \hat{z})\hat{z}$  ( $\mathbf{v}_{\mathcal{H}} \equiv \mathbf{v}/\mathcal{H}$ )

$$\delta \rightarrow \delta + \hat{z}^i \hat{z}^j \partial_i ((1 + \delta) v_{\mathcal{H}}^j) + \frac{1}{2} \hat{z}^i \hat{z}^j \hat{z}^k \hat{z}^l \partial_i \partial_j ((1 + \delta) v_{\mathcal{H}}^k v_{\mathcal{H}}^l) + \dots$$

Contact term renormalization, e.g.  $\delta v_{\mathcal{H}}^i \supset c_{\text{rs}} \partial_i \delta_1 / k_{\text{rs}}^2$

# Summary, part 1

## **EFTofLSS**

- organizes the fields expansion into fluctuations, gradients, etc.
- captures unknowns into free coefficients
- includes redshift-space distortions, IR displacements, baryons, massive neutrinos

## **Pros**

- robust (parametric control)
- flexible (analytic)

## **Cons**

- mildly nonlinear scales only
- many parameters

# *Tour d'horizon*

Cosmology with galaxies at long distances

# $\Lambda$ CDM

Given an observation  $\hat{\delta}_g$ , what cosmology  $\Omega$  can we learn, i.e.  $\mathcal{P}(\Omega|\hat{\delta}_g)$ ?

$$\delta_g \sim b_1(t)\delta_1(\mathbf{x}, t) + b_{\text{nl}}(t)\delta_{\text{nl}}(\mathbf{x}, t) + f(t)b_{\text{rs}}(t)\delta_{\text{rs}}(\mathbf{x}, t) + \dots$$

- linear perturbations  $\delta_1(\Omega)$  + nonlinear perturbations  $\delta_{\text{nl}}[\delta_1(\Omega)]$
- redshift-space distortions  $f(\Omega)$ ,  $\delta_{\text{rs}}[\delta_1(\Omega)]$

## In practice


*PyBird, CLASS-PT, Velocileptor, FAST-PT, CLASS-OneLoop, ...*

- Compute  $N$ -point functions, e.g.  $\langle \delta_g(\mathbf{x}_1)\delta_g(\mathbf{x}_2) \rangle$  ,  $\langle \delta_g(\mathbf{x}_1)\delta_g(\mathbf{x}_2)\delta_g(\mathbf{x}_3) \rangle$  ...
- Explore likelihood  $\mathcal{L}(\langle \hat{\delta}_g \hat{\delta}_g \rangle, \langle \hat{\delta}_g \hat{\delta}_g \hat{\delta}_g \rangle | \Omega)$

> <https://github.com/pierrexzyz/pybird> <

# $\Lambda$ CDM from galaxy surveys

w/ D'Amico, Donath, Lewandowski, Senatore 22b



# Dark energy and modified gravity

Given an observation  $\hat{\delta}_g$ , what cosmology  $\Omega$  can we learn, i.e.  $\mathcal{P}(\Omega|\hat{\delta}_g)$ ?

$$\delta_g \sim b_1 D(t) \delta_1 + b_{\text{nl}} g_{\text{nl}}(t) \delta_{\text{nl}} + f(t) b_{\text{rs}} g_{\text{rs}}(t) \delta_{\text{rs}} + \dots$$

D'Amico, Donath, Fujita, Lewandowski, Marinucci, Pietroni, Piga, Senatore, Taule, Vernizzi, Vlah, Zhang, ... 15-24

- Generalized nonlinear time evolution  $g_{\text{nl}}(\Omega)$ ,  $g_{\text{rs}}(\Omega)$
- (EFTofDE)  $\partial^2 \Phi \sim \mu_1 \delta + \mu_2 \delta^2 + \dots$

# New physics (long range)

Given an observation  $\hat{\delta}_g$ , what cosmology  $\Omega$  can we learn, i.e.  $\mathcal{P}(\Omega|\hat{\delta}_g)$ ?

$$\delta_g \sim b_1 \delta_1 + b_{\text{nl}} \delta_{\text{nl}} + f b_{\text{rs}} \delta_{\text{rs}} + b_* \delta_*$$

w/ D'Amico, Lewandowski, Senatore 22

- Primordial non-Gaussianity
  - (single-field Inflation)  $f_{\text{NL}} \phi^2$
  - (multi-field Inflation) Scale-dependent bias  $b_\phi f_{\text{NL}} \partial^{-2} \delta_1$

Lewandowski 19, Bottaro, Castorina, Costa, Redigolo, Salvioni 23

- Dark long-range force, beyond Horndeski ...
  - Galilean invariance breaking terms  $\delta_* [\delta_1, *]$

# Dark matter

Given an observation  $\hat{\delta}_g$ , what cosmology  $\Omega$  can we learn, i.e.  $\mathcal{P}(\Omega|\hat{\delta}_g)$ ?

$$\delta_g \sim b_1 \delta_1 + b_{\text{nl}} \delta_{\text{nl}} + f b_{\text{rs}} \delta_{\text{rs}}$$

- Speed-of-sound  $c_s^2(t)$ :  $\delta \supset c_s^2 \partial^2 \delta_1 / k_{\text{nl}}^2$ 
  - cold or not cold? comparison with simulations
  - *Issue*: degenerate with  $\partial^2 \delta_1 / k_{\text{M}}^2$ ,  $\partial^2 \delta_1 / k_{\text{rs}}^2$ , ...

Kousha, Hooshangi, Abolhasani 23

w/ D'Amico, Donath, Lewandowski, Senatore 22a

## Non-locality?

- Clean window onto dark matter from galaxies *in redshift space*
  - via *non-locally-contributing* (non degenerate) counterterms at  $2^{\text{nd}}$  order ( $4^{\text{th}}$  order in fluctuations)



# One-loop cosmology: today and tomorrow


w/ D'Amico, Lewandowski, Senatore 22

w/ Simon & Poulin 22

Bragança, Donath, Senatore, Zheng 23

w/ Spaar 23

w/ Racco & Zheng (to appear)



	$\sigma(f_{\text{NL}}^{\text{or}})$	$\sigma(f_{\text{NL}}^{\text{eq}})$
Planck	24	47
BOSS $P_\ell + B_0$	72	293
Future $P_\ell + B_0$	4	16

	$\sigma(w)$ [%]
Planck + BOSS BAO + ext	2.6
Planck + BOSS $P_\ell + B_0$ + ext	1.9
BOSS $P_\ell + \text{eBOSS } P_\ell + \text{ext}$	4.1
Future $P_\ell + B_0$	0.5

	$\sigma(\sum m_\nu)$
Planck	0.24 ( $< 2\sigma$ )
Planck BOSS $P_\ell + \text{eBOSS } P_\ell + \text{ext}$	0.09 ( $< 2\sigma$ )
BOSS $P_\ell + \text{eBOSS } P_\ell + \text{ext}$	0.26 ( $< 2\sigma$ )

# Final words

- **EFTofLSS**

- *Robust*: galaxies meet precision cosmology
- *Flexible*: great to include new physics !
- *Caveat*: mainly long range

- **Cosmology**: a large menu



- Next-decade BIG targets:  $\sum m_\nu$  detection,  $\sigma(w) < 1\%$ ,  $\sigma(f_{\text{NL}}) \sim \mathcal{O}(1 - 10)$ .
- Modified gravity, Inflation, dark matter, ...
  - Systematic inclusion (on going)
  - EFT  $\leftrightarrow$  UV dictionary?

- **Roads ahead**

- No *loops*, no chocolate: push theory, we are not yet here
- No *measures*, no chocolate: new ways to look at data (forward model, estimators)?

# *Supplements*

# The higher, the better



$$N_{\text{modes}}(\text{CMB}) \sim (\ell_{\text{max}})^2$$

$$N_{\text{modes}}^{\diamond}(\text{LSS}) \sim V_{\text{survey}} (k_{\text{max}}^{\diamond})^3 / (6\pi^2)$$

$\diamond = \text{yesterday, today, tomorrow}$

**One loop: not yet optimal!**

# Large IR displacements

Matsubara 07-08, Aviles, Baldauf, Blah, Garny, Ivanov, Lewandowski, Mirbabayi, Porto, Senatore, Seljak, Sibiryakov, Simonovic, Vlah, White, Zaldarriaga, ... 13-16

Lagrangian picture  $\mathbf{x}_L(t) = \mathbf{x}(t) + \psi(\mathbf{x}, t)$ ,  $\psi = \psi_1 + \psi_2 + \dots$

$$\psi_{\ell_{\text{BAO}}} \equiv \psi (|\mathbf{k}| < 2\pi/\ell_{\text{BAO}}) \sim \mathcal{O}(1)! \quad (\ell_{\text{BAO}} \sim 100 \text{ Mpc}/h)$$

$\psi_{\ell_{\text{BAO}}} \simeq \psi_1$ : keep **large** displacements  $\psi_1$ , expand in  $\psi_2, \dots$

$$\delta_g(\mathbf{x}_L) = b_1 \delta_1(\mathbf{x}_L) + b_2 \delta_2(\mathbf{x}_L) + \dots$$

$$= b_1 \delta_1(\mathbf{x} + \psi_1) + b_1 \partial_i ((1 + \delta_1(\mathbf{x})) \psi_2^i) + b_2 \delta_2(\mathbf{x} + \psi_1) + \dots$$

$$= b_1 \delta_1(\mathbf{x} + \psi_1) + b'_2 \delta_2(\mathbf{x} + \psi_1) + \dots$$

- $\psi_1^i = \frac{\partial_i}{\partial^2} \delta_1$
- $\psi_2^i = \frac{\partial_i}{\partial^2} \delta_2$

$$\delta_g(\mathbf{x}) \rightarrow \delta_g(\mathbf{x} + \psi_1)$$

