

COSMOLOGICAL RESULTS FROM THE FIRST eROSITA ALL SKY SURVEY

E. ARTIS on behalf of the eROSITA Clusters & Cosmology group


19 March 2024



Galaxy clusters:


- 85 % dark matter
- 13 % hot gas
- 2% galaxies

- Scale: a few Mpc(s)
- Hundreds to thousands of galaxies
- $\sim 10^{14} h^{-1}M_{\odot}$ and beyond




300 Mpc

Credits : TNG simulations




CLUSTER ABUNDANCE AS A COSMOLOGICAL PROBE




CLUSTER ABUNDANCE AS A COSMOLOGICAL PROBE

Abdalla et al., 2022




CLUSTER ABUNDANCE AS A COSMOLOGICAL PROBE

Abdalla et al., 2022



CLUSTER ABUNDANCE AS A COSMOLOGICAL PROBE

EA et al., 2024





Redshift and scale sensitivity of the different redshift probes, adapted from [Preston et al., 2023](#)

➤ Cluster abundance probe redshift and scale ranges complementary to other probes

➤ If discrepancies are to arise, it is important that they are tested in these different regimes

eROSITA X-RAY TELESCOPE ON SRG



- **eROSITA = extended ROentgen Survey** with an Imaging Telescope Array
- **X-ray all-sky survey** in **4** years (**8x6** month surveys)
- Expected lifetime **> 7** years
- **Sensitive to 0.2 - 10 keV band**
- **20-30 times more sensitive than ROSAT**
- **Grasp; FOV*Effective Area @1keV:**
 - **5 × XMM-Newton**
 - **100 × Chandra ACIS**



Credit : E. Bulbul

eROSITA X-RAY TELESCOPE ON SRG



- **eROSITA = extended ROentgen Survey** with an Imaging Telescope Array
- **X-ray all-sky survey** in **4** years (**8x6** month surveys)
- Expected lifetime **> 7** years
- **Sensitive to 0.2 - 10 keV band**
- **20-30 times more sensitive than ROSAT**
- **Grasp; FOV*Effective Area @1keV:**
 - **5 × XMM-Newton**
 - **100 × Chandra ACIS**



Credit : E. Bulbul


eROSITA X-RAY TELESCOPE ON SRG

- **eROSITA = extended ROentgen Survey** with an Imaging Telescope Array
- **X-ray all-sky survey** in **4** years (**8x6** month surveys)
- Expected lifetime **> 7** years
- **Sensitive to 0.2 - 10 keV band**
- **20-30 times more sensitive than ROSAT**
- **Grasp; FOV*Effective Area @1keV:**
 - **5 × XMM-Newton**
 - **100 × Chandra ACIS**




Credit : E. Bulbul

eROSITA SO FAR



Credit : E. Bulbul

eROSITA SO FAR




Cosmological results
from eRASS1 only

Credit : E. Bulbul


THE eRASS1 CLUSTER CATALOG

Bulbul et al., 2024



Credit : E. Bulbul

OPTICAL FOLLOW-UP WITH DESI LEGACY DR9 & 10




LS *grz* image of a cluster overlaid with eRASS1 X-ray contours

12,247 clusters with redshift measurement
86% purity

- Clean the catalog
- Get the photometric redshifts
- Get an additional mass proxy



Richness vs velocity dispersion from a subsample of the catalog

Kluge et al., 2024





THE eRASS1 CLUSTER CATALOG

Bulbul et al., 2024



X-RAY OBSERVABLE


C_R fitted using MBproj2D (Sanders et al., 2018)



Count rate to mass relation from the cluster catalog presented in Bulbul et al., 2024

THE eRASS1 CLUSTER SELECTION FUNCTION

Clerc et al., 2024




➤ Depends on the count rates, exposure times, hydrogen column density and background

➤ Quantity fitted on eRASS1 simulations


THE eRASS1 CLUSTER SELECTION FUNCTION

Clerc et al., 2024



The model is tested with eFEDS as a reference catalog


THE eRASS1 CLUSTER CATALOG



- 5,263 cluster candidates
- Total effective area: 12.8k deg²
- 95% purity

Ghirardini, Bulbul, EA et al., 2024

THE eRASS1 CLUSTER CATALOG



- 5,263 cluster candidates
- Total effective area: 12.8k deg²
- 95% purity

Largest ICM selected cosmology sample to date !

WEAK LENSING MASS CALIBRATION

Calibrate the $C_R - M$ and $\lambda - M$ relations

➤ **Dark energy Survey (DES)** Grandis et al., 2024


- 2201 tangential shears

➤ **Kilo Degree Survey (KIDS)** Kleinebreil et al., 2024


- 96 tangential shears

➤ **Hyper Supreme Cam Survey (HSC)** Chiu et al., 2024

- 236 tangential shears




Cosmological results from the first eROSITA All Sky Survey (eRASS1)



Grandis et al., 2024



Each individual shear profile is directly integrated in the cluster abundance likelihood




Grandis et al., 2024



Each individual shear profile is directly integrated in the cluster abundance likelihood


Grandis et al., 2024



Each individual shear profile is directly integrated in the cluster abundance likelihood


MIXTURE MODEL: REMOVING THE CONTAMINATION FRACTION

Ghirardini, Bulbul, EA et al., 2024



We account for AGN contamination and noise fluctuation. The fraction of these objects is fitted in the global likelihood

OVERVIEW : eRASS1 CLUSTER ABUNDANCE PIPELINE



Parameter	Units	Description	Prior
• Cosmology			
Ω_m	-	Mean matter density at present time	$\mathcal{U}(0.05, 0.95)$
$\log_{10} A_s$	-	Amplitude of the primordial power spectrum	$\mathcal{U}(-10, -8)$
H_0	$\frac{\text{km}}{\text{s Mpc}}$	Hubble expansion rate at present time	$\mathcal{N}(67.77, 0.6)$
Ω_b	-	Mean baryon density at present time	$\mathcal{U}(0.046, 0.052)$
n_s	-	Spectra index of the primordial power spectrum	$\mathcal{U}(0.92, 1.0)$
w_0	-	Dark energy equation of state. Fixed to -1 in Λ CDM	$\mathcal{U}(-2.5, -0.33)$
$\sum m_\nu$	eV	Summed neutrino masses. Fixed to 0 eV in Λ CDM	$\mathcal{U}(0, 1)$
• X-ray scaling relation			
A_X	-	Normalization of the $M - C_R$ scaling relation	$\mathcal{U}(0.01, 3)$
B_X	-	Mass slope of the $M - C_R$ scaling relation	$\mathcal{U}(0.1, 5)$
D_X	-	Luminosity distance evolution of the $M - C_R$ scaling relation	Fixed to -2
E_X	-	Scale factor evolution of the $M - C_R$ scaling relation	Fixed to 2
F_X	-	Redshift evolution of the mass slope of the $M - C_R$ scaling relation	$\mathcal{U}(-5, 5)$
G_X	-	Redshift evolution of the normalization of the $M - C_R$ scaling relation	$\mathcal{U}(-5, 5)$
σ_X	-	Intrinsic scatter of the $M - C_R$ scaling relation	$\mathcal{U}(0.05, 2)$
• Weak lensing mass calibration			
A_{WL}	-	Scatter in the weak lensing bias from the first principal component	$\mathcal{N}(0, 1)$
B_{WL}	-	Scatter in the weak lensing bias from the second principal component	$\mathcal{N}(0, 1)$
C_{WL}	-	Standardize mass slope of the weak lensing bias	$\mathcal{N}(0, 1)$
D_{WL}	-	Redshift dependent intrinsic scatter in the weak lensing bias	$\mathcal{N}(0, 1)$
$\rho_{M_{\text{WL}}, C_R}$	-	Intrinsic correlation between weak lensing mass and count rate	$\mathcal{U}(-0.9, 0.9)$
• Richness mass calibration			
$\log A_\lambda$	-	Normalization of the $M - \lambda$ scaling relation	$\mathcal{U}(1.4, 6)$
B_λ	-	Mass slope of the $M - \lambda$ scaling relation	$\mathcal{U}(0, 2)$
C_λ	-	Redshift evolution of the normalization of the $M - \lambda$ scaling relation	$\mathcal{U}(-2, 2)$
D_λ	-	Redshift evolution of the mass slope of the $M - \lambda$ scaling relation	$\mathcal{U}(-2, 2)$
σ_λ	-	Intrinsic scatter of the $M - \lambda$ scaling relation	$\mathcal{U}(0.05, 2)$
ρ_{λ, C_R}	-	Intrinsic correlation between richness and count rate	$\mathcal{U}(-0.9, 0.9)$
• Contamination modeling			
f_{AGN}	-	Fraction of AGN contaminants in the extended source sample	$\mathcal{U}(0, 0.1)$
f_{RS}	-	Fraction of RS contaminants in the extended source sample	$\mathcal{U}(0, 0.15)$
• Redshift uncertainty			
σ_z	-	Relative error on the measured redshift	$\mathcal{TN}(0.0050, 0.0011, 0, 1)$
b_z	-	Systematic bias in our redshift estimate	$\mathcal{N}(1.005, 0.037)$
c_z	-	Fraction of objects for which we measure a shifted redshift	$\mathcal{TN}(0.0013, 0.0010, 0, 1)$
$c_{\text{shift}, z}$	-	Amount of redshift shift for c_z fraction of objects	$\mathcal{U}(-0.1, 0.1)$

Parameter	Units	Description	Prior
• Cosmology			
Ω_m	-	Mean matter density at present time	$\mathcal{U}(0.05, 0.95)$
$\log_{10} A_s$	-	Amplitude of the primordial power spectrum	$\mathcal{U}(-10, -8)$
H_0	$\frac{\text{km}}{\text{s Mpc}}$	Hubble expansion rate at present time	$\mathcal{N}(67.77, 0.6)$
Ω_b	-	Mean baryon density at present time	$\mathcal{U}(0.046, 0.052)$
n_s	-	Spectra index of the primordial power spectrum	$\mathcal{U}(0.92, 1.0)$
w_0	-	Dark energy equation of state. Fixed to -1 in Λ CDM	$\mathcal{U}(-2.5, -0.33)$
$\sum m_\nu$	eV	Summed neutrino masses. Fixed to 0 eV in Λ CDM	$\mathcal{U}(0, 1)$
• X-ray scaling relation			
A_X	-	Normalization of the $M - C_R$ scaling relation	$\mathcal{U}(0.01, 3)$
B_X	-	Mass slope of the $M - C_R$ scaling relation	$\mathcal{U}(0.1, 5)$
D_X	-	Luminosity distance evolution of the $M - C_R$ scaling relation	Fixed to -2
E_X	-	Scale factor evolution of the $M - C_R$ scaling relation	Fixed to 2
F_X	-	Redshift evolution of the mass slope of the $M - C_R$ scaling relation	$\mathcal{U}(-5, 5)$
G_X	-	Redshift evolution of the normalization of the $M - C_R$ scaling relation	$\mathcal{U}(-5, 5)$
σ_X	-	Intrinsic scatter of the $M - C_R$ scaling relation	$\mathcal{U}(0.05, 2)$
• Weak lensing mass calibration			
A_{WL}	-	Scatter in the weak lensing bias from the first principal component	$\mathcal{N}(0, 1)$
B_{WL}	-	Scatter in the weak lensing bias from the second principal component	$\mathcal{N}(0, 1)$
C_{WL}	-	Standardize mass slope of the weak lensing bias	$\mathcal{N}(0, 1)$
D_{WL}	-	Redshift dependent intrinsic scatter in the weak lensing bias	$\mathcal{N}(0, 1)$
$\rho_{M_{\text{WL}}, C_R}$	-	Intrinsic correlation between weak lensing mass and count rate	$\mathcal{U}(-0.9, 0.9)$
• Richness mass calibration			
$\log A_\lambda$	-	Normalization of the $M - \lambda$ scaling relation	$\mathcal{U}(1.4, 6)$
B_λ	-	Mass slope of the $M - \lambda$ scaling relation	$\mathcal{U}(0, 2)$
C_λ	-	Redshift evolution of the normalization of the $M - \lambda$ scaling relation	$\mathcal{U}(-2, 2)$
D_λ	-	Redshift evolution of the mass slope of the $M - \lambda$ scaling relation	$\mathcal{U}(-2, 2)$
σ_λ	-	Intrinsic scatter of the $M - \lambda$ scaling relation	$\mathcal{U}(0.05, 2)$
ρ_{λ, C_R}	-	Intrinsic correlation between richness and count rate	$\mathcal{U}(-0.9, 0.9)$
• Contamination modeling			
f_{AGN}	-	Fraction of AGN contaminants in the extended source sample	$\mathcal{U}(0, 0.1)$
f_{RS}	-	Fraction of RS contaminants in the extended source sample	$\mathcal{U}(0, 0.15)$
• Redshift uncertainty			
σ_z	-	Relative error on the measured redshift	$\mathcal{TN}(0.0050, 0.0011, 0, 1)$
b_z	-	Systematic bias in our redshift estimate	$\mathcal{N}(1.005, 0.037)$
c_z	-	Fraction of objects for which we measure a shifted redshift	$\mathcal{TN}(0.0013, 0.0010, 0, 1)$
$c_{\text{shift}, z}$	-	Amount of redshift shift for c_z fraction of objects	$\mathcal{U}(-0.1, 0.1)$


Parameter	Units	Description	Prior
• Cosmology			
Ω_m	-	Mean matter density at present time	$\mathcal{U}(0.05, 0.95)$
$\log_{10} A_s$	-	Amplitude of the primordial power spectrum	$\mathcal{U}(-10, -8)$
H_0	$\frac{\text{km}}{\text{s Mpc}}$	Hubble expansion rate at present time	$\mathcal{N}(67.77, 0.6)$
Ω_b	-	Mean baryon density at present time	$\mathcal{U}(0.046, 0.052)$
n_s	-	Spectra index of the primordial power spectrum	$\mathcal{U}(0.92, 1.0)$
w_0	-	Dark energy equation of state. Fixed to -1 in Λ CDM	$\mathcal{U}(-2.5, -0.33)$
$\sum m_\nu$	eV	Summed neutrino masses. Fixed to 0 eV in Λ CDM	$\mathcal{U}(0, 1)$
• X-ray scaling relation			
A_X	-	Normalization of the $M - C_R$ scaling relation	$\mathcal{U}(0.01, 3)$
B_X	-	Mass slope of the $M - C_R$ scaling relation	$\mathcal{U}(0.1, 5)$
D_X	-	Luminosity distance evolution of the $M - C_R$ scaling relation	Fixed to -2
E_X	-	Scale factor evolution of the $M - C_R$ scaling relation	Fixed to 2
F_X	-	Redshift evolution of the mass slope of the $M - C_R$ scaling relation	$\mathcal{U}(-5, 5)$
G_X	-	Redshift evolution of the normalization of the $M - C_R$ scaling relation	$\mathcal{U}(-5, 5)$
σ_X	-	Intrinsic scatter of the $M - C_R$ scaling relation	$\mathcal{U}(0.05, 2)$
• Weak lensing mass calibration			
A_{WL}	-	Scatter in the weak lensing bias from the first principal component	$\mathcal{N}(0, 1)$
B_{WL}	-	Scatter in the weak lensing bias from the second principal component	$\mathcal{N}(0, 1)$
C_{WL}	-	Standardize mass slope of the weak lensing bias	$\mathcal{N}(0, 1)$
D_{WL}	-	Redshift dependent intrinsic scatter in the weak lensing bias	$\mathcal{N}(0, 1)$
$\rho_{M_{\text{WL}}, C_R}$	-	Intrinsic correlation between weak lensing mass and count rate	$\mathcal{U}(-0.9, 0.9)$
• Richness mass calibration			
$\log A_\lambda$	-	Normalization of the $M - \lambda$ scaling relation	$\mathcal{U}(1.4, 6)$
B_λ	-	Mass slope of the $M - \lambda$ scaling relation	$\mathcal{U}(0, 2)$
C_λ	-	Redshift evolution of the normalization of the $M - \lambda$ scaling relation	$\mathcal{U}(-2, 2)$
D_λ	-	Redshift evolution of the mass slope of the $M - \lambda$ scaling relation	$\mathcal{U}(-2, 2)$
σ_λ	-	Intrinsic scatter of the $M - \lambda$ scaling relation	$\mathcal{U}(0.05, 2)$
ρ_{λ, C_R}	-	Intrinsic correlation between richness and count rate	$\mathcal{U}(-0.9, 0.9)$
• Contamination modeling			
f_{AGN}	-	Fraction of AGN contaminants in the extended source sample	$\mathcal{U}(0, 0.1)$
f_{RS}	-	Fraction of RS contaminants in the extended source sample	$\mathcal{U}(0, 0.15)$
• Redshift uncertainty			
σ_z	-	Relative error on the measured redshift	$\mathcal{TN}(0.0050, 0.0011, 0, 1)$
b_z	-	Systematic bias in our redshift estimate	$\mathcal{N}(1.005, 0.037)$
c_z	-	Fraction of objects for which we measure a shifted redshift	$\mathcal{TN}(0.0013, 0.0010, 0, 1)$
$c_{\text{shift}, z}$	-	Amount of redshift shift for c_z fraction of objects	$\mathcal{U}(-0.1, 0.1)$

Parameter	Units	Description	Prior
• Cosmology			
Ω_m	-	Mean matter density at present time	$\mathcal{U}(0.05, 0.95)$
$\log_{10} A_s$	-	Amplitude of the primordial power spectrum	$\mathcal{U}(-10, -8)$
H_0	$\frac{\text{km}}{\text{s Mpc}}$	Hubble expansion rate at present time	$\mathcal{N}(67.77, 0.6)$
Ω_b	-	Mean baryon density at present time	$\mathcal{U}(0.046, 0.052)$
n_s	-	Spectra index of the primordial power spectrum	$\mathcal{U}(0.92, 1.0)$
w_0	-	Dark energy equation of state. Fixed to -1 in Λ CDM	$\mathcal{U}(-2.5, -0.33)$
$\sum m_\nu$	eV	Summed neutrino masses. Fixed to 0 eV in Λ CDM	$\mathcal{U}(0, 1)$
• X-ray scaling relation			
A_X	-	Normalization of the $M - C_R$ scaling relation	$\mathcal{U}(0.01, 3)$
B_X	-	Mass slope of the $M - C_R$ scaling relation	$\mathcal{U}(0.1, 5)$
D_X	-	Luminosity distance evolution of the $M - C_R$ scaling relation	Fixed to -2
E_X	-	Scale factor evolution of the $M - C_R$ scaling relation	Fixed to 2
F_X	-	Redshift evolution of the mass slope of the $M - C_R$ scaling relation	$\mathcal{U}(-5, 5)$
G_X	-	Redshift evolution of the normalization of the $M - C_R$ scaling relation	$\mathcal{U}(-5, 5)$
σ_X	-	Intrinsic scatter of the $M - C_R$ scaling relation	$\mathcal{U}(0.05, 2)$
• Weak lensing mass calibration			
A_{WL}	-	Scatter in the weak lensing bias from the first principal component	$\mathcal{N}(0, 1)$
B_{WL}	-	Scatter in the weak lensing bias from the second principal component	$\mathcal{N}(0, 1)$
C_{WL}	-	Standardize mass slope of the weak lensing bias	$\mathcal{N}(0, 1)$
D_{WL}	-	Redshift dependent intrinsic scatter in the weak lensing bias	$\mathcal{N}(0, 1)$
$\rho_{M_{\text{WL}}, C_R}$	-	Intrinsic correlation between weak lensing mass and count rate	$\mathcal{U}(-0.9, 0.9)$
• Richness mass calibration			
$\log A_\lambda$	-	Normalization of the $M - \lambda$ scaling relation	$\mathcal{U}(1.4, 6)$
B_λ	-	Mass slope of the $M - \lambda$ scaling relation	$\mathcal{U}(0, 2)$
C_λ	-	Redshift evolution of the normalization of the $M - \lambda$ scaling relation	$\mathcal{U}(-2, 2)$
D_λ	-	Redshift evolution of the mass slope of the $M - \lambda$ scaling relation	$\mathcal{U}(-2, 2)$
σ_λ	-	Intrinsic scatter of the $M - \lambda$ scaling relation	$\mathcal{U}(0.05, 2)$
ρ_{λ, C_R}	-	Intrinsic correlation between richness and count rate	$\mathcal{U}(-0.9, 0.9)$
• Contamination modeling			
f_{AGN}	-	Fraction of AGN contaminants in the extended source sample	$\mathcal{U}(0, 0.1)$
f_{RS}	-	Fraction of RS contaminants in the extended source sample	$\mathcal{U}(0, 0.15)$
• Redshift uncertainty			
σ_z	-	Relative error on the measured redshift	$\mathcal{TN}(0.0050, 0.0011, 0, 1)$
b_z	-	Systematic bias in our redshift estimate	$\mathcal{N}(1.005, 0.037)$
c_z	-	Fraction of objects for which we measure a shifted redshift	$\mathcal{TN}(0.0013, 0.0010, 0, 1)$
$c_{\text{shift}, z}$	-	Amount of redshift shift for c_z fraction of objects	$\mathcal{U}(-0.1, 0.1)$

Parameter	Units	Description	Prior
• Cosmology			
Ω_m	-	Mean matter density at present time	$\mathcal{U}(0.05, 0.95)$
$\log_{10} A_s$	-	Amplitude of the primordial power spectrum	$\mathcal{U}(-10, -8)$
H_0	$\frac{\text{km}}{\text{s Mpc}}$	Hubble expansion rate at present time	$\mathcal{N}(67.77, 0.6)$
Ω_b	-	Mean baryon density at present time	$\mathcal{U}(0.046, 0.052)$
n_s	-	Spectra index of the primordial power spectrum	$\mathcal{U}(0.92, 1.0)$
w_0	-	Dark energy equation of state. Fixed to -1 in Λ CDM	$\mathcal{U}(-2.5, -0.33)$
$\sum m_\nu$	eV	Summed neutrino masses. Fixed to 0 eV in Λ CDM	$\mathcal{U}(0, 1)$
• X-ray scaling relation			
A_X	-	Normalization of the $M - C_R$ scaling relation	$\mathcal{U}(0.01, 3)$
B_X	-	Mass slope of the $M - C_R$ scaling relation	$\mathcal{U}(0.1, 5)$
D_X	-	Luminosity distance evolution of the $M - C_R$ scaling relation	Fixed to -2
E_X	-	Scale factor evolution of the $M - C_R$ scaling relation	Fixed to 2
F_X	-	Redshift evolution of the mass slope of the $M - C_R$ scaling relation	$\mathcal{U}(-5, 5)$
G_X	-	Redshift evolution of the normalization of the $M - C_R$ scaling relation	$\mathcal{U}(-5, 5)$
σ_X	-	Intrinsic scatter of the $M - C_R$ scaling relation	$\mathcal{U}(0.05, 2)$
• Weak lensing mass calibration			
A_{WL}	-	Scatter in the weak lensing bias from the first principal component	$\mathcal{N}(0, 1)$
B_{WL}	-	Scatter in the weak lensing bias from the second principal component	$\mathcal{N}(0, 1)$
C_{WL}	-	Standardize mass slope of the weak lensing bias	$\mathcal{N}(0, 1)$
D_{WL}	-	Redshift dependent intrinsic scatter in the weak lensing bias	$\mathcal{N}(0, 1)$
$\rho_{M_{\text{WL}}, C_R}$	-	Intrinsic correlation between weak lensing mass and count rate	$\mathcal{U}(-0.9, 0.9)$
• Richness mass calibration			
$\log A_\lambda$	-	Normalization of the $M - \lambda$ scaling relation	$\mathcal{U}(1.4, 6)$
B_λ	-	Mass slope of the $M - \lambda$ scaling relation	$\mathcal{U}(0, 2)$
C_λ	-	Redshift evolution of the normalization of the $M - \lambda$ scaling relation	$\mathcal{U}(-2, 2)$
D_λ	-	Redshift evolution of the mass slope of the $M - \lambda$ scaling relation	$\mathcal{U}(-2, 2)$
σ_λ	-	Intrinsic scatter of the $M - \lambda$ scaling relation	$\mathcal{U}(0.05, 2)$
ρ_{λ, C_R}	-	Intrinsic correlation between richness and count rate	$\mathcal{U}(-0.9, 0.9)$
• Contamination modeling			
f_{AGN}	-	Fraction of AGN contaminants in the extended source sample	$\mathcal{U}(0, 0.1)$
f_{RS}	-	Fraction of RS contaminants in the extended source sample	$\mathcal{U}(0, 0.15)$
• Redshift uncertainty			
σ_z	-	Relative error on the measured redshift	$\mathcal{TN}(0.0050, 0.0011, 0, 1)$
b_z	-	Systematic bias in our redshift estimate	$\mathcal{N}(1.005, 0.037)$
c_z	-	Fraction of objects for which we measure a shifted redshift	$\mathcal{TN}(0.0013, 0.0010, 0, 1)$
$c_{\text{shift}, z}$	-	Amount of redshift shift for c_z fraction of objects	$\mathcal{U}(-0.1, 0.1)$

eRASS1 COSMOLOGICAL RESULTS : Λ CDM


Ghirardini, Bulbul, EA et al., 2024



- $\Omega_m = 0.29^{+0.01}_{-0.02}$ (7% precision)
- $\sigma_8 = 0.88 \pm 0.02$ (2% precision)

eRASS1 COSMOLOGICAL RESULTS : w CDM

Ghirardini, Bulbul, EA et al., 2024




DE equation of state

$$p = w\rho$$

$$w = -1.12 \pm 0.12$$

eRASS1 COSMOLOGICAL RESULTS : ν CDM


Ghirardini, Bulbul, EA et al., 2024



- $\sum m_\nu < 0.22$ (95% CL)

IMPACT ON THE S_8 TENSION



Ghirardini, Bulbul, EA et al., 2024



eRASS1 cluster abundance is in good agreement with the CMB

MORE : CLUSTER 2PT CORRELATION FUNCTION

Seppi et al., 2024




Riccardo Seppi

Volume limited selection of the eRASS1 cluster sample


CLUSTER 2PT CORRELATION FUNCTION

Seppi et al., 2024




Projected correlation function of the 4 volume selected eRASS1 cluster samples

CLUSTER 2PT CORRELATION FUNCTION



Seppi et al., 2024



Cosmological results obtained from the eRASS1 clusters correlation function

EVEN MORE : $f(R)$ GRAVITY

EA et al., 2024

Einstein-Hilbert action

$$S_{\text{EH}} = \int d^4x \sqrt{-g} \left(\frac{R}{16\pi G} + \mathcal{L}_m \right) \longrightarrow \tilde{S}_{\text{EH}} = \int d^4x \sqrt{-g} \left(\left[\frac{R + f(R)}{16\pi G} \right] + \mathcal{L}_m \right)$$

Einstein equations

$$G_{\mu\nu} - f_R R_{\mu\nu} - \left(\frac{f}{2} - \square f_R \right) g_{\mu\nu} - \nabla_\mu \nabla_\nu f_R = 8\pi G T_{\mu\nu}$$

Hu & Sawicki, 2007

$$f(R) = -2\Lambda - f_{R0} \frac{R_0^2}{R} \quad \text{with} \quad f_{R0} \ll 1$$

Impacts:

- The halo mass function
- The gravitationnal collapse of structures

EVEN MORE : $f(R)$ GRAVITY

EA et al., 2024

Einstein-Hilbert action

$$S_{\text{EH}} = \int d^4x \sqrt{-g} \left(\frac{R}{16\pi G} + \mathcal{L}_m \right) \longrightarrow \tilde{S}_{\text{EH}} = \int d^4x \sqrt{-g} \left(\left[\frac{R + f(R)}{16\pi G} \right] + \mathcal{L}_m \right)$$

Einstein equations

$$G_{\mu\nu} - f_R R_{\mu\nu} - \left(\frac{f}{2} - \square f_R \right) g_{\mu\nu} - \nabla_\mu \nabla_\nu f_R = 8\pi G T_{\mu\nu}$$

Hu & Sawicki, 2007

$$f(R) = -2\Lambda - f_{R0} \frac{R_0^2}{R} \quad \text{with} \quad f_{R0} \ll 1$$

Impacts:

- The halo mass function
- The gravitationnal collapse of structures

EVEN MORE : $f(R)$ GRAVITY

EA et al., 2024

Einstein-Hilbert action

$$S_{\text{EH}} = \int d^4x \sqrt{-g} \left(\frac{R}{16\pi G} + \mathcal{L}_m \right) \longrightarrow \tilde{S}_{\text{EH}} = \int d^4x \sqrt{-g} \left(\left[\frac{R + f(R)}{16\pi G} \right] + \mathcal{L}_m \right)$$

Einstein equations

$$G_{\mu\nu} \left[-f_R R_{\mu\nu} - \left(\frac{f}{2} - \square f_R \right) g_{\mu\nu} - \nabla_\mu \nabla_\nu f_R \right] = 8\pi G T_{\mu\nu}$$

Hu & Sawicki, 2007

$$f(R) = -2\Lambda - f_{R0} \frac{R_0^2}{R} \quad \text{with} \quad f_{R0} \ll 1$$

Impacts:

- The halo mass function
- The gravitationnal collapse of structures

EVEN MORE : $f(R)$ GRAVITY

EA et al., 2024

Einstein-Hilbert action

$$S_{\text{EH}} = \int d^4x \sqrt{-g} \left(\frac{R}{16\pi G} + \mathcal{L}_m \right) \longrightarrow \tilde{S}_{\text{EH}} = \int d^4x \sqrt{-g} \left(\left[\frac{R + f(R)}{16\pi G} \right] + \mathcal{L}_m \right)$$

Einstein equations

$$G_{\mu\nu} \left[-f_R R_{\mu\nu} - \left(\frac{f}{2} - \square f_R \right) g_{\mu\nu} - \nabla_\mu \nabla_\nu f_R \right] = 8\pi G T_{\mu\nu}$$

Hu & Sawicki, 2007


$$f(R) = -2\Lambda - f_{R0} \frac{R_0^2}{R} \quad \text{with} \quad f_{R0} \ll 1$$

Impacts:

- The halo mass function
- The gravitationnal collapse of structures

EVEN MORE : $f(R)$ GRAVITY

EA et al., 2024




Latest constraints on the Hu-Sawicki parameterization of $f(R)$ gravity from [Cataneo et al., 2014](#).

Impacts:

- The halo mass function
- The gravitationnal collapse of structures

$f(R)$ GRAVITY

EA et al., 2024





f_{R0} increases the number of massive clusters

$f(R)$ GRAVITY


EA et al., 2024

$\log |f_{R0}| < 4.12$ with clusters only




$f(R)$ COSMOLOGICAL CONSTRAINTS

EA et al., 2024



WHAT IS NEXT?



Abdalla et al., 2022



REDSHIFT EVOLUTION OF THE COSMOLOGICAL PARAMETERS

Preliminary


Redshift distribution of the clusters in each bin



REDSHIFT EVOLUTION OF THE COSMOLOGICAL PARAMETERS

Preliminary


eRASS1 cluster abundance + θ^*



REDSHIFT EVOLUTION OF THE COSMOLOGICAL PARAMETERS

Preliminary

eRASS1 cluster abundance + θ^*





REDSHIFT EVOLUTION OF THE COSMOLOGICAL PARAMETERS

Preliminary

Abbott et al., 2023

eRASS1 cluster abundance + θ^*





REDSHIFT EVOLUTION OF THE COSMOLOGICAL PARAMETERS

Preliminary


Abbott et al., 2023

eRASS1 cluster abundance + θ^*




REDSHIFT EVOLUTION OF THE COSMOLOGICAL PARAMETERS


Preliminary



REDSHIFT EVOLUTION OF THE COSMOLOGICAL PARAMETERS




REDSHIFT EVOLUTION OF THE COSMOLOGICAL PARAMETERS



DES weak lensing shear and **DESI targeted LRG 2x2pt + CMB** are in tension with Planck best fit model

REDSHIFT EVOLUTION OF THE COSMOLOGICAL PARAMETERS


Preliminary



Cluster abundance in good agreement with the predicted growth from Planck CMB

REDSHIFT EVOLUTION OF THE COSMOLOGICAL PARAMETERS

Preliminary



SUMMARY

- We measured the cosmological parameters with eRASS1, the largest ICM selected cluster sample to date
 - $\Omega_m = 0.29^{+0.01}_{-0.02}$ (7% precision)
 - $\sigma_8 = 0.88 \pm 0.02$ (2% precision)
 - $\sum m_\nu < 0.22$ (95% CL)
- We do not find traces of the S_8
- More results to come on the growth of structures

Thank you!