

High energy astrophysical neutrinos: Open questions and future prospects

Eli Waxman

Weizmann Institute of Science, Israel

The key goals of High Energy ν astronomy

- Identify the sources of (very) high energy (HE) cosmic-rays,
- Provide unique constraints on models of HE astrophysical sources,
- Possibly: Study ν /fundamental physics.

IceCube's extra-Galactic ν 's: What we have learned

50 TeV - 1 PeV

- The energy production rate densities in the local universe in ~ 100 TeV ν 's and in $>10^{10}$ GeV CRs are similar:

$$\sim 10^{44} \text{erg/Mpc}^3 \text{yr} (\Phi_\nu \approx \Phi_{\text{WB}})$$


→ The sources may be related, but no direct evidence, and the sources are not identified.

- Flux dominated by many weak sources:
 $n_s > 10^{-7}/\text{Mpc}^3$, $L_\nu < 10^{42.5}$ erg/s.

10 - 30 TeV

- $\Phi_\nu \approx 2\Phi_{\text{WB}}$ and is in tension with the 100 GeV γ background, as $\Phi_\gamma(100\text{GeV}) \approx \Phi_\nu$ is expected after EM cascades on the IR background.
- Suggests the existence of “hidden sources”, from which ν 's escape but γ 's don't.

IC 23, ν flux per flavor



Extra-Galactic ν 's: What we are missing

10 TeV – 10 PeV

- The spectrum measurement is crude.
- The flavor ratio measurement (consistent with 1:1:1) is crude.


$10^8 - 10^{10}$ GeV

- A flux measurement (10^{-9} GeV/cm²s sr) will constrain the UHE CR composition.

Sources

- The sources have not been identified.
- A $\sim 3\sigma$ association, including a 300 TeV ν – a Blazar (TXS 0506+056)
 - Models challenged by X- and γ - ray observations [e.g. Murase 18].
 - Blazars do not dominate the background [e.g. IC 17].
- A $\sim 4\sigma$ association, < 10 TeV – an AGN (Syefert)/Starburst galaxy (NGC1068)
 - (2 lower significance associations [Neronov et al. 24])
 - A hidden γ -ray source, $L_\nu > \sim 30 L_\gamma$ (1 GeV – 1 TeV)

IC 23, spectral index



NGC1068 association challenges: I. Physics

$$L_{\text{CR}} \sim L_{\text{Eddington}}$$

- Models postulate ν production at the $\sim 10 R_s$ vicinity of the BH, to obtain efficient ν production and strong suppression of 100GeV photon emission by pair production with UV photons. [Das et al. 24, Padovani et al. 24, Lemoine & Rieger 24, Inoue et al. 24]

- $100 \text{ TeV } p + \sim 1 \text{ keV } \gamma \rightarrow \sim 5 \text{ TeV } \nu$

$$100 \text{ TeV } p + \sim 10 \text{ eV } \gamma \rightarrow e^\pm$$

$$\frac{L_\pi}{L_\pm} \approx \frac{L_X}{L_{UV}} \approx 10^{-1.5}$$

since $(dE/E \times \sigma / \epsilon_\gamma)$ similar for pion & for pair production

$$\rightarrow L_{\text{CR}}(\sim 100 \text{ TeV}) \approx 2 L_\nu \frac{L_{UV}}{L_X} \approx 10^{44} \text{ erg/s} \approx 0.1 L_{\text{Eddington}}$$

$$L_{\text{CR}}(1\text{GeV} - 100\text{TeV}) \approx L_{\text{Eddington}}$$

- This would imply a modification of our basic understanding of AGN physics.

NGC1068 association challenges: II. Statistics

We are unique

- The average cosmic ν luminosity within a 10Mpc sphere is

$$10^{44} \text{erg/Mpc}^3 \text{yr} \times (10 \text{ Mpc sphere}) = 10^{40} \text{erg/s} .$$

The inferred ν luminosity of NGC1068 is $3 \times 10^{42} \text{erg/s}$

→ We are at a special location, with exceptionally (300 times) larger than average flux.

- If the ν luminosity of NGC1068 were typical for Seyferts, the resulting ν luminosity would exceed the observed one by a factor >100 :

$$3 \times 10^{42} \text{erg/s} \times (10^{-4} \text{ Mpc}^{-3}) = 10^{46} \text{erg/Mpc}^3 \text{yr}.$$

→ NGC1068 must be a “rare Seyfert”- e.g.:

Estimated large intrinsic X-ray luminosity corresponding to $\sim 10^{-5} \text{ Mpc}^{-3}$,
exceeding the observed flux by a factor >10 .

ν 's from Supernova driven shocks:

Alternative “ γ -ray hidden” sources of $>\sim 10$ TeV ν 's

- Collisionless SN shocks driven into the circum/inter stellar medium (CSM/ISM) are likely CR accelerators
- CR-CSM/ISM pp interaction is a likely source of ν 's

[e.g. Murase et al. 11; Petropoulou et al. 17; Sarmah et al. 22]


- For typical CSM wind/ISM
 - ν emission over months-years
 - Low ν luminosity, small contribution to the ν background
 - Accompanied by HE γ -rays (low pair production optical depth)
- Much larger ν luminosity may be produced if the progenitor star is surrounded by a compact optically thick CSM

[Katz, Sapir & EW 11, Li 19]


Evidence for the prevalence of optically thick CSM at $\sim 10^{14.5}$ cm around Type II SN progenitors

- Most type II SN progenitors show $>10^{47}$ erg precursors, ~ 1 yr preceding SN
Unbinding ~ 0.1 -1 solar mass, ejected to $\sim (100 \text{ km/s} \times 1 \text{ yr}) = 10^{14.5}$ cm
[Strotjohannet al. 21]
- 50% of regular type II SNe show a slow, ~ 3 -d, rise of the light-curve to large, $\sim 10^{43.5}$ erg/s, UV luminosity –
A SN shock breaking through an optically thick CSM,
 ~ 0.1 solar mass out to $\sim (10,000 \text{ km/s} \times 3\text{-d}) = 10^{14.5}$ cm
[Irani et al. 24]
- $>50\%$ of regular type II SNe show narrow lines of ionized species, disappearing after ~ 2 -3 days –
Emission from ~ 0.1 solar mass CSM excited by the shock UV luminosity,
disappearing after shock crossing at $\sim (10,000 \text{ km/s} \times 3\text{-d}) = 10^{14.5}$ cm
[Bruch et al. 23]

SN 2023ixf, 6.4 Mpc



[Zimmerman et al. 24]



ν 's from SN shock breakouts through compact optically thick CSM

- ν 's produced by pp collisions during and shortly after the transition of the SN shock from being radiation-mediated to collisionless

$$L_{\text{br, UV}} \sim 10^{43.5} \text{ erg/s}, \quad L_{\text{br, } \nu} \sim 0.05 L_{\text{br, UV}}, \quad \Delta t \sim 3\text{d}$$


- Significant contribution to the $> \sim 10$ TeV ν background if ~ 0.1 solar-mass ejection is prevalent on a year time scale preceding explosion

$$0.05 \times 10^{43.5} \text{ erg/s} \times 3\text{d} \times (10^{-4} / \text{Mpc}^3 \text{ yr}) = 5 \times 10^{43} \text{ erg/Mpc}^3 \text{ yr}$$

- The pair production optical depth at $> 100\text{GeV}$ is $> 10^4$, due to the compact size and large UV luminosity
- The rate of SNe producing > 1 (ν induced μ) events in a 1 km^2 detector is $< \sim 1/10\text{yr}$ (SN 2023ixf – 0.2ν induced μ). A single association will be significant at 99.9% CL.

Identifying >10 TeV steady sources: ($10 \text{ km}^2 \times 10 \text{ yr}$) required, beyond 2040

- ~ 1 atmospheric ν_μ per 1 squared degree.
 - Source density $>10^{-7}/\text{Mpc}^3$, source number $>10^6$.
 - 1% of astrophysical ν 's, i.e. $<\sim 10$, originate at $d < 100$ Mpc.
- Association with sources by correlation with catalogs is difficult.
- ($10 \text{ km}^2 + \Delta\theta < 0.5 \text{ deg}$) $\times 10 \text{ yr}$
+ complete source catalog to 200 Mpc
required for a 3σ association with nearby sources.




Probability for a 3σ detection, for 10 yrs IceCubeGen2, and complete catalogs to 0.2 Gpc

Identifying >10 TeV transient sources:

Which X/ γ -ray flares are viable candidates?

The number density of active X/ γ -ray flares



$L_\gamma < 10^{45}$ erg/s

$L_\gamma > 10^{50}$ erg/s
< $\sim 1\%$ of the flux

Identifying >10 TeV transient sources: Unlikely with X/ γ -ray observations

- Coincident ν /EM transient detection increases the significance of an angular association for transient duration $\Delta t \sim \text{days} \ll T \sim 1 \text{ yr}$.

- A handful of events per year- $d > 1 \text{ Gpc}$ sources.

$$L_\gamma < 10^{45} \text{ erg/s} \text{ (assuming } L_\nu \leq L_\gamma \text{)}.$$

- The required sensitivity:

$$f_\gamma < \frac{10^{44} \text{ erg/s}}{4\pi(1\text{Gpc})^2} = 10^{-12} \text{ erg/cm}^2\text{s}$$

- May be possible with XRT ($\sim 1 \text{ keV}$; 0.1 sq. deg FOV).


Well below the sensitivity of

BAT/GBM ($\sim 1 \text{ MeV}$), Fermi LAT ($\sim 1 \text{ GeV}$), HESS, MAGIC, LHAASO (sub-TeV).

Marginal for CTA/LST (sub-TeV).

Identifying >10 TeV transient sources: UV/Optical surveys open new opportunities

- Many candidate sources are expected to be UV bright.
 - Supernovae:
 - Jet-driven explosions,
(LL GRBs),
Ejecta – Circumstellar Medium interaction,
 - Tidal disruption events.
- The ULTRASAT UV space telescope will enable a systematic detection and detailed study of these fast transients, and possibly coincident ν /EM detections.




A handful of ν - γ associations for the nearest, yet quite distant – 0.5 Gpc, sources, will not enable a systematic detection and study of the transient sources.


Sensitive, wide FOV UV/Optical surveys are key for systematic study and understanding.

Identifying >10 TeV transient sources: UV/Optical surveys open new opportunities

- Many candidate sources are expected to be UV bright.
 - Supernovae: Jet-driven explosion (LL GRBs), Ejecta – Circumstellar interaction,
 - Tidal disruption
- The ULTRASAT will enable a systematic and detailed study of these fast transients necessary coincident ν /EM detections.



ULTRASAT



A handful of ν - γ associations for the nearest, yet quite distant – 0.5 Gpc, sources, will not enable a systematic detection and study of the transient sources.

Sensitive, wide FOV UV/Optical surveys are key for systematic study and understanding.

Summary

- HE ν astronomy has the potential to
 - Provide unique constraints on models of HE astrophysical sources, and
 - Identify the sources of (very) HE cosmic-rays.
- Fulfilling the potential relies on the EM identification of the neutrino sources.
- $M_{\text{eff}} \sim 10 \text{ Gton} @ 10^5 - 10^8 \text{ GeV}$ (IceCube Gen2 + KM3NeT/ GVD/ P1/ TRIDENT/ HUNT) is required to
 - Detect multiple events from few nearby sources (eg starbursts),
 - Possibly detect luminous transients (GRB/TDE-jet) contributing $\sim 1\%$ of the flux,
 - Obtain accurate ν spectrum, angular distribution and flavor content.
- EM follow-up observations may identify hour-day long transient sources, e.g. SN CSM breakouts.
Crucial for a systematic study of the sources.
EM detector requirements: $\text{FOV} > 1 \text{ deg}^2$, Sensitivity better than $10^{-13} \text{ erg/cm}^2\text{s}$.
 - May be possible at X-ray (XRT), marginal at sub-TeV (CTA).
 - UV/O (ULTRASAT) surveys are key for systematic study.
- $10^8 - 10^{10} \text{ GeV}$: A flux measurement ($10^{-9} \text{ GeV/cm}^2\text{s sr}$) will constrain the UHE CR composition (Radio).