

CP-violation in complex-singlet extensions of 2HDM (2HDMS)

Jayita Lahiri

II. Institut für Theoretische Physik, Universität Hamburg
3rd ECFA Workshop, Paris, 2024

Work in progress in collaboration with Gudrid Moortgat-Pick

CP-violation

- Baryon-asymmetry of the universe \rightarrow additional sources of CP-violation beyond SM is 'necessary'.
- It is possible to have additional CPV in models with extended scalar sectors.
- Constraints comes from :
 - 1 EDM experiments
 - 2 Collider experiments
 - 3 Requirement from observed baryon-asymmerty.

CP-violation in 2HDM

The most general 2HDM scalar potential :

$$\begin{aligned} V_{2HDM} = & -m_{11}^2 \Phi_1^\dagger \Phi_1 - m_{22}^2 \Phi_2^\dagger \Phi_2 - [m_{12}^2 \Phi_1^\dagger \Phi_2 + h.c.] + \frac{\lambda_1}{2} (\Phi_1^\dagger \Phi_1)^2 \\ & + \frac{\lambda_2}{2} (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) + \lambda_4 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) \\ & + \left[\frac{\lambda_5}{2} (\Phi_1^\dagger \Phi_2) + \lambda_6 (\Phi_1^\dagger \Phi_1) + \lambda_7 (\Phi_2^\dagger \Phi_2) \right] (\Phi_1^\dagger \Phi_2) + h.c \end{aligned}$$

- Exact \mathcal{Z}_2 symmetry i.e. $m_{12}^2 = \lambda_6 = \lambda_7 = 0 \rightarrow$ no CP-violation, the phase of λ_5 can be rotated away, by a global U(2) transformation of the potential. No CP-violation. *Ilya F. Ginzburg and Maria Krawczyk (Arxiv:hep-ph/0408011)*
- Softly-broken \mathcal{Z}_2 -symmetry i.e. $m_{12}^2 \neq 0, \lambda_6, \lambda_7 = 0$ CP-violation is possible. In the alignment limit, CP-violation becomes negligible. Imaginary part of *all* possible U(2)-invariants ≈ 0 . *John F. Gunion and Howard E Haber (Arxiv:hep-ph/0506227)*

Yukawa-aligned 2HDM

- Hard breaking of \mathcal{Z}_2 i.e. $m_{12}^2, \lambda_6, \lambda_7 \neq 0$, CP-violation can be significant.

S. Kanemura, M. Kubota and K. Yagyu (Arxiv:2004.03943)

$$\mathcal{L}_{\text{Yukawa}} = \sum_{k=1}^2 \left(\bar{Q}_{LYu,k}^\dagger \tilde{\Phi}_k u_R + \bar{Q}_{LYd,k} \Phi_k d_R + \bar{L}_{LYe,k} \Phi_k e_R \right)$$

- In the absence of \mathcal{Z}_2 symmetry, to avoid tree-level FCNC Yukawa matrices associated with the two doublets are assumed to be proportional to each other. *A. Pich and P. Tuzon (Arxiv:0908.1554)*

$$y_{f,2} = \zeta_f y_{f,1}$$

- ζ can be complex and the source of CP-violation.
- In 2HDM (Yukawa-aligned), in the *exact* alignment limit, **the CP-violation stems from Yukawa sector and not from the CP-mixing in the scalar sector.**
- Also the Yukawa interaction of the 125 GeV Higgs is CP-conserving in the alignment limit.

2HDM + complex singlet (2HDMS)

There are two major motivation to go to the complex singlet extension of 2HDM are :

- 1 The scalar sector of 2HDMS resembles that of NMSSM, when the complex scalar is charged under a \mathcal{Z}_3 symmetry.
- 2 The model can accommodate a dark matter component when the complex scalar is charged under a \mathcal{Z}'_2 symmetry, as well as an excess such as 95 GeV observed at CMS as well as LEP in $\gamma\gamma$ and $b\bar{b}$ final state.
- 3 We would investigate whether, there are additional (physical) sources of CP-violation in 2HDMS.

2HDMS potential- \mathcal{Z}'_2 symmetric case

$$V_{2\text{HDMS}} = V_{2\text{HDM}} + V_S$$

$$\begin{aligned} V_S = & m_S^2 S^\dagger S + \left[\frac{m_S'^2}{2} S^2 + h.c. \right] + \left[\frac{\lambda_1''}{24} S^4 + h.c. \right] + \left[\frac{\lambda_2''}{6} (S^2 S^\dagger S) + h.c. \right] \\ & + \frac{\lambda_3''}{4} (S^\dagger S)^2 + S^\dagger S [\lambda_1' \Phi_1^\dagger \Phi_1 + \lambda_2' \Phi_2^\dagger \Phi_2] + [S^2 (\lambda_4' \Phi_1^\dagger \Phi_1 + \lambda_5' \Phi_2^\dagger \Phi_2) + h.c.] \end{aligned}$$

2HDMS potential- \mathcal{Z}'_2 symmetric case

$$V_{2\text{HDMS}} = V_{2\text{HDM}} + V_S$$

$$\begin{aligned} V_S = & m_S^2 S^\dagger S + \left[\frac{m_S'^2}{2} S^2 + h.c. \right] + \left[\frac{\lambda_1''}{24} S^4 + h.c. \right] + \left[\frac{\lambda_2''}{6} (S^2 S^\dagger S) + h.c. \right] \\ & + \frac{\lambda_3''}{4} (S^\dagger S)^2 + S^\dagger S [\lambda_1' \Phi_1^\dagger \Phi_1 + \lambda_2' \Phi_2^\dagger \Phi_2] + [S^2 (\lambda_4' \Phi_1^\dagger \Phi_1 + \lambda_5' \Phi_2^\dagger \Phi_2) + h.c.] \\ & + [\lambda_6' \Phi_1^\dagger \Phi_2 S^\dagger S + h.c.] + [\lambda_7' \Phi_1^\dagger \Phi_2 S^2 + h.c.] \end{aligned}$$

2HDMS potential- \mathcal{Z}'_2 symmetric case

$$V_{2\text{HDMS}} = V_{2\text{HDM}} + V_S$$

$$\begin{aligned} V_S = & m_S^2 S^\dagger S + \left[\frac{m_S'^2}{2} S^2 + h.c. \right] + \left[\frac{\lambda_1''}{24} S^4 + h.c. \right] + \left[\frac{\lambda_2''}{6} (S^2 S^\dagger S) + h.c. \right] \\ & + \frac{\lambda_3''}{4} (S^\dagger S)^2 + S^\dagger S [\lambda_1' \Phi_1^\dagger \Phi_1 + \lambda_2' \Phi_2^\dagger \Phi_2] + [S^2 (\lambda_4' \Phi_1^\dagger \Phi_1 + \lambda_5' \Phi_2^\dagger \Phi_2) + h.c.] \\ & + [\lambda_6' \Phi_1^\dagger \Phi_2 S^\dagger S + h.c.] + [\lambda_7' \Phi_1^\dagger \Phi_2 S^2 + h.c.] \end{aligned}$$

- Although $m_S'^2$, λ_1'' , λ_2'' , λ_4' , λ_5' , λ_6' and λ_7' , are all in principle complex, only $\text{Im}(\lambda_6')$ and $\text{Im}(\lambda_7')$, can introduce mixing between scalar and pseudoscalars, due to the presence of $\Phi_1^\dagger \Phi_2$ term.
- \mathcal{Z}'_2 -breaking of the 2HDM potential is essential here as well for CP-violation.

\mathcal{Z}'_2 symmetric case - can we get dark matter and CP-violation simultaneously?

- In order to accommodate a dark matter candidate, we need $S = v_S + h_S + ia_S$ and one of the component fields acquire zero vev and two separate \mathcal{Z}_2 symmetry should be imposed on the two fields.
- Two separate \mathcal{Z}_2 on the two component fields of S can be imposed, only when the coefficients are real.
- Two separate \mathcal{Z}_2 , is sufficient to get dark matter, not necessary.
- The necessary conditions are $\lambda'_4, \lambda'_5, \lambda'_7, m_S'^2$ are real and $\text{Im}[\lambda_1''] = -2 \times \text{Im}[\lambda_2'']$.
- In that case we will be left with two independent phases, of λ'_6 and λ_1'' .
- In addition, to be in the alignment limit, one needs $\text{Re}[\lambda_1'] = -2 \times \text{Re}[\lambda_4']$.

Minimization of the potential

$$m_{11}^2 = \frac{1}{2}\lambda_1 v^2 + \frac{1}{2}\lambda'_1 v_S^2 + \text{Re}[\lambda'_4] v_S^2,$$

$$m_{12}^2 = \frac{1}{2}(\lambda_6 v^2 + \lambda'_6 v_S^2 + \lambda'_7 v_S^2)$$

$$m_S^2 = -(\text{Re}[m_S'^2] + \frac{1}{2}\lambda'_1 v^2 + \text{Re}[\lambda'_4] v^2) + \left(\frac{\text{Re}[\lambda_1'']}{12} + \frac{\text{Re}[\lambda_2'']}{3} + \frac{\text{Re}[\lambda_3'']}{4} \right) v_S^2$$

$$\text{Im}[m_S'^2] = \left(\frac{\text{Im}[\lambda_1'']}{12} + \frac{\text{Im}[\lambda_2'']}{6} \right) v_S^2 + \text{Im}[\lambda'_4] v^2$$

Dark Matter mass

$$m_{\text{DM}}^2 = -2\text{Re}[m_S'^2] - \frac{1}{3}v_S^2(\text{Re}[\lambda_1''] + \text{Re}[\lambda_2'']) - 2v^2\text{Re}[\lambda'_4]$$

Mass-matrix and CP-mixing in the scalar sector

In the Higgs-basis

$$\mathcal{M}_{ij}^2 = \left(\begin{array}{c|ccc|c} m_{11} & 0 & 0 & 0 & 0 \\ \hline 0 & m_{22} & 0 & m_{24} & 0 \\ 0 & 0 & m_{33} & m_{34} & 0 \\ 0 & m_{24} & m_{34} & m_{44} & 0 \\ \hline 0 & 0 & 0 & 0 & m_{55} \end{array} \right)$$

$$m_{11} = \lambda_1 v^2 = m_h^2; m_h = 125\text{GeV}$$

$$m_{22} = -m_{22}^2 + \left(\frac{\lambda_2' + \text{Re}[\lambda_5']}{2} \right) v_S^2 + \left(\frac{\lambda_3 + \lambda_4 + \text{Re}[\lambda_5]}{2} \right) v^2$$

$$m_{24} = v v_S \text{Re}[\lambda_6']$$

$$m_{33} = -m_{22}^2 + \left(\frac{\lambda_2' + \text{Re}[\lambda_5']}{2} \right) v_S^2 + \left(\frac{\lambda_3 + \lambda_4 - \text{Re}[\lambda_5]}{2} \right) v^2$$

$$m_{34} = v v_S \text{Im}[\lambda_6']$$

$$m_{44} = \frac{1}{6} v_S^2 (\text{Re}[\lambda_1''] + 4\text{Re}[\lambda_2''] + 3\text{Re}[\lambda_3''])$$

$$m_{55} = -2\text{Re}[m_S'^2] - \left(\frac{\text{Re}[\lambda_1''] + \text{Re}[\lambda_2'']}{3} \right) v_S^2 - 2\text{Re}[\lambda_4'] v^2 = m_{\text{DM}}^2$$

Yukawa sector

In terms of fermion mass eigenstates,

$$\begin{aligned}\mathcal{L}_{\text{Yukawa}} = & - \sum_{f=u,d,e} \left\{ \bar{f}_L M_f f_R + \sum_{j=1}^3 \bar{f}_L \left(\frac{M_f}{v} \kappa_f^j \right) f_R H_j^0 + h.c. \right\} \\ & - \frac{\sqrt{2}}{v} \left\{ -\zeta_u \bar{u}_R (M_u^\dagger V_{\text{CKM}}) d_L + \zeta_d \bar{u}_L (V_{\text{CKM}} M_d) d_R + \zeta_e \bar{\nu}_L M_e e_R \right\} H^+ + h.c.\end{aligned}$$

$$\kappa_f^j = \mathcal{R}_{1j} + [\mathcal{R}_{2j} + i(-2I_f)\mathcal{R}_{3j}] |\zeta_f| e^{i(-2I_f)\theta_f}$$

- In 2HDM, in the alignment limit ($R_{ij} = \delta_{ij}$), the CP-violation in the Yukawa sector can not come from the CP-mixing in the scalar sector. It must come from the phases of the Yukawa matrices.
- In 2HDMS, there can be additional source of CP-violation from the scalar sector mixing.
- In both cases the Yukawa couplings of the H_1^0 does not contain any CP-violating phases and therefore SM-like.

Electric Dipole Moments

$$H_{\text{EDM}} = -d_f \frac{\vec{S}}{|\vec{S}|} \cdot \vec{E}$$

Under the time reversal transformation:

$\mathcal{T}(\vec{S}) = -\vec{S}$ and $\mathcal{T}(\vec{E}) = +\vec{E}$ the sign of this term H_{EDM} is flipped. CP symmetry is broken.



In EFT language,

$$\mathcal{L}_{\text{EDM}} = -\frac{d_f}{2} \bar{f} \sigma^{\mu\nu} (i\gamma^5) f F_{\mu\nu}$$

The most recent bound on electron EDM comes from ACME collaboration via measurement of thorium-monoxide EDM.

$$|d_e| < 1.1 \times 10^{-29} \text{e.cm}$$

Bar-Zee diagrams





$$d_f = d_f(\text{fermion}) + d_f(\text{Higgs}) + d_f(\text{gauge})$$

Each contribution $d_f(X)$ further consists of

$$d_f(X) = d_f^\gamma(X) + d_f^Z(X) + d_f^W(X)$$

- The gauge boson loops contribute negligibly in the alignment limit.
- The fermion and scalar boson loops contribute at equivalent strength.
- One loop contribution is suppressed by at least 4-5 orders of magnitude.

Results



S. Kanemura, M. Kubota and K. Yagyu (Arxiv:2004.03943)




Figure: Orange : $m_{h_4}=95$ GeV, Maroon : $m_{h_4}=200$ GeV

I chose the benchmark in Yukawa-aligned 2HDM scenario with
 $[\theta_u, \theta_7] = [\frac{\pi}{2}, \frac{\pi}{2}]$, $m_{h_2} = 280$ GeV, $m_{h_3} = m_{h^\pm} = 230$ GeV.

Summary

- Hard breaking of \mathcal{Z}_2 -symmetry of 2HDM in the alignment limit is necessary for CP-violation. This statement holds even in complex-singlet extension (\mathcal{Z}'_2 -symmetric) of 2HDM.
- It is possible to accommodate DM and CP-violation in 2HDMS, with restrictions on complex couplings.
- The fine-tuned cancellations required to satisfy EDM bounds in Yukawa-aligned 2HDM can be alleviated in 2HDMS.

Further things to do

- What happens when the \mathcal{Z}_3 symmetric (NMSSM-like) complex singlet sector?
- Imposing existing experimental constraints on the parameter space.
- Constructing CP-odd observables to probe CP-violating effects, eg. azimuthal angles, asymmetries, impact of beam polarization in lepton colliders.
- Interrelation with dark matter observables.
- Can the amount of allowed CP-violation in this model, be sufficient for baryogenesis?