

MAPTMD24: New insights on flavor dependence in TMD extractions

JHEP 08 (2024) 232

Resummation, Evolution and Factorization 2024

Lorenzo Rossi

MAP Collaboration

October 15th




Istituto Nazionale di Fisica Nucleare




**UNIVERSITÀ
DEGLI STUDI
DI MILANO**

TMD factorization – Drell-Yan process

TMD factorization – Drell-Yan process




TMD factorization – Drell-Yan process



In $q_T^2 \ll Q^2$ and $M^2 \ll Q^2$ region:


TMD factorization – Drell-Yan process



In $q_T^2 \ll Q^2$ and $M^2 \ll Q^2$ region:

$$F_{UU}^1(x_A, x_B, \mathbf{q}_T, Q) = x_A x_B \mathcal{H}^{DY}(Q; \mu) \sum_a c_a(Q^2) \int d|\mathbf{b}_T| |\mathbf{b}_T| J_0(|\mathbf{q}_T| |\mathbf{b}_T|) \hat{f}_1^a(x_A, \mathbf{b}_T^2; \mu, \zeta_A) \hat{f}_1^b(x_B, \mathbf{b}_T^2; \mu, \zeta_B)$$


TMD factorization – Drell-Yan process



In $q_T^2 \ll Q^2$ and $M^2 \ll Q^2$ region:

$$F_{UU}^1(x_A, x_B, \mathbf{q}_T, Q) = x_A x_B \mathcal{H}^{DY}(Q; \mu) \sum_a c_a(Q^2) \int d|\mathbf{b}_T| |\mathbf{b}_T| J_0(|\mathbf{q}_T| |\mathbf{b}_T|) \hat{f}_1^a(x_A, \mathbf{b}_T^2; \mu, \zeta_A) \hat{f}_1^b(x_B, \mathbf{b}_T^2; \mu, \zeta_B)$$


TMD factorization – Drell-Yan process



In $q_T^2 \ll Q^2$ and $M^2 \ll Q^2$ region:

$$F_{UU}^1(x_A, x_B, \mathbf{q}_T, Q) = x_A x_B \mathcal{H}^{DY}(Q; \mu) \sum_a c_a(Q^2) \int d|\mathbf{b}_T| |\mathbf{b}_T| J_0(|\mathbf{q}_T| |\mathbf{b}_T|) \hat{f}_1^a(x_A, \mathbf{b}_T^2; \mu, \zeta_A) \hat{f}_1^b(x_B, \mathbf{b}_T^2; \mu, \zeta_B)$$


TMD factorization – Drell-Yan process



In $q_T^2 \ll Q^2$ and $M^2 \ll Q^2$ region:

$$F_{UU}^1(x_A, x_B, \mathbf{q}_T, Q) = x_A x_B \mathcal{H}^{DY}(Q; \mu) \sum_a c_a(Q^2) \int d|\mathbf{b}_T| |\mathbf{b}_T| J_0(|\mathbf{q}_T| |\mathbf{b}_T|) \hat{f}_1^a(x_A, \mathbf{b}_T^2; \mu, \zeta_A) \hat{f}_1^b(x_B, \mathbf{b}_T^2; \mu, \zeta_B)$$

TMD factorization – Drell-Yan process




In $q_T^2 \ll Q^2$ and $M^2 \ll Q^2$ region:

$$F_{UU}^1(x_A, x_B, \mathbf{q}_T, Q) = x_A x_B \mathcal{H}^{DY}(Q; \mu) \sum_a c_a(Q^2) \int d|\mathbf{b}_T| |\mathbf{b}_T| J_0(|\mathbf{q}_T| |\mathbf{b}_T|) \hat{f}_1^a(x_A, \mathbf{b}_T^2; \mu, \zeta_A) \hat{f}_1^b(x_B, \mathbf{b}_T^2; \mu, \zeta_B)$$


W term

TMD Factorization - SIDIS process



$$\begin{aligned}
 F_{UU,T}(x.z; \mu_F, \mathbf{P}_{hT}^2, Q^2) = & x \sum_a H_{UU,T}^a(Q^2, \mu^2) \int d^2 \mathbf{k}_\perp d^2 \mathbf{P}_\perp f_1^{\mathbf{a}}(x, \mathbf{k}_\perp^2; \mu^2) D_1^{\mathbf{a} \rightarrow \mathbf{h}}(z, \mathbf{P}_\perp^2; \mu^2) \delta^{(2)}(z \mathbf{k}_\perp - \mathbf{P}_{hT} + \mathbf{P}_\perp) \\
 & + Y_{UU,T}(Q^2, \mathbf{P}_{hT}^2) + \mathcal{O}(M^2/Q^2)
 \end{aligned}$$


TMD Factorization - SIDIS process



$$F_{UU,T}(x \cdot z; \mu_F, \mathbf{P}_{hT}^2, Q^2) = \boxed{x \sum_a H_{UU,T}^a(Q^2, \mu^2) \int d^2 \mathbf{k}_\perp d^2 \mathbf{P}_\perp f_1^a(x, \mathbf{k}_\perp^2; \mu^2) D_1^{\mathbf{a} \rightarrow \mathbf{h}}(z, \mathbf{P}_\perp^2; \mu^2) \delta^{(2)}(z \mathbf{k}_\perp - \mathbf{P}_{hT} + \mathbf{P}_\perp)} \\ + Y_{UU,T}(Q^2, \mathbf{P}_{hT}^2) + \mathcal{O}(M^2/Q^2)$$

W Term


TMD Factorization - SIDIS process



TMD Factorization - SIDIS process

TMD FF

TMD PDF



$$F_{UU,T}(x \cdot z; \mu_F, \mathbf{P}_{hT}^2, Q^2) = x \sum_a H_{UU,T}^a(Q^2, \mu^2) \int d^2 \mathbf{k}_\perp d^2 \mathbf{P}_\perp f_1^a(x, \mathbf{k}_\perp^2; \mu^2) D_1^{a \rightarrow h}(z, \mathbf{P}_\perp^2; \mu^2) \delta^{(2)}(z \mathbf{k}_\perp - \mathbf{P}_{hT} + \mathbf{P}_\perp)$$


$$+ Y_{UU,T}(Q^2, \mathbf{P}_{hT}^2) + \mathcal{O}(M^2/Q^2)$$

W Term

TMD Factorization - SIDIS process

TMD FF

TMD PDF



$$F_{UU,T}(x \cdot z; \mu_F, \mathbf{P}_{hT}^2, Q^2) = x \sum_a H_{UU,T}^a(Q^2, \mu^2) \int d^2 \mathbf{k}_\perp d^2 \mathbf{P}_\perp f_1^a(x, \mathbf{k}_\perp^2; \mu^2) D_1^{a \rightarrow h}(z, \mathbf{P}_\perp^2; \mu^2) \delta^{(2)}(z \mathbf{k}_\perp - \mathbf{P}_{hT} + \mathbf{P}_\perp)$$

$$+ Y_{UU,T}(Q^2, \mathbf{P}_{hT}^2) + \mathcal{O}(M^2/Q^2)$$


W Term

- The W term dominates in the region where $\mathbf{q}_T \ll \mathbf{Q}$

TMD Factorization - SIDIS process

TMD FF

TMD PDF



$$F_{UU,T}(x \cdot z; \mu_F, \mathbf{P}_{hT}^2, Q^2) = x \sum_a H_{UU,T}^a(Q^2, \mu^2) \int d^2 \mathbf{k}_\perp d^2 \mathbf{P}_\perp f_1^a(x, \mathbf{k}_\perp^2; \mu^2) D_1^{a \rightarrow h}(z, \mathbf{P}_\perp^2; \mu^2) \delta^{(2)}(z \mathbf{k}_\perp - \mathbf{P}_{hT} + \mathbf{P}_\perp)$$

$$+ Y_{UU,T}(Q^2, \mathbf{P}_{hT}^2) + \mathcal{O}(M^2/Q^2)$$

W Term

- The W term dominates in the region where $q_T \ll Q$
- The Y term has been excluded in the MAP analysis

TMD Factorization - structure of TMDs

$$\begin{aligned}\hat{f}_1^q(x_B, \mathbf{b}_T; \mu_F, \zeta_F) &= [C \otimes f_1](x_B, b_\star; \mu_{b_\star}, \mu_{b_\star}^2) \exp \left\{ \int_{\mu_{b_\star}}^{\mu_F} \frac{d\mu'}{\mu'} \gamma(\mu', \zeta_F) \right\} \\ &\times \left(\frac{\zeta}{\mu_{b_\star}^2} \right)^{K(b_\star, \mu_{b_\star})/2} \left[\frac{\zeta}{Q_0} \right]^{-g_K(\mathbf{b}_T)/2} f_1^{NP}(x, \mathbf{b}_T; \zeta, Q_0)\end{aligned}$$

TMD Factorization - structure of TMDs

Matching coeff.
(perturbative calculable)

$$\hat{f}_1^q(x_B, \mathbf{b}_T; \mu_F, \zeta_F) = [C \otimes f_1](x_B, b_\star; \mu_{b_\star}, \mu_{b_\star}^2) \exp \left\{ \int_{\mu_{b_\star}}^{\mu_F} \frac{d\mu'}{\mu'} \gamma(\mu', \zeta_F) \right\}$$
$$\times \left(\frac{\zeta}{\mu_{b_\star}^2} \right)^{K(b_\star, \mu_{b_\star})/2} \left[\frac{\zeta}{Q_0} \right]^{-g_K(\mathbf{b}_T)/2} f_1^{NP}(x, \mathbf{b}_T; \zeta, Q_0)$$

TMD Factorization - structure of TMDs

Matching coeff.
(perturbative calculable)

Collinear PDFs
(previous fit)

$$\hat{f}_1^q(x_B, \mathbf{b}_T; \mu_F, \zeta_F) = [C \otimes f_1](x_B, b_\star; \mu_{b_\star}, \mu_{b_\star}^2) \exp \left\{ \int_{\mu_{b_\star}}^{\mu_F} \frac{d\mu'}{\mu'} \gamma(\mu', \zeta_F) \right\}$$
$$\times \left(\frac{\zeta}{\mu_{b_\star}^2} \right)^{K(b_\star, \mu_{b_\star})/2} \left[\frac{\zeta}{Q_0} \right]^{-g_K(\mathbf{b}_T)/2} f_1^{NP}(x, \mathbf{b}_T; \zeta, Q_0)$$

TMD Factorization - structure of TMDs

Matching coeff.
(perturbative calculable)

Collinear PDFs
(previous fit)

Perturbative Sudakov
evolution factor

$$\hat{f}_1^q(x_B, \mathbf{b}_T; \mu_F, \zeta_F) = [C \otimes f_1](x_B, b_\star; \mu_{b_\star}, \mu_{b_\star}^2) \exp \left\{ \int_{\mu_{b_\star}}^{\mu_F} \frac{d\mu'}{\mu'} \gamma(\mu', \zeta_F) \right\}$$
$$\times \left(\frac{\zeta}{\mu_{b_\star}^2} \right)^{K(b_\star, \mu_{b_\star})/2} \left[\frac{\zeta}{Q_0} \right]^{-g_K(\mathbf{b}_T)/2} f_1^{NP}(x, \mathbf{b}_T; \zeta, Q_0)$$

TMD Factorization - structure of TMDs

Matching coeff.
(perturbative calculable)

Collinear PDFs
(previous fit)

Perturbative Sudakov
evolution factor

$$\hat{f}_1^q(x_B, \mathbf{b}_T; \mu_F, \zeta_F) = [C \otimes f_1](x_B, b_\star; \mu_{b_\star}, \mu_{b_\star}^2) \exp \left\{ \int_{\mu_{b_\star}}^{\mu_F} \frac{d\mu'}{\mu'} \gamma(\mu', \zeta_F) \right\}$$
$$\times \left(\frac{\zeta}{\mu_{b_\star}^2} \right)^{K(b_\star, \mu_{b_\star})/2} \left[\frac{\zeta}{Q_0} \right]^{-g_K(\mathbf{b}_T)/2} f_1^{NP}(x, \mathbf{b}_T; \zeta, Q_0)$$

Collins-Soper
kernel

TMD Factorization - structure of TMDs

Matching coeff.
(perturbative calculable)

Collinear PDFs
(previous fit)

Perturbative Sudakov
evolution factor

$$\hat{f}_1^q(x_B, \mathbf{b}_T; \mu_F, \zeta_F) = [C \otimes f_1](x_B, b_\star; \mu_{b_\star}, \mu_{b_\star}^2) \exp \left\{ \int_{\mu_{b_\star}}^{\mu_F} \frac{d\mu'}{\mu'} \gamma(\mu', \zeta_F) \right\}$$
$$\times \left(\frac{\zeta}{\mu_{b_\star}^2} \right)^{K(b_\star, \mu_{b_\star})/2} \left[\frac{\zeta}{Q_0} \right]^{-g_K(\mathbf{b}_T)/2} f_1^{NP}(x, \mathbf{b}_T; \zeta, Q_0)$$

Collins-Soper
kernel

NP part of
Collins-Soper Kernel

TMD Factorization - structure of TMDs

Matching coeff.
(perturbative calculable)

Collinear PDFs
(previous fit)

Perturbative Sudakov
evolution factor

$$\hat{f}_1^q(x_B, \mathbf{b}_T; \mu_F, \zeta_F) = [C \otimes f_1](x_B, b_\star; \mu_{b_\star}, \mu_{b_\star}^2) \exp \left\{ \int_{\mu_{b_\star}}^{\mu_F} \frac{d\mu'}{\mu'} \gamma(\mu', \zeta_F) \right\}$$

$$\times \left(\frac{\zeta}{\mu_{b_\star}^2} \right)^{K(b_\star, \mu_{b_\star})/2} \left[\frac{\zeta}{Q_0} \right]^{-g_K(\mathbf{b}_T)/2} f_1^{NP}(x, \mathbf{b}_T; \zeta, Q_0)$$

Collins-Soper
kernel

NP part of
Collins-Soper Kernel

Non perturbative part
of TMDs

TMD Factorization - structure of TMDs

$$\hat{f}_1^q(x_B, \mathbf{b}_T; \mu_F, \zeta_F) = [C \otimes f_1](x_B, b_\star; \mu_{b_\star}, \mu_{b_\star}^2) \exp \left\{ \int_{\mu_{b_\star}}^{\mu_F} \frac{d\mu'}{\mu'} \gamma(\mu', \zeta_F) \right\}$$

$$\times \left(\frac{\zeta}{\mu_{b_\star}^2} \right)^{K(b_\star, \mu_{b_\star})/2} \left[\frac{\zeta}{Q_0} \right]^{-g_K(\mathbf{b}_T)/2} f_1^{NP}(x, \mathbf{b}_T; \zeta, Q_0)$$

Collins-Soper
kernel

NP part of
Collins-Soper Kernel

Non perturbative part
of TMDs


Fit extraction

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points

MAPTMD22 extraction – starting point


- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points



MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points

Drell-Yan data
484



MAPTMD22 extraction – starting point


- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points

Drell-Yan data

484

SIDIS data

1547



MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **$N^3 LL^-$**

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **N^3LL^-**

Accuracy	H and C	K and γ_F	γ_K	PDFs/FFs and as evol.
LL	0	-	1	-
NLL	0	1	2	LO
NLL'	1	1	2	NLO
NNLL	1	2	3	NLO
NNLL'	2	2	3	NNLO
N^3LL^-	2	3	4	NNLO + NLO
N^3LL	2	3	4	NNLO
N^3LL'	3	3	4	N^3LO

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **N^3LL^-**

Accuracy	H and C	K and γ_F	γ_K	PDFs/FFs and as evol.
LL	0	-	1	-
NLL	0	1	2	LO
NLL'	1	1	2	NLO
NNLL	1	2	3	NLO
NNLL'	2	2	3	NNLO
N^3LL^-	2	3	4	NNLO + NLO
N^3LL	2	3	4	NNLO
N^3LL'	3	3	4	N^3LO

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **N^3LL^-**

Accuracy	H and C	K and γ_F	γ_K	PDFs/FFs and as evol.
LL	0	-	1	-
NLL	0	1	2	LO
NLL'	1	1	2	NLO
NNLL	1	2	3	NLO
NNLL'	2	2	3	NNLO
N^3LL^-	2	3	4	NNLO + NLO
N^3LL	2	3	4	NNLO
N^3LL'	3	3	4	N^3LO MMHT2014nnlo DSS14-17 NLO

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **$N^3 LL^-$**
- ***Normalization*** of SIDIS multiplicities beyond NLL

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **$N^3 LL^-$**
- ***Normalization*** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **$N^3 LL^-$**
- ***Normalization*** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**

$$f_{1\text{NP}}(x, b_T^2) \propto \text{F.T. of} \left(e^{-\frac{k_\perp^2}{g^{1A}}} + \lambda_B k_\perp^2 e^{-\frac{k_\perp^2}{g^{1B}}} + \lambda_C e^{-\frac{k_\perp^2}{g^{1C}}} \right) \quad g_1(x) = N_1 \frac{(1-x)^\alpha x^\sigma}{(1-\hat{x})^\alpha \hat{x}^\sigma}$$

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **$N^3 LL^-$**
- ***Normalization*** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**

$$f_{1\text{NP}}(x, b_T^2) \propto \text{F.T. of} \left(e^{-\frac{k_\perp^2}{g^{1A}}} + \lambda_B k_\perp^2 e^{-\frac{k_\perp^2}{g^{1B}}} + \lambda_C e^{-\frac{k_\perp^2}{g^{1C}}} \right) \quad g_1(x) = N_1 \frac{(1-x)^\alpha x^\sigma}{(1-\hat{x})^\alpha \hat{x}^\sigma}$$

$$D_{\text{NP}}(z, b_T; Q_0) = \text{F.T.} \left(e^{-P_\perp^2/g_{3A}(z)} + \lambda_F P_\perp^2 e^{-P_\perp^2/g_{3B}(z)} \right)$$

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **$N^3 LL^-$**
- **Normalization** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**

$$f_{1\text{NP}}(x, b_T^2) \propto \text{F.T. of} \left(e^{-\frac{k_\perp^2}{g^{1A}}} + \lambda_B k_\perp^2 e^{-\frac{k_\perp^2}{g^{1B}}} + \lambda_C e^{-\frac{k_\perp^2}{g^{1C}}} \right) \quad g_1(x) = N_1 \frac{(1-x)^\alpha x^\sigma}{(1-\hat{x})^\alpha \hat{x}^\sigma}$$

$$D_{\text{NP}}(z, b_T; Q_0) = \text{F.T.} \left(e^{-P_\perp^2/g_{3A}(z)} + \lambda_F P_\perp^2 e^{-P_\perp^2/g_{3B}(z)} \right)$$

$$g_K(b_T^2) = -g_2^2 \frac{b_T^2}{4}$$

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **$N^3 LL^-$**
- **Normalization** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**

11 parameters for TMD PDF
+ 1 for NP evolution + 9 for
TMD FF

$$f_{1\text{NP}}(x, b_T^2) \propto \text{F.T. of} \left(e^{-\frac{k_\perp^2}{g^{1A}}} + \lambda_B k_\perp^2 e^{-\frac{k_\perp^2}{g^{1B}}} + \lambda_C e^{-\frac{k_\perp^2}{g^{1C}}} \right) \quad g_1(x) = N_1 \frac{(1-x)^\alpha x^\sigma}{(1-\hat{x})^\alpha \hat{x}^\sigma}$$

$$D_{\text{NP}}(z, b_T; Q_0) = \text{F.T.} \left(e^{-P_\perp^2/g_{3A}(z)} + \lambda_F P_\perp^2 e^{-P_\perp^2/g_{3B}(z)} \right)$$

$$g_K(b_T^2) = -g_2^2 \frac{b_T^2}{4}$$

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **$N^3 LL^-$**
- ***Normalization*** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**

MAPTMD22 extraction – starting point

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **$N^3 LL^-$**
- ***Normalization*** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**
- Extremely good description: **$\chi^2/N_{\text{data}} = 1.06$**

MAPTMD22



MAPTMD24

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
- Perturbative accuracy: **$N^3 LL^-$**
- ***Normalization*** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**
- Extremely good description: **$\chi^2/N_{\text{data}} = 1.06$**

MAPTMD22



MAPTMD24

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
Same global data set
- Perturbative accuracy: **$N^3 LL^-$**
- ***Normalization*** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**
- Extremely good description: **$\chi^2/N_{\text{data}} = 1.06$**

MAPTMD22



MAPTMD24

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
Same global data set
- Perturbative accuracy: **$N^3 LL$**
- ***Normalization*** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**
- Extremely good description: **$\chi^2/N_{\text{data}} = 1.06$**

MAPTMD22



MAPTMD24

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points
Same global data set
- Perturbative accuracy: **$N^3 LL$** **MMHT2014nnlo**
 DSS14-17 NLO
- **Normalization** of SIDIS multiplicities beyond NLL
- Number of fitted parameters: **21**
- Extremely good description: **$\chi^2/N_{\text{data}} = 1.06$**

MAPTMD22



MAPTMD24

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points

Same global data set

- Perturbative accuracy: **$N^3 LL$**

MMHT2014nnlo
DSS14-17 NLO



NNPDF31NNLO
MAPFF10NNLO

- **Normalization** of SIDIS multiplicities beyond NLL

- Number of fitted parameters: **21**

- Extremely good description: **$\chi^2/N_{\text{data}} = 1.06$**

MAPTMD22



MAPTMD24

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points

Same global data set

- Perturbative accuracy: **$N^3 LL$**

MMHT2014nnlo
DSS14-17 NLO



NNPDF31NNLO
MAPFF10NNLO

- **Normalization** of SIDIS multiplicities beyond NLL

Same approach

- Number of fitted parameters: **21**

- Extremely good description: **$\chi^2/N_{\text{data}} = 1.06$**

MAPTMD22



MAPTMD24

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points

Same global data set

- Perturbative accuracy: **$N^3 LL$**

MMHT2014nnlo

DSS14-17 NLO



NNPDF31NNLO

MAPFF10NNLO

- **Normalization** of SIDIS multiplicities beyond NLL

Same approach

- Number of fitted parameters: **21**

Same parameterization

- Extremely good description: **$\chi^2/N_{\text{data}} = 1.06$**

MAPTMD22



MAPTMD24

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points

Same global data set

- Perturbative accuracy: **$N^3 LL$**

MMHT2014nnlo

DSS14-17 NLO



NNPDF31NNLO

MAPFF10NNLO

- **Normalization** of SIDIS multiplicities beyond NLL

Same approach

- Number of fitted parameters: **21**

Same parameterization

- Extremely good description: **$\chi^2/N_{\text{data}} = 1.40$**

MAPTMD22



MAPTMD24

- Global analysis of Drell-Yan and Semi-Inclusive DIS data sets: **2031** data points

Same global data set

- Perturbative accuracy: **$N^3 LL$** **MMHT2014nnlo** **DSS14-17 NLO** **NNPDF31NNLO** **MAPFF10NNLO**

- **Normalization** of SIDIS multiplicities beyond NLL

Same approach

- Number of fitted parameters: **21**

Same parameterization

- Extremely good description: $\chi^2/N_{\text{data}} = 1.40$

**Why does the
 χ^2 get worse?**

MAPTMD24 extraction



Collinear sets	Data set χ^2_0/N_{dat}		
	DY total	SIDIS total	Total
MMHT + DSS (MAP22)	1.66	0.87	1.06
NNPDF + MAPFF (MAP24 FI)	1.58	1.34	1.40

MAPTMD24 extraction

Collinear sets	Data set χ^2_0/N_{dat}		
	DY total	SIDIS total	Total
MMHT + DSS (MAP22)	1.66	0.87	1.06
NNPDF + MAPFF (MAP24 FI)	1.58	1.34	1.40



MAPTMD24 extraction

	Data set χ^2_0/N_{dat}		
Collinear sets	DY total	SIDIS total	Total
MMHT + DSS (MAP22)	1.66	0.87	1.06
NNPDF + MAPFF (MAP24 FI)	1.58	1.34	1.40



MAPTMD24 extraction

	Data set χ^2_0/N_{dat}		
Collinear sets	DY total	SIDIS total	Total
MMHT + DSS (MAP22)	1.66	0.87	1.06
NNPDF + MAPFF (MAP24 FI)	1.58	1.34	1.40




TMD PDFs are compatible with MAP22

MAPTMD24 extraction

Collinear sets	Data set χ^2_0/N_{dat}		
	DY total	SIDIS total	Total
MMHT + DSS (MAP22)	1.66	0.87	1.06
NNPDF + MAPFF (MAP24 FI)	1.58	1.34	1.40


MAPTMD24 extraction

	Data set χ^2_0/N_{dat}		
Collinear sets	DY total	SIDIS total	Total
MMHT + DSS (MAP22)	1.66	0.87	1.06
NNPDF + MAPFF (MAP24 FI)	1.58	1.34	1.40



MAPTMD24 extraction

	Data set χ^2/N_{dat}		
Collinear sets	DY total	SIDIS total	Total
MMHT + DSS (MAP22)	1.66	0.87	1.06
NNPDF + MAPFF (MAP24 FI)	1.58	1.34	1.40



MAPFF1.Onnlo

- Approx NNLO
 - NN approach
 - New behaviors
 - Smaller uncertainties

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

NNPDF + MAPFF

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

NNPDF + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06

MMHT + DSS (MAP22)

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

NNPDF + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.43
Dy fixed target total	233	0.75
HERMES total	344	0.95
COMPASS total	1203	0.88
SIDIS total	1547	0.90
Total	2031	1.07

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

NNPDF + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.43
Dy fixed target total	233	0.75
HERMES total	344	0.95
COMPASS total	1203	0.88
SIDIS total	1547	0.90
Total	2031	1.07

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

NNPDF + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.43
Dy fixed target total	233	0.75
HERMES total	344	0.95
COMPASS total	1203	0.88
SIDIS total	1547	0.90
Total	2031	1.07

Good agreement



Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06

NNPDF + DSS

MMHT + DSS (MAP22)

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

NNPDF + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.43
Dy fixed target total	233	0.75
HERMES total	344	0.95
COMPASS total	1203	0.88
SIDIS total	1547	0.90
Total	2031	1.07

NNPDF + DSS

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.01
Dy fixed target total	233	1.11
HERMES total	344	2.51
COMPASS total	1203	0.99
SIDIS total	1547	1.33
Total	2031	1.39

MMHT + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06



MMHT + DSS (MAP22)

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

NNPDF + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.43
Dy fixed target total	233	0.75
HERMES total	344	0.95
COMPASS total	1203	0.88
SIDIS total	1547	0.90
Total	2031	1.07

NNPDF + DSS

Good agreement



Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.01
Dy fixed target total	233	1.11
HERMES total	344	2.51
COMPASS total	1203	0.99
SIDIS total	1547	1.33
Total	2031	1.39

MMHT + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06

MMHT + DSS (MAP22)

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

NNPDF + MAPFF

Agreement
↔

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.01
Dy fixed target total	233	1.11
HERMES total	344	2.51
COMPASS total	1203	0.99
SIDIS total	1547	1.33
Total	2031	1.39

MMHT + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.43
Dy fixed target total	233	0.75
HERMES total	344	0.95
COMPASS total	1203	0.88
SIDIS total	1547	0.90
Total	2031	1.07

Good agreement
↔

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06

NNPDF + DSS

MMHT + DSS (MAP22)

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

NNPDF + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.43
Dy fixed target total	233	0.75
HERMES total	344	0.95
COMPASS total	1203	0.88
SIDIS total	1547	0.90
Total	2031	1.07

NNPDF + DSS


Agreement



Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.01
Dy fixed target total	233	1.11
HERMES total	344	2.51
COMPASS total	1203	0.99
SIDIS total	1547	1.33
Total	2031	1.39

MMHT + MAPFF

Similar
worsening



Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06

MMHT + DSS (MAP22)

Good agreement



MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

NNPDF + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.43
Dy fixed target total	233	0.75
HERMES total	344	0.95
COMPASS total	1203	0.88
SIDIS total	1547	0.90
Total	2031	1.07

Similar
worsening

MMHT + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06

NNPDF + DSS

MMHT + DSS (MAP22)

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.01
Dy fixed target total	233	1.11
HERMES total	344	2.51
COMPASS total	1203	0.99
SIDIS total	1547	1.33
Total	2031	1.39

NNPDF + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.43
Dy fixed target total	233	0.75
HERMES total	344	0.95
COMPASS total	1203	0.88
SIDIS total	1547	0.90
Total	2031	1.07

Similar
worsening

MMHT + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06

NNPDF + DSS

MMHT + DSS (MAP22)

MAPTMD24 extraction

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.14
Dy fixed target total	233	0.68
HERMES total	344	2.72
COMPASS total	1203	0.99
SIDIS total	1547	1.38
Total	2031	1.40

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.01
Dy fixed target total	233	1.11
HERMES total	344	2.51
COMPASS total	1203	0.99
SIDIS total	1547	1.33
Total	2031	1.39

NNPDF + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.43
Dy fixed target total	233	0.75
HERMES total	344	0.95
COMPASS total	1203	0.88
SIDIS total	1547	0.90
Total	2031	1.07

MMHT + MAPFF

Data set	N_{dat}	χ^2_0/N_{dat}
DY collider total	251	2.06
Dy fixed target total	233	1.24
HERMES total	344	0.71
COMPASS total	1203	0.92
SIDIS total	1547	0.87
Total	2031	1.06

NNPDF + DSS

Similar
worsening

MMHT + DSS (MAP22)

MAPTMD24 extraction

Data set	N_{dat}	χ^2/N_{dat}
DY collider to		
Dy fixed targ		
HERMES tot		
COMPASS to		
SIDIS total		
Total		

NNPDF



NNPDF + DSS

Data set	N_{dat}	χ^2/N_{dat}
251	2.01	
233	1.11	
344	2.51	
1203	0.99	
1547	1.33	
2031	1.39	

APFF

Data set	N_{dat}	χ^2/N_{dat}
251	2.06	
233	1.24	
344	0.71	
1203	0.92	
1547	0.87	
Total	2031	1.06

MMHT + DSS (MAP22)

MAPTMD24 extraction

MAPTMD24 extraction


Solution:

MAPTMD24 extraction

Solution: *Flavour dependence*

MAPTMD24 extraction

Solution: **Flavour dependence**




u, d

\bar{u}, \bar{d}

s (*sea*)

MAPTMD24 extraction

Solution: **Flavour dependence**



u, d

\bar{u}, \bar{d}


s (*sea*)

$u \rightarrow \pi^+, \dots$

$d \rightarrow \pi^+, \dots$

MAPTMD24 extraction

Solution: **Flavour dependence**



u, d

\bar{u}, \bar{d}

s (*sea*)

$u \rightarrow \pi^+, \dots$

$d \rightarrow \pi^+, \dots$


$u \rightarrow K^+, \dots$

$\bar{s} \rightarrow K^+, \dots$

$d \rightarrow K^+, \dots$

MAPTMD24 extraction

Solution: **Flavour dependence**



u, d

\bar{u}, \bar{d}

s (*sea*)

$u \rightarrow \pi^+, \dots$

$d \rightarrow \pi^+, \dots$

$u \rightarrow K^+, \dots$

$\bar{s} \rightarrow K^+, \dots$

$d \rightarrow K^+, \dots$

Negative fragmenting mesons: charge conjugation

MAPTMD24 extraction

HERMES

MAPTMD24 extraction

$$e + p \rightarrow e' + \pi^+ + X$$

$$e + p \rightarrow e' + \pi^- + X$$





$$e + p \rightarrow e' + K^+ + X$$

$$e + p \rightarrow e' + K^- + X$$

+ deuteron target

HERMES

MAPTMD24 extraction




+ deuteron target

high sensitivity to flavour dependence

MAPTMD24 extraction

HERMES



+ deuteron target


high sensitivity to flavour dependence

COMPASS

deuteron target & unidentified final state hadron

MAPTMD24 extraction

HERMES



+ deuteron target

high sensitivity to flavour dependence


COMPASS

deuteron target & unidentified final state hadron

low sensitivity to flavour dependence

MAPTMD24 extraction

HERMES



+ deuteron target

high sensitivity to flavour dependence

COMPASS

deuteron target & unidentified final state hadron


low sensitivity to flavour dependence

Drell-Yan

$q\bar{q}$ in the initial state

MAPTMD24 extraction

HERMES



high sensitivity to flavour dependence

COMPASS

deuteron target & unidentified final state hadron

low sensitivity to flavour dependence

Drell-Yan

$q\bar{q}$ in the initial state


lowest sensitivity to flavour dependence

MAPTMD24 extraction - Results $\chi^2/N_{data} = 1.08$

Data set	N ³ LL			
	N_{dat}	χ^2_D	χ^2_λ	χ^2_0
DY collider total	251	1.37	0.28	1.65
DY fixed-target total	233	0.63	0.31	0.94
<i>HERMES total</i>	344	0.81	0.24	1.05
<i>COMPASS total</i>	1203	0.67	0.27	0.94
SIDIS total	1547	0.70	0.26	0.96
Total	2031	0.81	0.27	1.08


MAPTMD24 extraction - Results $\chi^2/N_{data} = 1.08$

N ³ LL				
Data set	N_{dat}	χ^2_D	χ^2_λ	χ^2_0
DY collider total	251	1.37	0.28	1.65
DY fixed-target total	233	0.63	0.31	0.94
<i>HERMES total</i>	344	0.81	0.24	1.05
<i>COMPASS total</i>	1203	0.67	0.27	0.94
SIDIS total	1547	0.70	0.26	0.96
Total	2031	0.81	0.27	1.08



MAPTMD24 extraction - Results $\chi^2/N_{data} = 1.08$


N ³ LL				
Data set	N_{dat}	χ^2_D	χ^2_λ	χ^2_0
DY collider total	251	1.37	0.28	1.65
DY fixed-target total	233	0.63	0.31	0.94
<i>HERMES total</i>	344	0.81	0.24	1.05
<i>COMPASS total</i>	1203	0.67	0.27	0.94
SIDIS total	1547	0.70	0.26	0.96
Total	2031	0.81	0.27	1.08



MAPTMD24 extraction - Results $\chi^2/N_{data} = 1.08$

N ³ LL				
Data set	N_{dat}	χ^2_D	χ^2_λ	χ^2_0
DY collider total	251	1.37	0.28	1.65
DY fixed-target total	233	0.63	0.31	0.94
<i>HERMES total</i>	344	0.81	0.24	1.05
<i>COMPASS total</i>	1203	0.67	0.27	0.94
SIDIS total	1547	0.70	0.26	0.96
Total	2031	0.81	0.27	1.08


Very good agreement



MAPTMD24 extraction - Results $\chi^2/N_{data} = 1.08$

N ³ LL				
Data set	N_{dat}	χ^2_D	χ^2_λ	χ^2_0
DY collider total	251	1.37	0.28	1.65
DY fixed-target total	233	0.63	0.31	0.94
<i>HERMES total</i>	344	0.81	0.24	1.05
<i>COMPASS total</i>	1203	0.67	0.27	0.94
SIDIS total	1547	0.70	0.26	0.96
Total	2031	0.81	0.27	1.08


Very good agreement



MAPTMD24 extraction - Results $\chi^2/N_{data} = 1.08$

N ³ LL				
Data set	N_{dat}	χ^2_D	χ^2_λ	χ^2_0
DY collider total	251	1.37	0.28	1.65
DY fixed-target total	233	0.63	0.31	0.94
<i>HERMES total</i>	344	0.81	0.24	1.05
<i>COMPASS total</i>	1203	0.67	0.27	0.94
SIDIS total	1547	0.70	0.26	0.96
Total	2031	0.81	0.27	1.08

Very good agreement




MAPTMD24 extraction - Results $\chi^2/N_{data} = 1.08$

N ³ LL				
Data set	N_{dat}	χ^2_D	χ^2_λ	χ^2_0
DY collider total	251	1.37	0.28	1.65
DY fixed-target total	233	0.63	0.31	0.94
<i>HERMES total</i>	344	0.81	0.24	1.05
<i>COMPASS total</i>	1203	0.67	0.27	0.94
SIDIS total	1547	0.70	0.26	0.96
Total	2031	0.81	0.27	1.08

Very good agreement

Quite good agreement




MAPTMD24 extraction - Results $\chi^2/N_{data} = 1.08$


Data set	N ³ LL			
	N_{dat}	χ^2_D	χ^2_λ	χ^2_0
DY collider total	251	1.37	0.28	1.65
DY fixed-target total	233	0.63	0.31	0.94
<i>HERMES total</i>	344	0.81	0.24	1.05
<i>COMPASS total</i>	1203	0.67	0.27	0.94
SIDIS total	1547	0.70	0.26	0.96
Total	2031	0.81	0.27	1.08

Very good agreement


Quite good agreement




MAPTMD24 extraction - TMD PDFs




MAPTMD24 extraction - TMD PDFs



MAPTMD24 extraction - TMD PDFs




MAPTMD24 extraction - TMD PDFs



Very different k_\perp - behaviours!


MAPTMD24 extraction - TMD PDFs



Very different k_\perp - behaviours!

It changes also by varying x

MAPTMD24 extraction - TMD PDFs




Very different k_\perp - behaviours!

It changes also by varying x

MAPTMD24 extraction - TMD PDFs

The sea is the least constrained




Very different k_{\perp} - behaviours!

It changes also by varying x

MAPTMD24 extraction - TMD PDFs

The sea is the least constrained




Very different k_{\perp} - behaviours!

It changes also by varying x

MAPTMD24 extraction - TMD PDFs

The sea is the least constrained


The up quark is the most one




Very different k_{\perp} - behaviours!

It changes also by varying x


MAPTMD24 extraction - TMD FFs




MAPTMD24 extraction - TMD FFs



MAPTMD24 extraction - TMD FFs





MAPTMD24 extraction - TMD FFs



Some signals of differences between favoured and unfavoured channels



MAPTMD24 extraction - TMD FFs

The favoured is better constrained than the unfavoured one





Some signals of differences between favoured and unfavoured channels



MAPTMD24 extraction - TMD FFs



MAPTMD24 extraction - TMD FFs




MAPTMD24 extraction - TMD FFs



Strong differences between different hadron fragmentations!

MAPTMD24 extraction - Scatter plots



Conclusions and outlook

Conclusions and outlook

- **MAPTMD24** is the *first flavour dependent* extraction of unpolarized quarks TMDs in the proton from a *global* fit

Conclusions and outlook

- **MAPTMD24** is the *first flavour dependent* extraction of unpolarized quarks TMDs in the proton from a *global* fit
- We find ***significant*** differences between the flavors in the ***TMD PDFs***.

Conclusions and outlook

- **MAPTMD24** is the *first flavour dependent* extraction of unpolarized quarks TMDs in the proton from a *global* fit
- We find ***significant*** differences between the flavors in the ***TMD PDFs***.
- We find ***significant*** differences between different final hadrons in the ***TMD FFs***.

Conclusions and outlook


- **MAPTMD24** is the *first flavour dependent* extraction of unpolarized quarks TMDs in the proton from a *global* fit
- We find ***significant*** differences between the flavors in the ***TMD PDFs***.
- We find ***significant*** differences between different final hadrons in the ***TMD FFs***.
- We find a weak signal between different flavors in the same final hadron.

BACKUP

MAPTMD22: Normalization of SIDIS


MAPTMD22: Normalization of SIDIS

High Energy Drell-Yan



MAPTMD22: Normalization of SIDIS


High Energy Drell-Yan




The description improves at high orders

MAPTMD22: Normalization of SIDIS

High Energy Drell-Yan



SIDIS




The description improves at high orders

Strange behaviors at higher orders


MAPTMD22: Normalization of SIDIS

COMPASS multiplicities (one of many bins)




MAPTMD22: Normalization of SIDIS

COMPASS multiplicities (one of many bins)




MAPTMD22: Normalization of SIDIS

COMPASS multiplicities (one of many bins)



MAPTMD22: Normalization of SIDIS

COMPASS multiplicities (one of many bins)



For different orders the discrepancy amounts to a nearly constant factor

MAPTMD22: Normalization of SIDIS

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} \cancel{dP}_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$


MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ dz dP_{hT}} \Bigg/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ dz}$$




MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ dz dP_{hT}} \Bigg/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ dz}$$



MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$

$$\frac{d\sigma}{dx dQ \cancel{dz}} = \int \cancel{dP_{hT}} \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$

$$\frac{d\sigma}{dx dQ \cancel{dz}} = \int \cancel{dP_{hT}} \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \stackrel{?}{=} \int \cancel{dP_{hT}} \quad W\text{-term}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$

$$\frac{d\sigma}{dx dQ \cancel{dz}} = \int \cancel{dP_{hT}} \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \stackrel{?}{=} \int \cancel{dP_{hT}} \quad W\text{-term}$$

$$\left. \frac{d\sigma}{dx dz dQ} \right|_{\text{LO}} = \simeq \int dq_T W \Big|_{\text{NLL}} \propto f_1^q(x, Q) D_1^{q \rightarrow h}(z, Q)$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$

$$\frac{d\sigma}{dx dQ \cancel{dz}} = \int \cancel{dP_{hT}} \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \stackrel{?}{=} \int \cancel{dP_{hT}} \quad W\text{-term}$$

$$\left. \frac{d\sigma}{dx dz dQ} \right|_{\text{LO}} = \simeq \int dq_T W \Big|_{\text{NLL}} \propto f_1^q(x, Q) D_1^{q \rightarrow h}(z, Q)$$

$$\int dq_T W \Big|_{\text{NNLL}} \neq \left. \frac{d\sigma}{dx dz dQ} \right|_{\text{NLO}}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$

$$\frac{d\sigma}{dx dQ \cancel{dz}} = \int \cancel{dP_{hT}} \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \stackrel{?}{=} \int \cancel{dP_{hT}} \quad W\text{-term}$$

$$\left. \frac{d\sigma}{dx dz dQ} \right|_{\text{LO}} = \left. \int dq_T W \right|_{\text{NLL}} \propto f_1^q(x, Q) D_1^{q \rightarrow h}(z, Q)$$

At higher orders
something is missing
(Y-term? Power corrections?)

$$\left. \int dq_T W \right|_{\text{NNLL}} \neq \left. \frac{d\sigma}{dx dz dQ} \right|_{\text{NLO}}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Big/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$

$$\int dP_{hT} \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} = \frac{d\sigma}{dx dQ \cancel{dz}}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ \cancel{dz} dP_{hT}} \Bigg/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ \cancel{dz}}$$

$$\int dP_{hT} \frac{d\sigma}{dx dQ dz dP_{hT}} = \frac{d\sigma}{dx dQ dz}$$

$$w(x, z, Q) = \frac{d\sigma}{dx dQ \cancel{dz}} \Bigg/ \int dP_{hT} \frac{d\sigma}{dx dQ dz dP_{hT}}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ dz dP_{hT}} \Bigg/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ dz}$$

$$\int dP_{hT} \frac{d\sigma}{dx dQ dz dP_{hT}} = \frac{d\sigma}{dx dQ dz}$$

$$w(x, z, Q) = \frac{d\sigma}{dx dQ dz} \Bigg/ \int dP_{hT} \frac{d\sigma}{dx dQ dz dP_{hT}}$$

$$M(x, z, P_{hT}, Q) = w(x, z, Q) \frac{d\sigma}{dx dQ dz dP_{hT}} \Bigg/ \frac{d\sigma}{dx dQ}$$

MAPTMD22: Normalization of SIDIS

SIDIS multiplicity

$$M(x, z, P_{hT}, Q) = \frac{d\sigma}{dx dQ dz dP_{hT}} \Bigg/ \frac{d\sigma}{dx dQ}$$

Collinear SIDIS cross section

$$\frac{d\sigma}{dx dQ dz}$$

$$\int dP_{hT} \frac{d\sigma}{dx dQ dz dP_{hT}} = \frac{d\sigma}{dx dQ dz}$$

***Fitting parameters
independent***

$$w(x, z, Q) = \frac{d\sigma}{dx dQ dz} \Bigg/ \int dP_{hT} \frac{d\sigma}{dx dQ dz dP_{hT}}$$

$$M(x, z, P_{hT}, Q) = w(x, z, Q) \frac{d\sigma}{dx dQ dz dP_{hT}} \Bigg/ \frac{d\sigma}{dx dQ}$$

BACKUP

Data set	N ³ LL			
	N_{dat}	χ^2_D	χ^2_λ	χ^2_0
<i>Tevatron total</i>	71	1.10	0.07	1.17
<i>LHCb total</i>	21	3.56	0.96	4.52
<i>ATLAS total</i>	72	3.54	0.82	4.36
<i>CMS total</i>	78	0.38	0.05	0.43
PHENIX 200	2	2.76	1.04	3.80
STAR 510	7	1.12	0.26	1.38
DY collider total	251	1.37	0.28	1.65
E288 200 GeV	30	0.13	0.40	0.53
E288 300 GeV	39	0.16	0.26	0.42
E288 400 GeV	61	0.11	0.08	0.19
E772	53	0.88	0.20	1.08
E605	50	0.70	0.22	0.92
DY fixed-target total	233	0.63	0.31	0.94
<i>HERMES total</i>	344	0.81	0.24	1.05
<i>COMPASS total</i>	1203	0.67	0.27	0.94
SIDIS total	1547	0.70	0.26	0.96
Total	2031	0.81	0.27	1.08

BACKUP


Error propagation



100 Monte Carlo
replicas of data

100 Monte Carlo
replicas of PDFs

100 Monte Carlo
replicas of FFs



BACKUP

Kinematic power corrections in TMD factorization theorem

#3

Alexey Vladimirov (Madrid U.) (Jul 24, 2023)

Published in: *JHEP* 12 (2023) 008 • e-Print: [2307.13054](#) [hep-ph]


 pdf

 DOI

 cite


 claim

 reference search


 6 citations

Estimations made in sec. 5.3 demonstrate that including KPCs results in an almost constant increment of the cross-section. The magnitude of this correction depends on Q and x . For typical LHC kinematics, the correction is around 1%, while at $Q \sim 4 - 5$ GeV, the correction can reach 100%. Interestingly, the deficiency in normalization for the TMD factorization at low energies has been reported by multiple groups. One could expect that these problems will be resolved with the inclusion of KPCs.


MAPTMD24 extraction - Collins Soper Kernel



MAPTMD24 extraction - Collins Soper Kernel




MAPTMD24 extraction - Collins Soper Kernel



*Independent of our non
perturbative choices*


MAPTMD24 extraction - Collins Soper Kernel



Independent of our non perturbative choices

Quite flat behaviour

MAPTMD24 extraction - Collins Soper Kernel



Independent of our non perturbative choices

Quite flat behaviour

Compatible with latest lattice calculation

PLB 852 (2024) 138617