

Superfluid fraction in the inner crust of neutron stars

Giorgio Almirante


IJCLab, Orsay, France




Outline

- Introduction: inner crust of neutron stars and its superfluid fraction
- Formalism: Hartre-Fock-Bogoliubov and Andreev-Bashkin two fluid model
- Results: a whole lasagna and some spaghetti

Inner crust of neutron stars...




W.G. Newton et al,
Sym.En.,In.Crust,Gl.Mod. (2011)




H. Dinh Thi et al,
A&A 654, A114 (2021)

...and its superfluid fraction



Radhakrishnan & Manchester,
Nature 222, 228-229 (1969)



Martin & Urban,
Phys. Rev. C 94, 065801 (2016)

Hartree-Fock-Bogoliubov in the pasta phases

$$\begin{pmatrix} h - \mu & -\Delta \\ -\Delta^\dagger & -\bar{h} + \mu \end{pmatrix} \begin{pmatrix} U_\alpha^* \\ -V_\alpha \end{pmatrix} = E_\alpha \begin{pmatrix} U_\alpha^* \\ -V_\alpha \end{pmatrix}$$

in momentum space only integer indices, diagonal in the Bloch and parallel momenta

$$h_{kk'} = \left(\frac{\hbar^2}{2m^*} \right)_{kk'} k \cdot k' + U_{kk'} - k \cdot v \delta_{kk'} \quad (1)$$

$$U_{kk'} = - \sum_{pp'} V_{kp k' p'} \rho_{p' p} \rightarrow \text{Skyrme potential} \quad (2)$$

$$\Delta_{kk'} = - \sum_{pp'} V_{kk' p' p} \kappa_{p' p} \rightarrow \text{separable interaction} \quad (3)$$

Andreev-Bashkin formalism

Our setup gets us access to densities, currents and other relevant quantities. Then with the formalism due to Andreev and Bashkin we can compute the superfluid fraction

$$\vec{\rho}_n = (\rho_n - \rho_S)\vec{v} + \rho_S \vec{V}_n \quad ; \quad \vec{\rho}_p = \rho_p \vec{v}$$



This relation has to be understood in an average sense

$$\vec{V}_n = \int_V \frac{dx^3}{V} \frac{\hbar}{2m} \vec{\nabla} \phi \quad \text{where} \quad \Delta(q, x) = |\Delta(q, x)| e^{i\phi(x)}$$

$$(V_n)_x = \iint \frac{dydz}{V} \frac{\hbar}{2m} \left(\phi(L_x, y, z) - \phi(0, y, z) \right)$$



Since our quantities are periodic the average superfluid velocity is zero, thus we are working in the reference frame in which the superfluid component carries no momentum

Lasagna statics



Almirante & Urban, Phys. Rev. C 109, 045805 (2024)



Lasagna dynamics



Almirante & Urban, Phys. Rev. C 109, 045805 (2024)



Spaghetti square

$$\rho_b = 0.062 \text{ fm}^{-3} ; \rho_S/\rho_n = 0.94$$



Spaghetti hexagon

$$\rho_b = 0.062 \text{ fm}^{-3} ; \rho_S/\rho_n = 0.94$$



Results summary and comparison

Spaghetti

μ_n (MeV)	L (fm)	ρ_b (fm ⁻³)	ρ_S/ρ_n (our%)	ρ_S/ρ_n (Chamel%)
12	24	0.0619	94.5	75
	28	0.0617	95.7	
12.5	24	0.0670	95.4	82
	28	0.0668	96.7	

Lasagna

μ_n (MeV)	L (fm)	ρ_b (fm ⁻³)	ρ_S/ρ_n (our%)	ρ_S/ρ_n (Chamel%)
13	20	0.0723	96.3	93
	24	0.0720	96.2	
13.5	20	0.0768	97.2	94
	24	0.0766	97.1	

Thanks for your attention!

