

Rencontre du groupe de travail ”méthodes d’analyse des données” du GdR Ondes Gravitationnelles Description

Rapport sur les contributions

ID de Contribution: 1

Type: Non spécifié

A test for LISA foregrounds Gaussianity and stationarity. II. Extreme-mass-ratio inspirals

mercredi 16 octobre 2024 12:00 (20 minutes)

Extreme Mass Ratio Inspirals (EMRIs) are key targets expected to be observed by the Laser Interferometer Space Antenna (LISA) mission. Unresolvable EMRI signals contribute to forming a gravitational wave background (GWB).

Characterizing the statistical features of the GWB from EMRIs is of great importance, as EMRIs will ubiquitously affect large segments of the inference scheme.

In this work, we apply a frequentist test for GWB Gaussianity and stationarity, exploring three astrophysically-motivated EMRI populations. We construct the resulting signal by combining state-of-the-art EMRI waveforms and a detailed description of the LISA response with first-generation time-delay interferometry variables.

Depending on the brightness of the GWB, our analysis demonstrates that the resultant EMRI foregrounds show varying degrees of departure from the usual statistical assumptions that the GWBs are both Gaussian and Stationary.

If the GWB background is non-stationary with non-Gaussian features, this will challenge the robustness of Gaussian-likelihood model, when applied to global inference results, e.g. foreground estimation, background detection, and individual-source parameters reconstruction.

Auteur principal: PIARULLI, Manuel (L2IT, Université Toulouse III - Paul Sabatier)

Co-auteurs: SESANA, Alberto (Università degli Studi di Milano-Bicocca); POZZOLI, Federico (Università degli Studi Milano-Bicocca); M. BONETTI, Matteo (Università degli Studi di Milano-Bicocca); BURKE, Ollie (L2I Toulouse, CNRS/IN2P3, UT3); Dr BUSCICCHIO, Riccardo (Università di Milano-Bicocca)

Orateur: PIARULLI, Manuel (L2IT, Université Toulouse III - Paul Sabatier)

Classification de Session: Contributed talks

Reconstructing gravitational wave polarizations with bivariate signal processing and deep generative models

mercredi 16 octobre 2024 10:45 (45 minutes)

This work addresses the inverse problem of the reconstruction of the impinging gravitational-wave (GW) polarizations (h_+, h_\times) from the interferometric detector measurements. It is specifically focused on situations where the polarization pattern is non-stationary and evolves over the duration of the signal. For instance, this is the case for GW signals from compact binaries coalescence (CBC) subject to precession of their orbital plane; precession may be due to misaligned component spins. The GWTC-3 catalog (LIGO Scientific Collaboration et al., 2023) includes several such binaries like GW200129\065458.

Two new approaches are proposed that improve upon the methods available in the literature such as BayesWave (Cornish and Littenberg, 2015). BayesWave defines the GW signal model as a combination of wavelets with independent or constant elliptical polarization. It is therefore unable to capture both polarizations correctly or accurately.

1. The first approach builds on a previous work (Cano, 2022), that is based on the geometrical parametrization of the polarization content through Stokes parameters (Flamant, 2018). The regularization of the inversion is performed using a term to penalize the misalignment between reconstructed and predefined target Stokes parameters. This earlier work is restricted to constant Stokes parameters and suffers for the same limitations as BayesWave. The new approach enforces a variable but smooth evolution of the signal Stokes parameters in the reconstructed polarizations (Pilavci et al., 2024).
2. The second approach is based on the plug-and-play (PnP) method (Hurault, 2023) that exploits a deep generative model. PnP methods constitute the current state-of-the-art in image processing and consist in learning a prior from image examples. We apply this technique here on spectrograms of GW simulated signals.

The two proposed approaches are complementary. They resort to different levels of priors for the regularization. These priors are exploited in different ways in practice. The first approach is rather model-agnostic and encodes desirable characteristics of GW signals such as regularity or smoothness of the polarization evolution. In contrast, the second approach ensures that reconstructed GWs belong to the true GW spectrogram distribution learnt from the numerical model.

References

- Cano, Cyril (2022). “Mathematical tools and signal processing algorithms for the study of gravitational waves polarization”. PhD thesis. Université Grenoble Alpes.
- The LIGO Scientific Collaboration et al. (2023). “GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run”. Phys. Rev. X 13.4, p. 041039.
- Cornish, Neil J. and Tyson B. Littenberg (2015). “Bayeswave: Bayesian inference for gravitational wave bursts and instrument glitches”. Class. Quantum Grav. 32.13, p. 135012.
- Flamant, Julien (2018). “A general approach for the analysis and filtering of bivariate signals”. PhD thesis. Ecole Centrale de Lille.
- Hurault, Samuel (2023). “Convergent plug-and-play methods for image inverse problems with explicit and nonconvex deep regularization”. PhD thesis. Université de Bordeaux.

Pilavci, Yusuf Yigit et al. (2024). "Denoising bivariate signals via smoothing and polarization priors "(European Signal Processing Conference (EUSIPCO)).

Auteurs principaux: PALUD, Pierre (APC); PILAVCI, Yusuf yigit; CHASSANDE-MOTTIN, Eric (CNRS AstroParticule et Cosmologie); CHAINAIS, Pierre (Centrale Lille - CRISTAL)

Orateurs: PALUD, Pierre (APC); PILAVCI, Yusuf yigit

Classification de Session: Contributed talks

ID de Contribution: 5

Type: Non spécifié

Addressing gaps in LISA data

mercredi 16 octobre 2024 13:30 (30 minutes)

In this talk, we will discuss the impact of data gaps on parameter estimation in the context of the Laser Interferometer Space Antennae (LISA). Data gaps, for LISA, are unavoidable: whether it is due to antennae repointing due to drift in the orbit, or instrumental malfunctions that are then masked, it is paramount that we can account for data gaps in our parameter estimation pipelines. In this talk, we will discuss a number of general methods that can be used to account for data gaps in LISA data. We will discuss advantages and disadvantages for dealing with data gaps in the frequency domain, time domain and, if time allows, the time-frequency domain.

Auteur principal: MARSAT, Sylvain (L2I Toulouse, CNRS/IN2P3, UT3)

Co-auteur: BURKE, Ollie (L2I Toulouse, CNRS/IN2P3, UT3)

Orateurs: BURKE, Ollie (L2I Toulouse, CNRS/IN2P3, UT3); MARSAT, Sylvain (L2I Toulouse, CNRS/IN2P3, UT3)

Classification de Session: Contributed talks

ID de Contribution: 6

Type: Non spécifié

Inferring astrophysics and cosmology with individual compact binary coalescences and their gravitational-wave stochastic background

mercredi 16 octobre 2024 11:30 (30 minutes)

This work introduces a method to infer the Hubble constant H_0 by combining dark siren gravitational wave sources (without electromagnetic counterparts) with the stochastic gravitational wave background (SGWB). Traditional H_0 measurement techniques, such as the local distance ladder and cosmic microwave background observations, face significant challenges and yield conflicting results. Gravitational Waves dark sirens can measure the Hubble constant by using a calibration given by the source mass spectrum. The proposed framework integrates SGWB data, which contains signals from numerous unresolved sources, to determine the mass spectrum and hence H_0 . This method leverages complementary information from both sources. Although preliminary analysis has not shown yet a significant improvement in the H_0 precision with projected O5 sensitivity, considering also the other population parameters unknown might result in an improvement.

Auteur principal: FERRAIUOLO, Sarah (CPPM at Aix-Marseille University (AMU) & Università di Roma, La Sapienza)

Orateur: FERRAIUOLO, Sarah (CPPM at Aix-Marseille University (AMU) & Università di Roma, La Sapienza)

Classification de Session: Contributed talks

ID de Contribution: 7

Type: **Non spécifié**

LISA data analysis

mercredi 16 octobre 2024 09:00 (30 minutes)

Status of LISA data analysis

Orateur: PETITEAU, Antoine (CEA/IRFU/DPhP)

Classification de Session: Reviews

ID de Contribution: **8**

Type: **Non spécifié**

PTA data analysis (remote)

mercredi 16 octobre 2024 09:30 (30 minutes)

Status of PTA data analysis

Orateur: QUELQUEJAY-LECLÈRE, Hippolyte (APC)

Classification de Session: Reviews

Matched Filtering

mercredi 16 octobre 2024 10:00 (30 minutes)

Orateur: DAL CANTON, Tito (UPSAclay, CNRS/IN2P3, IJCLab)

Classification de Session: Reviews

Rencontre du gro... / Rapport sur les contributions

Population inference

ID de Contribution: **10**

Type: **Non spécifié**

Population inference

mercredi 16 octobre 2024 14:00 (30 minutes)

Auteur principal: TOUBIANA, Alexandre (Max Planck Institute for Gravitational Physics)

Orateur: TOUBIANA, Alexandre (Max Planck Institute for Gravitational Physics)

Classification de Session: Contributed talks