

Heavy Sterile Neutrinos in B Decays and new QCD Corrections to their semi-hadronic Decay Rates

Tim Kretz

Based on work with

Florian Bernlochner, Marco Fedele, Ulrich Nierste and Markus T. Prim

JHEP 01 (2025) 040 [2410.11945]

EPS-HEP 2025 - Marseille 10.07.2025

1. Overview

2. Sterile Neutrinos from $B \rightarrow D^* \ell N$

- Theory Description
- Belle II Measurements
- Results

3. Hadronic Sterile Neutrino Decay

- Tree-Level
- QCD Corrections
- Results

1. Overview

- A **Sterile Neutrino = Heavy Neutral Lepton (HNL)** arises in many NP models on e.g. Dark Matter, ν Oscillations and baryon asymmetry (see e.g. Bodarenko et al. [1805.08567](#))
- Most commonly sterile neutrino interactions are parametrised by a mixing parameter i.e. $V_{N\alpha}$

$$\mathcal{L}_I = \frac{g V_{N\alpha}}{\sqrt{2}} W_\mu^+ \bar{N}^c \gamma^\mu P_L \ell_\alpha^- + \frac{g V_{N\alpha}}{\cos \theta_w} Z_\mu \bar{N}^c \gamma^\mu P_L \nu_\alpha + \text{h.c.}$$

with **weak coupling** g and **weak mixing angle** θ_w and $P_L = (1 - \gamma_5)/2$


- $B \rightarrow D^* \ell N$ deviates from mixing parameter description!

2. Sterile Neutrinos from $B \rightarrow D^* \ell N$

- Idea: Data sample on Standard Model (SM) process $B \rightarrow D^* \ell \nu$ could contain a new-physics (NP) contribution $B \rightarrow D^* \ell N$
- SM and NP decay are 4-body decays: $B \rightarrow D^* [\rightarrow D\pi] \ell \nu$ with $\ell = e, \mu$
- The SM decay was investigated by Belle II
→ access to angular distributions!
- SM described by dimension-6 Fermi operator

$$\mathcal{O}^{(6)} = \bar{c}_L \gamma_\mu b_L \bar{\ell}_L \gamma^\mu \nu_{\ell, L}$$

Angles of the decay distribution



graphic taken from Bečirević et al., [1907.02257](#)

Differential Decay Rate of $B \rightarrow D^* \ell \nu$

$$\frac{32\pi}{9} \frac{d^4\Gamma}{dq^2 d\cos\theta_\ell d\cos\theta_D d\chi} = (J_{1s} + J_{2s} \cos 2\theta_\ell + J_{6s} \cos \theta_\ell) \sin^2 \theta_D +$$
$$(J_{1c} + J_{2c} \cos 2\theta_\ell + J_{6c} \cos \theta_\ell) \cos^2 \theta_D +$$
$$(J_3 \cos 2\chi + J_9 \sin 2\chi) \sin^2 \theta_D \sin^2 \theta_\ell +$$
$$(J_4 \cos \chi + J_8 \sin \chi) \sin 2\theta_D \sin 2\theta_\ell +$$
$$(J_5 \cos \chi + J_7 \sin \chi) \sin 2\theta_D \sin 2\theta_\ell$$

➡ J_i coefficients measurable in experiment!

N new physics contribution to B decay

- SM and NP sum incoherently $J_i = J_i^{SM} + J_i^{NP}(\{g_j\})$
- Sterile neutrinos described by four energy dimension-6 operators

$$\mathcal{H}_{\text{eff}} = \frac{4G_F}{\sqrt{2}} V_{cb} \left[(\bar{c}_L \gamma_\mu b_L) (\bar{\ell}_L \gamma^\mu \nu_{\ell,L}) + g_{V_R}^N (\bar{c}_R \gamma_\mu b_R) (\bar{\ell}_R \gamma^\mu N_R) + g_{S_L}^N (\bar{c}_R b_L) (\bar{\ell}_L N_R) \right. \\ \left. + g_{S_R}^N (\bar{c}_L b_R) (\bar{\ell}_L N_R) + g_T^N (\bar{c}_L \sigma_{\mu\nu} b_R) (\bar{\ell}_L \sigma^{\mu\nu} N_R) + \text{h.c.} \right]$$

Robinson, Shakya and Zupan, [1807.04753](#)

N new physics contribution to B decay


- Higher energy dimension operators are neglected
- E.g. operator with left-handed quarks is dimension-8

$$\mathcal{O}_{V_L} = (\bar{Q}_L \tilde{H} \gamma_\mu H^\dagger Q_L) (\bar{\ell}_R \gamma_\mu N_R)$$

- This description is a deviation from the mixing angle description. In terms of quark operators a mixing angle would be energy dimension-7

Belle II Measurements

- Belle II measured these J_i coefficients:



Hadronic recoil parameter:


$$w = \frac{m_B^2 + m_{D^*}^2 - q^2}{2m_B m_{D^*}}$$

Normalized angular coefficient:

$$\hat{J}_i^{(n)} = \frac{\int_{\Delta w^{(n)}} dw J_i(w)}{\int_{w_{\min}}^{w_{\max}} dw \frac{d\Gamma}{dw}}$$

Prim et al., [2310.20286](#)

Belle II Measurements



Prim et al., 2310.20286

Parameter Analysis with decay distributions from Belle II


Bernlochner, Fedele, TK, Nierste, Prim [2410.11945]:

- We have fitted the J_i to the recent Belle II data
- Bayesian analysis, fitted parameters: (g_j^N, m_N, FF) . Two scenarios: One non-zero WC and varying all WC at the same time.
- Analysis performed for both $\ell = e$ and $\ell = \mu$
- Fit insensitive to choice of form factors (FNAL/MILC, JLQCD, ...)

Results



- Angular coefficients only sensitive for $m_N \leq 62.5 \text{ MeV}$

$$\ell = e$$





Results

$$\ell = e$$





Results

$$\ell = \mu$$




Results

$$\ell = \mu$$




Results

$$\ell = e$$




$$\ell = \mu$$




Results

- Hint at sterile neutrino with a mass of $m_N = 354 \text{ MeV}$





Results



3. Hadronic Sterile Neutrino Decays


TK, Nierste 2025:

- Return to **mixing angle** description
- Sterile neutrinos with $m_N > 1 \text{ GeV}$ can decay into leptons and into hadrons $\pi^0, \pi^\pm, K^\pm, D^\pm \dots$ via the weak gauge bosons W^\pm, Z
- At tree-level this is known



Multi-Hadron Final States

- While tree-level results for $N \rightarrow \ell q \bar{q}$ and $N \rightarrow \nu q \bar{q}$ are well known, decay rates into multi-hadron final states (i.e. $N \rightarrow \ell \pi \pi \pi$) and QCD corrections are completely unknown.
- In principle these contributions could be sizeable
- QCD behaviour is fully governed by W^*, Z^* couplings to quarks
- W^*, Z^* correlation functions have already been calculated [1] up to $\mathcal{O}(\alpha_s^4)$!



[1] see e.g. Baikov, Chetyrkin and Kühn, [0801.1821](#)

Fully Inclusive Decay Rate $N \rightarrow \ell \text{ had.}$

- Fully inclusive decay width calculable using the **gauge boson correlator**
- Neutral current decay similar albeit more subtle due to triangles!

$$\Gamma(N \rightarrow \ell \text{ had.}) = \frac{G_F^2 m_N^5 |V_{N\ell}|^2}{192\pi^3} 12\pi \times \int_0^{(1-x_\ell)^2} dx \sqrt{\lambda(1, x, x_\ell^2)} \left((1-x+x_\ell^2)(1+2x+x_\ell^2) - 4x_\ell^2 \right) \Im(\Pi^{(1+0)}(m_N^2 x))$$


here: $x_\ell = \frac{m_\ell}{m_N}$

QCD Correlator

- Here $\Pi^{(1+0)}(s) = \Pi^{(1)}(s) + \Pi^{(0)}(s)$ is the sum of the transversal and longitudinal part of the correlator ($s = q^2$):

$$\Pi_{\mu\nu,ij}^{V/A} = (-g_{\mu\nu}q^2 + q_\mu q_\nu)\Pi_{ij,V/A}^{(1)}(s) + q_\mu q_\nu \Pi_{ij,V/A}^{(0)}(s)$$

- Corresponds to QCD corrections to W like





Results

- We calculated the fully inclusive decay width up to $\mathcal{O}(\alpha_s^4)$ in chiral limit $m_q = 0$ for the **charged current**
- Up to $\mathcal{O}(\alpha_s^3)$ fully analytical
- At $\mathcal{O}(\alpha_s^4)$ semi-analytical
- We can estimate the stability of the perturbative expansion
- Neutral current decay to follow soon.



Results

$$\ell = \mu$$







Results

$$\ell = \tau$$







Results

$$\ell = \mu$$





Results

$$\ell = \tau$$



Results

$$\kappa_{\ell}^h = \frac{Br(N \rightarrow \ell h)}{Br(N \rightarrow \ell X)} = \frac{\Gamma(N \rightarrow \ell h)}{\Gamma(N \rightarrow \ell X)}$$



Conclusion

- Bump Hunt in Belle II hints at $m_N = 354 \text{ MeV}$ (not statistically significant)
- Angular Coefficients show no preference for new physics
- Heavy Sterile Neutrinos can decay semi-hadronically
- We calculated the charged current inclusive decay rate up to $\mathcal{O}(\alpha_s^4)$
- For $\ell = e, \mu$ perturbativity ok for $m_N > 1.5 \text{ GeV}$
- For $\ell = \tau$ perturbativity ok for $m_N > 3 \text{ GeV}$
- Neutral Current comparable in magnitude




Backup: Contour Integration

- Integration over **Keyhole contour**
- Low energy regime does not contribute!

$$\Gamma(N \rightarrow \ell x) \propto 12\pi \int_0^{m_N^2} ds \left(1 - \frac{s}{m_N^2}\right)^2 \left[\left(1 + 2\frac{s}{m_N^2}\right) \Im \Pi^{(1+0)}(s) - 2\frac{s}{m_N^2} \Im \Pi^{(0)}(s) \right]$$

$$= 6\pi i \oint_{s=m_N^2} ds \left(1 - \frac{s}{m_N^2}\right)^2 \left[\left(1 + 2\frac{s}{m_N^2}\right) \Pi^{(1+0)}(s) - 2\frac{s}{m_N^2} \Pi^{(0)}(s) \right]$$



see e.g. Braaten, Narison and Pich; [Nucl.Phys.B 373 \(1992\) 581-612](#)