



Contribution ID: 800

Type: Parallel

General Linblad equation for in-QGP quarkonia evolution

Accurate modeling and understanding of quarkonium production in AA collisions requires a formalism that preserves the quantum properties of a microscopic $Q\bar{Q}$ system while treating the interaction of such pairs with the QGP. The open quantum system approach has recently emerged as one of the most fruitful schemes to meet such requirements. However, the quantum master equations obtained so far in this approach and currently used to make predictions for the upsilon suppression at RHIC and LHC are derived assuming a strict ordering between the QGP temperature and the energy gaps (ΔE) of the upsilon bound states. This limits their predictive power as the QGP cooling interpolates between the quantum Brownian regime ($T \gtrsim \Delta E$) and the quantum optical regime ($T \lesssim \Delta E$).

In our contribution, we derive and present a more general non-abelian quantum master equation of the Linblad type, which does not suffer from these limitations and thus allows to faithfully describe the quantum evolution of the $b\bar{b}$ pairs during the whole QGP evolution. Preliminary results will be shown showing the interpolating features of our equation between the two regimes. We also establish the contact with the QME previously obtained in the QBM and QO regimes and give some concrete perspectives for its efficient solution.

Secondary track

Author: Dr DADDI HAMMOU, Aoumeur (Subatech)

Co-authors: Dr BLAIZOT, Jean-Paul (IPHT); GOUSSET, Thierry (Subatech); GOSSIAUX, pol bernard (subatech)

Presenter: Dr DADDI HAMMOU, Aoumeur (Subatech)

Session Classification: T04

Track Classification: T04 - Ultra-relativistic Nuclear Collisions