

Precision Measurements - Theory

Andreas von Manteuffel
Rene Poncelet
Raoul Röntschi



THE HENRYK NIEWODNICZAŃSKI
INSTITUTE OF NUCLEAR PHYSICS
POLISH ACADEMY OF SCIENCES



Current issues

NNLO + PS

- generality
- accuracy

Mixed EW x QCD corrections

- amplitudes
- subtraction

FO subtraction schemes at NNLO

- generality & efficiency
- numerical integration

Multi-loop amplitudes

- Methods
- Automation

Theory uncertainties


- fixed-order
- resummation
- PDFs (aN3LO/flavour schemes)

Flavoured jets

- flavoured algorithms
- experimental implementation

Toponium/top-pair threshold

- bound state
- resummation



[Credit: SHERPA]


...

Theory uncertainties

Precision example: strong coupling from pT(Z)


[ATLAS 2309.12986]

Sensitivity of Z-boson's recoil to the strong coupling constant:



→ at low pT resummation regime!

→ theory uncertainty?



Theory uncertainties in Higgs production

[LHCH(XS)WG YR4' 16]

N4LO approximation
[Das, Moch, Vogt '20]

aN3LO PDFs
[MSHT'22]
[NNPDF'24]

Exact top-mass dependence
through NNLO QCD
[Czakon, Harlander, Klappert, Niggetiedt'21]

Input parameters

$\sqrt{S}$	13 TeV
m_h	125 GeV
PDF	PDF4LHC15_nnlo_100
$\alpha_s(m_Z)$	0.118
$m_t(m_t)$	162.7 GeV ($\overline{MS}$)
$m_b(m_b)$	4.18 GeV ($\overline{MS}$)
$m_c(3GeV)$	0.986 GeV ($\overline{MS}$)
$\mu = \mu_R = \mu_F$	62.5 GeV ($= m_H/2$)

$\delta(\text{scale})$	$\delta(\text{trunc})$	$\delta(\text{PDF-TH})$	$\delta(\text{EW})$	$\delta(t, b, c)$	$\delta(1/m_t)$
+0.10 pb -1.15 pb	± 0.18 pb	± 0.56 pb	± 0.49 pb	± 0.40 pb	± 0.49 pb
+0.21% -2.37%	$\pm 0.37\%$	$\pm 1.16\%$	$\pm 1\%$	$\pm 0.83\%$	$\pm 1\%$

N3LO HEFT
[Mistlberger'18]

Improved QCD-EW predictions
[Bonetti, Melnikov, Trancredi'18] [Anastasiou et al '19]
[Bonetti et al. '20][Bechetti et al. '21] [Bonetti, Panzer, Trancredi '22]


Bottom-top-interference
[Czakon, Eschment, Niggetiedt,
Poncelet, Schellenberger '23, '24]

Uncertainties in precision phenomenology

Experiments are getting more precise → theory uncertainties matter!

Sources of theory uncertainties:

- parametric (values of coupling parameters etc.)
→ variation of parameters within their uncertainties
- parton distribution functions (PDFs)
→ different error propagation methods (fit parameter, replicas,...)
- non-perturbative parameters in Monte Carlo simulations.
→ needs data constraints by definition. Problematic if dominant effect...
- missing higher orders in fixed-order and resummed predictions (MHOU)
→ tricky because we are trying to estimate the unknown....



[Credit: SHERPA]

Missing higher orders

Notation from: [\[Tackmann 2411.18606\]](#)

Beyond Scale Variations: Perturbative Theory Uncertainties from Nuisance Parameters

Generic perturbative expansion:

$$f(\alpha) = f_0 + f_1\alpha + f_2\alpha^2 + f_3\alpha^3 + \dots$$

f_i : the coefficient of the series, potentially unknown

We can compute the truncated series: $\hat{f}_i$: the true value, i.e. a value we actually computed

$$f^{\text{LO}}(\alpha) = \hat{f}_0 \quad f^{\text{NLO}}(\alpha) = \hat{f}_0 + \hat{f}_1\alpha \quad f^{\text{NNLO}}(\alpha) = \hat{f}_0 + \hat{f}_1\alpha + \hat{f}_2\alpha^2$$

The missing terms are the source of uncertainty.

(assume convergence $\rightarrow$ the first missing is the dominant one)

$$f^{\text{LO}+1}(\alpha) = \hat{f}_0 + f_1\alpha \quad f^{\text{NLO}+1}(\alpha) = \hat{f}_0 + \hat{f}_1\alpha + f_2\alpha^2 \quad f^{\text{NNLO}+1}(\alpha) = \hat{f}_0 + \hat{f}_1\alpha + \hat{f}_2\alpha^2 + f_3\alpha^3$$

Challenge: how to estimate $f_1, f_2, f_3, \dots$ without computing them?



Theory uncertainties from scale variations

Lets focus on QCD as an example: $\alpha = \alpha_s(\mu_0)$


$$d\sigma = d\sigma^{(0)} + \alpha_s d\sigma^{(1)} + \alpha_s^2 d\sigma^{(2)} + \dots \xrightarrow{\text{RGE}} \mu \frac{d\sigma^{(n)}}{d\mu} = \mathcal{O}((n_0+n+1))$$

Of same order as the next dominant term
 $\rightarrow$ exploiting this to estimate size of $d\sigma^{(n+1)}$

Sometimes it does works ... sometimes not



[ATLAS 2301.09351]



[Grazzini, talk]

Short comings of scale variations

- **not always reliable** ... however in most cases issues are understood/expected:
new channels, phase space constraints, etc. → often we can design workarounds
- however, some issues are more fundamental:
 - how to choose the **central scale**? → **not a physical parameter**, no 'true' value
(Principle of fastest apparent convergence, principle of minimal sensitivity,...)
 - how to propagate the estimated uncertainty, **no statistical interpretation!**
 - what about **correlations**? Based on 'fixed form' of the lower orders and RGE.

(At the moment) two alternative approaches under investigation:

"Bayesian"

[Cacciari,Houdeau 1105.5152]

[Bonvini 2006.16293]

[Duhr, Huss, Mazeliauskas, Szafron 2106.04585]

"Theory Nuisance Parameter"

[Tackmann 2411.18606]

→ W mass extraction: [CMS 2412.13872]


[Cal,Lim, Scott,Tackmann Waalewijn 2408.13301]

[Lim, Poncelet, 2412.14910]


Example: uncertainties in jet cross sections

[1903.12563]

DiJets, R-dependence fit to $10^4 \cdot (a + b \log(R) + cR^2)$, $\mu_{R/F} = p_T^{Jet}$




[2503.21866]



Flavoured jets


Flavoured jets

jet that initiated from a "hard scatter" product of specific flavour:
bottom, charm , "quark/gluon"




Flavoured jets

jet that initiated from a "hard scatter" product of specific flavour:
bottom, charm , "quark/gluon"




Flavoured jets

jet that initiated from a "hard scatter" product of specific flavour:
bottom, charm , "quark/gluon"




Flavoured jets

jet that initiated from a "hard scatter" product of specific flavour:
bottom, charm , "quark/gluon"



Infrared safety issues with flavoured jets I




- IRC unsafe due to $g \rightarrow \text{quark-anti-quark}$ splitting
- Quarks massless: cross-section not defined
- Quarks massive: logarithmic sensitivity to quark mass
- Can resolved by proper flavour recombination schemes:

scheme	jet contents			
	b	$b + \bar{b}$	$b + b$	
"any flavour"	b	b	b	simplest experimentally (but collinear unsafe for $m_b \rightarrow 0$)
net flavour	b	g	$2b$	theoretically "ideal" definition; but not robust wrt B-Bbar oscillations
flavour modulo 2	b	g	g	theoretically OK; robust wrt B-Bbar oscillations

Picture from
[Gauld et al. 2302.12844]

[Salam]

Infrared safety issues with flavoured jets II




- Collinear unsafe if pT requirement on the quark is present
- Not implementable in pQCD with massless quarks
→ proper treatment needs fragmentation functions
→ NNLO QCD example:

B-hadron production in NNLO QCD: application to LHC ttbar events with leptonic decays,
Czakon, Generet, Mitov and Poncelet, 2102.08267

Picture from
[Gauld et al. 2302.12844]

Infrared safety issues with flavoured jets III



- Starting at NNLO QCD:
→ Soft singularity from quark pairs
- Massless quarks → cross section not defined
- Massive quarks → logarithmic IRC sensitivity $\ln\left(\frac{m}{p_T}\right)$
- Needs modified jet algorithms!

Picture from
[Gauld et al. 2302.12844]

New proposals for flavour-safe anti-kT jets


- Flavour with Soft-drop **Practical Jet Flavour Through NNLO**
Caletti, Larkoski, Marzani, Reichelt 2205.01109
- Flavour anti-kT **Infrared-safe flavoured anti-kT jets,**
Czakon, Mitov, Poncelet 2205.11879
- Fragmentation approach **A Fragmentation Approach to Jet Flavor**
Caletti, Larkoski, Marzani, Reichelt 2205.01117
B-hadron production in NNLO QCD: application to LHC ttbar events with leptonic decays,
Czakon, Generet, Mitov and Poncelet, 2102.08267
- Flavour dressing → standard anti-kT + flavour assignment
QCD-aware partonic jet clustering for truth-jet flavour labelling
Buckley, Pollard 1507.00508
A dress of flavour to suit any jet
Gauld, Huss, Stagnitto 2208.11138
- Interleaved flavour neutralisation
Flavoured jets with exact anti-kT kinematics and tests of infrared and collinear safety
Caola, Grabarczyk, Hutt, Salam, Scyboz, Thaler 2306.07314
- TBC...

Outcome LH23


Today on the arxiv

Flavoured jet algorithms: a comparative study

Arnd Behring,¹ Simone Caletti,² Francesco Giuliani,³ Radosław Grabarczyk,^{4,5} Andreas Hinzmann,⁶ Alexander Huss,⁷ Joey Huston,⁸ Ezra D. Lesser,⁷ Simone Marzani,⁹ Davide Napoletano,¹⁰ Rene Poncelet,¹¹ Daniel Reichelt,^{7,12} Alberto Rescia,⁹ Gavin P. Salam,^{5,14} Ludovic Scyboz,¹⁵ Federico Sforza,⁹ Andrzej Siódmok,^{7,16} Giovanni Stagnitto,¹⁰ James Whitehead,¹⁶ Ruide Xu¹⁷





Many analyses from theory and experiment groups → improved understanding on all sides




Open Questions

- impact of hadronisation?
- impact on jet substructure?
- how to incorporate in experiment?
- revisit the truth level labels in the training of ML taggers?
- can we get an experimental handle on $g \rightarrow f\bar{f}$?



Amplitudes

Typical multi-loop calculation work-flow



Integration-By-Parts reduction

$$a_i^{(L),p} = \sum_i c_{j,i}(\{p\}, \epsilon) \mathcal{I}(\{p\}, \epsilon) \quad \rightarrow \text{prohibitively large number of integrals}$$

$$\mathcal{I}_i(\{p\}, \epsilon) \equiv \mathcal{I}(\vec{n}_i, \{p\}, \epsilon) = \int \frac{d^d k_1}{(2\pi)^d} \frac{d^d k_2}{(2\pi)^d} \prod_{k=1}^{11} D_k^{-n_{i,k}}(\{p\}, \{k\})$$

Integration-By-Parts identities connect different integrals $\rightarrow$ system of equations
 $\rightarrow$ only a small number of independent “master” integrals

$$0 = \int \frac{d^d k_1}{(2\pi)^d} \frac{d^d k_2}{(2\pi)^d} l_\mu \frac{\partial}{\partial l_\mu} \prod_{k=1}^{11} D_k^{-n_{i,k}}(\{p\}, \{k\}) \quad \text{with} \quad l \in \{p\} \cap \{k\}$$



$$a_i^{(L),p} = \sum_i d_{j,i}(\{p\}, \epsilon) \text{MI}(\{p\}, \epsilon)$$

Master integrals

Differential Equations: $d\vec{MI} = dA(\{p\}, \epsilon)\vec{MI}$

[Remiddi, 97]

[Gehrmann, Remiddi, 99]

[Henn, 13]

Canonical basis: $d\vec{MI} = \epsilon d\tilde{A}(\{p\})\vec{MI}$

Simple iterative solution




$$MI_i = \sum_w \epsilon^w \tilde{MI}_i^w \quad \text{with} \quad \tilde{MI}_i^w = \sum_j c_{i,j} m_j$$

Chen-iterated integrals
"Pentagon"-functions


[Chicherin, Sotnikov, 20]

[Chicherin, Sotnikov, Zoia, 21]


Reconstruction of Amplitudes



$$A_{34;q}^{(2),N_c^2}, A_{\delta;q}^{(2),N_c}$$



$$A_{34;q}^{(2),1}, A_{\delta;q}^{(2),N_c}, A_{\delta;q}^{(2),1/N_c}$$



$$A_{34;l}^{(2),N_c}, A_{34;l}^{(2),1/N_c}, A_{\delta;l}^{(2),1/N_c^2}$$

[Badger et al '23]

Many possible optimizations



- Syzygy's to simplify IBPs
- Exploitation of Q-linear relations
- Denominator Ansatz
- On-the-fly partial fractioning
- ...

amplitude	helicity	original	stage 1	stage 2	stage 3	stage 4
$A_{34;q}^{(2),1}$	- + + - +	94/91	74/71	74/0	22/18	22/0
$A_{34;q}^{(2),1}$	- + - + +	93/89	90/86	90/0	24/14	18/0
$A_{34;q}^{(2),1/N_c^2}$	- + + - +	90/88	73/71	73/0	23/18	22/0
$A_{34;q}^{(2),1/N_c^2}$	- + - + +	90/86	86/82	86/0	24/14	19/0
$A_{34;l}^{(2),1/N_c}$	- + - + +	89/82	74/67	73/0	27/14	20/0
$A_{34;l}^{(2),1/N_c}$	- + + - +	85/81	61/58	60/0	27/18	20/0
$A_{34;q}^{(2),N_c^2}$	- + - + +	58/55	54/51	53/0	20/18	20/0

Reduction of complexity

Numerical evaluation of two-loop matrix elements

[De Laurentis, Ita, Sotnikov'23][De Laurentis, Ita, Page, Sotnikov'25]



Efficient function basis representation power the speed.

If too expensive $\rightarrow$ numerical interpolation grids (used a lot for 2D problems)?

Methods for high dimensions?


Topics for amplitude discussions

- Geometry, elliptic functions, Calabi-Yau, ...
- Dimensional regularization and γ_5
- Automation @ two-loops
- Standardisation of interfaces
- Efficiency:
 - ML
 - **Interpolation**
 - partial fractioning
 - singularities

Toponium / top-pair threshold


Surprises at the threshold

[ATLAS'23] Entanglement




Threshold QCD enhancements

[Nason,Re,Rottoli'25]



Sounds like super topic for a discussion :)

[CMS'25] Toponium



Numerical integration and non-positive integrands or negatively weighted events

The problem

- Numerical integration of highly dimensional integrands $\rightarrow$ Monte Carlo Sampling

Integral

$$I = \int_{\mathbf{x} \in \Omega} d\mathbf{x} f(\mathbf{x})$$

MC estimate

$$\hat{I} = \frac{1}{N} \sum_{i=1}^N f(\mathbf{x}_i),$$

MC error estimate

$$\delta \hat{I} = \sqrt{\frac{1}{N-1} \left(\frac{1}{N} \sum_{i=1}^N f^2(\mathbf{x}_i) - \hat{I}^2 \right)}$$

- Variance reduction techniques improve performance, mapping $\mathbf{H} : \Omega \rightarrow \Omega, \mathbf{x} \mapsto \mathbf{H}(\mathbf{x})$

$$I = \int_{\mathbf{H}(\mathbf{x}) \in \Omega} d\mathbf{H} \frac{f(\mathbf{x})}{h(\mathbf{x})}$$


$$h(\mathbf{x}) = \left| \det \left(\frac{\partial \mathbf{H}(\mathbf{x})}{\partial \mathbf{x}} \right) \right|$$

Find with $h(\mathbf{x})$ adaptive MC techniques: VEGAS [Lepage'78], Parni [Hameren'14],
ML techniques: Normalising Flows Iflow [Bothmann'20] Madnis [Heimel'22], ...

Non-positive definite integrands

- Non-definite integrands introduce computational challenges

→ cancellation between +/- parts increase the variance

- Consider extreme case: $|f(x)|/h(x) = w = \text{const.}$



MC estimate: $\hat{I} = w \frac{N_+ - N_-}{N} \equiv w(2\alpha - 1) \quad \alpha = N_+/N$

- Lower bound on variance:

$$\text{Var}(\hat{I}) = w^2 - w^2(2\alpha - 1)^2 = w^2(4\alpha(1 - \alpha))$$

→ relative uncertainty: $\frac{\delta \hat{I}}{\hat{I}} = \frac{1}{\sqrt{N-1}} \frac{\sqrt{\alpha(1-\alpha)}}{\alpha - \frac{1}{2}}$

Rephrased: at some point it doesn't matter any more how good your adaptive MC is...



Negative Weights

How to improve convergence
of non-positive definite MC integrals?

→ phase space splitting

[Janssen, Schumann, Poncelet'25]


How to avoid negative weighted events?

→ Design a positive matching
for example ESME, KrKNLO, ...

[van Beekveld, Ravasio, Helliwell, Karlberg, Salam,
Scyboz, Soto-Ontoso, Soyez, Zanolì'25]

[Whitehead, Sarmah, Siódmok'24]

[Andersen, Maier, Maitre, 2023]



I have negative weights
→ How to get rid of them?

Cell-resampling [Anderson, Maier'21]

Current issues

NNLO + PS

- generality
- accuracy

Mixed EW x QCD corrections

- amplitudes
- subtraction

FO subtraction schemes at NNLO

- generality & efficiency
- numerical integration

Multi-loop amplitudes

- Methods
- Automation

Theory uncertainties


- fixed-order
- resummation
- PDFs (aN3LO/flavour schemes)

Flavoured jets

- flavoured algorithms
- experimental implementation

Toponium/top-pair threshold

- bound state
- resummation



[Credit: SHERPA]

...

Summary

Looking forward to a productive week with lots of interesting discussions!

