

# Flavour physics and tests of fundamental interactions

GT2 – Symposium ESPPU

Yasmine Amhis, Giulio Dujany, Christopher Smith

20 January 2025


# The intensity frontier

Probe New Physics pushing the experiment's luminosity rather than the energy scale

Strategy:

- Look for deviations from theory predictions in precise measurements of SM processes
- Search for hugely suppressed or forbidden processes in SM


Require large datasets and precise measurements



**“Symmetry” (08/01/2008)**

# Why the flavour sector

- Majority of SM parameters
- Unexplained patterns and hierarchies in masses and CKM elements
- Accidental symmetries (baryon and lepton numbers)
- Indirect measurements are powerful prediction and discovery tool
  - Predicted half the quarks before their direct discovery!
  - Strong constraints on masses of c, t, H before their discovery



## Link to open questions

- Matter/antimatter imbalance
- Strong CP problem
- Dark matter

# Inputs to GT2

- 11 direct contributions on [esppu.in2p3.fr](https://esppu.in2p3.fr) about
  - Current and proposed future experiments (Belle II, LHCb, n2EDM, GrAHal, PIONEER)
  - Precision measurements at FCC-ee
  - Theory (Phenomenology and lattice QCD)
  - Challenges in an heterogeneous computing environment with exabytes of data
  - New-physics searches with radioactive ions beams
- Dedicated discussion at GDR-Inf annual workshop ([indico.in2p3.fr/event/33443/](https://indico.in2p3.fr/event/33443/))

Amhis	Yasmine	UCLab	072 074	Nouvelle expérience <b>Sauvages</b> <b>QCD</b>	↳ Intérêt de la communauté française pour une participation au programme de physique et aux améliorations du détecteur LHCb pour la période 2035-2041 (LHCb Upgrade II)
Blossier	Benoît	UCLab Orsay	072 074	Théorie/Phénoménologie <b>Calcul et données</b> <b>BSM / Nouvelle physique</b> <b>Sauvages</b> <b>QCD</b>	↳ QCD sur réseau, texte en anglais
DELAHAYE	Pierre	GANIL, CAEN, France	072 071	Physique sur accélérateurs <b>Physique hors-accelérateurs</b> <b>R&amp;D détecteurs</b> <b>Nouvelle expérience</b> <b>Calcul / Nouvelle physique</b> <b>Physique électrofaible</b> <b>Sauvages</b>	↳ "PIONEER: A Next-Generation Pion Decay Experiment at PSI"
Delmastro	Marco	LAPP Annecy	071 072	Physique sur accélérateurs <b>Higgs</b> <b>Physique électrofaible</b> <b>Sauvages</b>	↳ Expression d'intérêt pour les mesures de précision au FCC-ee
Farget	Fanny	GANIL	072	Physique hors-accelérateurs <b>BSM / Nouvelle physique</b>	↳ Radioactive Ion Beams opportunities to reach Physics Beyond the Standard Model
Farget	Fanny	GANIL	071 072	Physique hors-accelérateurs <b>BSM / Nouvelle physique</b>	↳ Exploring physics beyond the Standard Model with the new DESIR facility at GANIL
Gligorov	Vladimir	LPNHE	071 072	Physique sur accélérateurs <b>Physique hors-accelérateurs</b> <b>R&amp;D détecteurs</b> <b>Calcul et données</b> <b>Développement durable</b>	↳ Reflection sur les défis technologiques dans calcul et données
Lefort	Thomas	LPC Caen	072 076	Physique hors-accelérateurs <b>BSM / Nouvelle physique</b>	↳ neutron EDM
NORTIER	Florian	IP2I Lyon	071 072	Physique sur accélérateurs <b>Théorie/Phénoménologie</b> <b>Higgs</b> <b>BSM / Nouvelle physique</b> <b>Physique électrofaible</b> <b>Sauvages</b>	↳ Contribution of the IP2I Theory Group to the ESPPU 2025
Pugnat	Pierre	LNCMI, CNRS, Grenoble	072	Physique hors-accelérateurs <b>Nouvelle expérience</b> <b>Nouveaux instruments expérimentaux</b> <b>BSM / Nouvelle physique</b>	↳ Grenoble Axion Haloscopes: From BabyGrAHal to GrAHal for Axion Dark Matter Search
Serrano	Justine	CPPM	072	Physique sur accélérateurs <b>R&amp;D détecteurs</b> <b>BSM / Nouvelle physique</b> <b>Sauvages</b>	↳ The Belle II experiment and its upgrade

6-8 nov. 2024  
Cabourg


# Two-pillars strategy

- Big experiments with a large physics program
- Smaller experiments dedicated to
  - Measurement of crucial observables (EDM,  $g-2$ , LFV experiments)
  - Dedicated searches for long-lived particles (Axion, ALPs, ...)




# Big experiments with a large physics program

# Big experiments with a large physics program










- Hermetic detector
  - Allow inclusive and missing energy analyses
- Asymmetric  $e^+e^-$  collider at  $\Upsilon(4S)$ 
  - Clean environment
  - Entangled B pair
  - ~100% trigger efficiency for BB events
  - Similar BB, cc and  $\tau\tau$  production x-section
  - Good efficiency for photons and neutral particles



- General purpose detector in the forward region
- pp collisions
  - Large bb and cc production cross section
  - All B-hadron produced
  - Large boost
- Complementary program of QCD and EW measurements
- Fully software trigger since Run 3

# Main physics interests

1. CP violation and unitary triangle
2. Rare B decays sensitive to New Physics
3. Tests of lepton flavour violation and lepton flavour universality violation
4. Charm and tau physics
5. Spectroscopy
6. ...





# Upgrade opportunities

- Without upgrades expect  $50 \text{ fb}^{-1}$  for LHCb (2033) and  $5\text{-}10 \text{ ab}^{-1}$  for Belle II (2032)
- Most of key observables still statistically limited
- Calls for larger datasets
  - Increase instantaneous luminosity
  - Upgrade detectors and acquisition systems to cope with increased luminosity




- Goal:  $50 \text{ ab}^{-1}$
- When:  $\sim 2032$
- More robust against beam background
- Improved performances
- Possibly polarised  $e^-$  beam for precise measurements of electroweak and  $\tau$  parameters



- Goal:  $300 \text{ fb}^{-1}$
- When: 2034 (Long Shutdown 4), some already in 2027 (Long Shutdown 3)
- Precise timing and high granularity essential to reduce pileup
- Heterogeneous trigger
- Probe a 90% higher new physics mass scale for fixed couplings

# Computing challenges

## A. Cerri, University of Sussex



- Flavour physics at the forefront of the Exascale computing
- Need for highly efficient and energy-saving heterogeneous high performance computing
- Need for interdisciplinary teams of physicists and computer engineers

# Longer term wish: Tera-Z + Giga-W on a $e^+e^-$ machine with a suitable detector

- A Tera-Z factory has several advantages of LHCb and Belle II combined
- $W^+W^-$  run allows direct access to CKM elements



Attribute	$\Upsilon(4S)$	$pp$	$Z^0$
All hadron species		✓	✓
High boost		✓	✓
Enormous production cross-section		✓	
Negligible trigger losses	✓		✓
Low backgrounds	✓		✓
Initial energy constraint	✓		(✓)

[Eur.Phys.J.Plus 136 \(2021\) 8, 837](#)

- Preferred option is a circular  $e^+e^-$  collider
  - A smaller number of Z from a linear collider will clearly provide less opportunities for World-leading measurements (considering the available samples after Belle II and LHCb)
  - Multiple collision points allow for an experiment optimized for the needs of flavour physics

# Opportunities at Tera-Z + Giga-W

- Improved precision on CKM parameters (angles and sides Unitarity triangle,  $\Phi_s$ ; direct access to  $V_{cb}$  from W decays)
- Unique sensitivity to very rare FCNC B decays ( $B_{(s)} \rightarrow \mu\mu$ ,  $B_s \rightarrow \pi\pi$ ,  $b \rightarrow s(d)\ell\ell$ ,  $b \rightarrow s(d)\pi\pi$ ,  $b \rightarrow s(d)uu$ ) and challenging  $B_c$  decays ( $B_c \rightarrow \mu u$ ,  $B_c \rightarrow \tau u$ ) sensitive to possible new physics
- Improve measurement of  $\tau$  properties and reach for LFV decays
- Opportunities in charm physics, heavy flavour spectroscopy



# A suitable detector for flavour physics


- Good momentum resolution down to 100 MeV
- Good vertex resolution to separate secondary vertices and perform time-dependent measurements
- pion/kaon/proton separation across full kinematical range
- Electromagnetic calorimeter with good resolution and granularity (transverse and longitudinal) down to low energies
- Good efficiency and precision in reconstructing  $K_S$  and  $\Lambda$

# Smaller dedicated experiments

# Smaller dedicated experiments

CP violation: Very constrained in the SM : - Only one phase in the CKM matrix,  
 - Strong CP violation is (puzzlingly) tiny.


Tiny EDMs (leptons, nucleons, atomic), nuclear CP asymmetries, ...  
 Does not suffice to induce the baryon asymmetry of the universe,



Most BSM bring many new CP-violating phases and break this pattern.  
 Scales up to  $O(100 \text{ TeV})$  can be probed.

n2EDM@PSI for the neutron EDM,  
 Ultra cold neutrons, comagnetometer.

DESIR@GANIL for asymmetries,  
 Beta decay spectra, radioactive ion beams.




# Smaller dedicated experiments

**Lepton flavor violation:** In the SM, charged leptonic flavors appear essentially conserved.

Many tests using B, D, K,  $\tau$  decays + dedicated experiments for  $\mu$ .

COMET experiment (COherent Muon to Electron Transition) at JPARC



Sensitivity up to 1000 TeV.



Many other initiatives (conversion,  $\mu \rightarrow e\gamma$ ,  $\mu \rightarrow eee$ )

**Lepton flavor universality:** In the SM, gauge interactions do not care about flavors.

Again, many tests both heavy meson decay, or directly via EW-scale colliders.

Dedicated experiment for the pion:  $\frac{\pi \rightarrow \mu\nu}{\pi \rightarrow e\nu}$  @ **PIEER**



# Smaller dedicated experiments


**V<sub>ud</sub>:** It is the only 'naturally occurring' CKM element, best measured at low energy

Precise test of the unitarity of CKM:  $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$ .

Nuclear (superallowed):



Pion decay  $\pi \rightarrow \ell \nu$ :



**Baryon and lepton numbers:** Accidentally but exactly conserved in the SM Lagrangian.

	$\Delta B$	$\Delta L$
Proton decay	-1	$\pm 1, \pm 3$
$0\nu\beta\beta$	0	$\pm 2$
$n - \bar{n}$ oscillations	$\pm 2$	0
Dinucleon trans.	-2	$x$

These are flavored transitions, but mostly addressed by the GT3-neutrinos.


# Smaller dedicated experiments



New light particle: Are we sure nothing else exists at low energy?  
Dark matter is presumably made of particles!

A whole dark sector may exist with dark fermions, photons, (pseudo)scalars,...

Generalist searches (e.g. beam dump, nuclear recoil, colliders, ...)

Precision observables:  $\frac{1}{(\text{many TeV})^n} \rightarrow \frac{(\text{very small})}{(\leq \text{GeV})^n}$

Many searches for axions as solutions to the strong CP puzzle (including ).



Dielectric disks for interference + resonance

GrAHal

RF resonant cavity in a large 42T hybrid magnet @LNCMI

Most flavor observables involve light hadrons:

Observables  $\sim |(\text{SD physics, calculable}) \times (\text{hadronic matrix element})|^2$

↑  
Lattice calculations are crucial to match the experimental precision

Challenges to attain the percent level accuracy:

Needs to control QED and  $m_u \neq m_d$  effects,  
(directly on fine lattice and/or indirectly via low-energy EFT)

Road to inclusiveness requires dealing with many hadron states,

Developments for nuclear observables (from  $V_{ud}$  to  $0\nu\beta\beta$  or proton decay).

Past few decades have seen great achievements,

...but the next generation experiments ask to push even further!

## Short-medium term

- Fully exploit LHCb and Belle II and support their proposed upgrades
- Support smaller dedicated experiments (n2EDM, COMET, GrAhaI, MADMAX, Gbar, ...)
- Pursue synergy with NuPECC for precise measurements at the boundary between particle and nuclear physics (e.g. with DESIR at GANIL)

## Long-term

- Tera-Z + Giga-W on a  $e^+e^-$  machine with a well designed detector
- Important to continue investing in smaller dedicated experiments with potential high-impact discoveries (including probes for a possible dark sector)



## Specific needs from Theory, Detector and Computing

# BACKUP

# Discussion items

-

# LHCb Upgrade I



- 5 time larger instantaneous luminosity
- Greatly improved tracker & PID granularity, 30 MHz detector readout & GPU tracking trigger
- More than doubled the efficiency for hadronic signals

# LHCb-U2 physics opportunities

Observable	Old LHCb (up to $9 \text{ fb}^{-1}$ )	LHCb Upgrade 2 Scoping Document		
		Upgrade I ( $23 \text{ fb}^{-1}$ )	Upgrade I ( $50 \text{ fb}^{-1}$ )	Upgrade II ( $300 \text{ fb}^{-1}$ )
<b>CKM tests</b>				
$\gamma (B \rightarrow DK, \text{ etc.})$	$2.8^\circ$ [18, 19]	$1.3^\circ$	$0.8^\circ$	$0.3^\circ$
$\phi_s (B_s^0 \rightarrow J/\psi\phi)$	20 mrad [22]	12 mrad	8 mrad	3 mrad
$ V_{ub} / V_{cb}  (A_b^0 \rightarrow p\mu^-\bar{\nu}_\mu, \text{ etc.})$	6% [55, 56]	3%	2%	1%
<b>Charm</b>				
$\Delta A_{CP} (D^0 \rightarrow K^+K^-, \pi^+\pi^-)$	$29 \times 10^{-5}$ [25]	$13 \times 10^{-5}$	$8 \times 10^{-5}$	$3.3 \times 10^{-5}$
$A_\Gamma (D^0 \rightarrow K^+K^-, \pi^+\pi^-)$	$11 \times 10^{-5}$ [29]	$5 \times 10^{-5}$	$3.2 \times 10^{-5}$	$1.2 \times 10^{-5}$
$\Delta x (D^0 \rightarrow K_S^0\pi^+\pi^-)$	$18 \times 10^{-5}$ [57]	$6.3 \times 10^{-5}$	$4.1 \times 10^{-5}$	$1.6 \times 10^{-5}$
<b>Rare decays</b>				
$\mathcal{B}(B^0 \rightarrow \mu^+\mu^-)/\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-)$	69% [30, 31]	41%	27%	11%
$S_{\mu\mu} (B_s^0 \rightarrow \mu^+\mu^-)$	—	—	—	0.2
$A_\Gamma^{(2)} (B^0 \rightarrow K^{*0}e^+e^-)$	0.10 [58]	0.060	0.043	0.016
$S_{\phi\gamma} (B_s^0 \rightarrow \phi\gamma)$	0.32 [59]	0.093	0.062	0.025
$\alpha_\gamma (A_b^0 \rightarrow \Lambda\gamma)$	$^{+0.17}_{-0.29}$ [60]	0.148	0.097	0.038



# Comparison expected yields


Working point	Lumi. / IP [ $10^{34} \text{ cm}^{-2} \cdot \text{s}^{-1}$ ]	Total lumi. (2 IPs)	Run time	Physics goal
Z first phase	100	26 $\text{ab}^{-1}$ /year	2	
Z second phase	200	52 $\text{ab}^{-1}$ /year	2	150 $\text{ab}^{-1}$


Particle production ( $10^9$ )	$B^0$	$B^-$	$B_s^0$	$\Lambda_b$	$c\bar{c}$	$\tau^- \tau^+$
Belle II	27.5	27.5	n/a	n/a	65	45
FCC- <i>ee</i>	400	400	100	100	800	220

[Eur. Phys. J. C 79 \(2019\) 474](#)


# Current and planned b-factories



All b hadrons  
Large production rate



Clean environment  
Entanglement  
Neutral decay products  
Hermiticity (neutrinos)  
 $\tau$  factory



Many of the  
advantages of  
LHCb and Belle II