ID de Contribution: 6

Type: Non spécifié

## The ParB-CTP Cycle Activates Phase Separation in Bacterial DNA Segregation

Cell function relies on liquid-like organelles formed through phase transitions, yet the mechanisms ensuring their specificity and rapid assembly remain poorly understood. In bacterial chromosome segregation via the ParABS system, hundreds of ParB proteins are recruited around the centromere-like *parS* sequence forming the partition complex. Recent studies have shown that ParB binds CTP and undergoes cycles of binding and unbinding near *parS*, however, this accounts for the recruitment of only a small fraction of ParB molecules, leaving its role unclear. Separately, a lattice gas model with fixed interaction energy has been proposed to describe ParB phase separation, but it fails to explain key experimental observations, including the absence of droplets in ParB variants. We reconcile these two perspectives by proposing that the ParB-CTP cycle acts as a molecular switch that enhances ParB-ParB interactions, triggering phase transition from vapor to liquid-like condensates.

Our hypothesis is supported by numerical simulations of droplet formation and experiments showing that ParB variants disrupting the CTP cycle fail to undergo phase separation. These findings establish a mechanistic framework for ParB-CTP-mediated phase transitions and may have broader implications for understanding the spatial control of intracellular condensate formation.

Auteur principal: Mlle DELIMI, Linda (Laboratoire Charles Coulomb)

**Co-auteurs:** REVOIL, Perrine (Université Toulpuse III Paul Sabatier - Laboratoire de Microbiologie et Génétique Moléculaires); M. RECH, Jérome (Université Toulouse III Paul Sabatier - Laboratoire de Microbiologie et Génétique Moléculaires); BOUET, Jean-Yves (CNRS); Dr WALTER, Jean-Charles (L2C, CNRS & Montpellier Université)

**Orateur:** Mlle DELIMI, Linda (Laboratoire Charles Coulomb)