

Letter of Intent

Virtual Access for Partial Wave Analysis (VA-PWA)

Kai Habermann, Mikhail Mikhasenko, Sebastian Neubert

1 Research Objectives

Physical background

The study of exclusive hadronic processes is of great importance in flavor physics, particularly for elucidating CP violation and for quantifying hadronic contributions crucial for precision measurements. The exploration of hadronic multi-body processes often relies on disentangling contributions of subchannel resonances, detailed by hadron spectroscopy. The field itself remains vibrant, rich with complex phenomena and a significant rate of new discoveries.

A primary analytical tool in this domain is the partial-wave expansion of decay matrix elements. Partial-Wave Analysis (PWA) decomposes complex decays into a set of cascades of two-body transitions, exploiting the fact that the spin and parity quantum numbers of intermediate states are unambiguously and model-independently determined by the angular distributions of the decay products. This technique stands as a cornerstone of modern hadron physics, as it offers the only reliable method for disentangling overlapping hadronic resonances. Its application, however, remains a highly non-trivial undertaking.

Recent collaborative efforts by research groups in Bonn and Bochum ¹ have achieved a significant milestone by establishing a consistent and comprehensive framework within the helicity formalism for describing hadronic systems of arbitrary complexity. This technical advancement extends beyond previously tractable analytic cases, such as two-body and three-body systems, by proposing, implementing, and validating a numerical algorithm applicable to the most general exclusive decays. Furthermore, this development has unified various widely-used branches of the helicity formalism, including plain and canonical representations, and reconciled different phase conventions.

The platform project

We propose to lower entry barriers for amplitude analysis by developing a user-friendly, web-based platform. This platform will allow users to construct and manipulate amplitudes for arbitrary decay processes and generate executable code for amplitude evaluation for various established frameworks. The interface will guide users through the construction process, with real-time setup validation. Decay chains are defined by specifying initial and final states and inserting appropriate resonances. Configurations can be exported in human- and machine-readable formats for reproducibility and downstream integration. Integration with the Particle Data Group (PDG) API provides access to particle and resonance data, including their quantum numbers.

The backend architecture is designed for on-demand code generation, initially for Python. We also plan to support the existing Julia ecosystem and C++ routines. Comprehensive documentation will detail procedures for integrating further custom analysis frameworks, using a standardized intermediate format for broad applicability.

A functional prototype of the platform, shown in Fig. 1, with essential backend capabilities, is accessible online. The prototype allows users to generate valid and executable amplitude code in Python. However, it currently operates on limited infrastructure (a free-tier of the hosting platform *Render*), and substantial development is needed for production readiness.

While a single, universal framework is unlikely to emerge in the community, easy access to a tested and well-documented portal will aid in benchmarking new tools. The platform will

¹<https://doi.org/10.1103/PhysRevD.111.056015>

