

Constraining the dark sector and the initial conditions of the Univers with the EFTofLSS

Théo SIMON

ADE - 06/11/2025





theo.simon@umontpellier.fr

Introduction: the epistemic context




Part I: *How to obtain the most robust and optimised constraints on Λ CDM cosmological parameters from galaxy clustering data?*

Part II: How can clustering data be used to constrain the nature of dark energy and the initial conditions?




Introduction: large-scale structure data

The Sloan Digital Sky Survey (SDSS)



www.sdss.org



BOSS DR12 LRG (Luminous Red Galaxies)

Galaxies (~ 1.5 million) selected in two redshift ranges:

→ LOWZ (SGC/NGC): $0.2 < z < 0.43$ ($z_{\text{eff}} = 0.32$)

→ CMASS (SGC/NGC): $0.43 < z < 0.7$ ($z_{\text{eff}} = 0.57$)

BOSS Collaboration [arXiv:1607.03155]

eBOSS DR16 QSO


Quasars ($\sim 300\,000$) selected in one redshift range:

$0.8 < z < 2.2$ ($z_{\text{eff}} = 1.5$)

eBOSS Collaboration [arXiv:2007.08991]

Introduction: large-scale structure data

DESI



DESI Collaboration [arxiv:2404.03002]

DESI Collaboration [arXiv:2503.14738]

Tracer	Redshift range	N_{tracer}	V_{eff} (Gpc ³)
BGS	0.1 – 0.4	1,188,526	3.8
LRG1	0.4 – 0.6	1,052,151	4.9
LRG2	0.6 – 0.8	1,613,562	7.6
LRG3+ELG1	0.8 – 1.1	4,540,343	14.8
ELG2	1.1 – 1.6	3,797,271	8.3
QSO	0.8 – 2.1	1,461,588	2.7
Lya	1.8 – 4.2	1,289,874	—
LRG3	0.8 – 1.1	1,802,770	9.8
ELG1	0.8 – 1.1	2,737,573	5.8

I- Introduction to the Effective Field Theory of Large-Scale Structures (EFTofLSS)

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Before EFTofLSS...

There are **two main** historical ways of using LSS data:


1. From the linear perturbation theory:

$$P_g(z, k, \mu) \simeq [b_1(z) + f\mu^2]^2 P_m(z, k) \quad \text{Kaiser '87}$$


b_1 : bias parameter, f : growth rate and $\mu = \hat{z} \cdot \hat{k}$

2. BAO angles + Redshift Space Distortion (RSD) information: $\text{BAO}/f\sigma_8$

LSS collaborations conventionally used the second method



Planck Collaboration [arXiv:1807.06209]



The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Before EFTofLSS...

There are **two main** historical ways of using LSS data:

1. From the linear perturbation theory:

$$P_g(z, k, \mu) \simeq [b_1(z) + f\mu^2]^2 P_m(z, k)$$


Lack of precision

Kaiser '87


b_1 : bias parameter, f : growth rate and $\mu = \hat{z} \cdot \hat{k}$

2. BAO angles + Redshift Space Distortion (RSD) information: $\text{BAO}/f\sigma_8$

LSS collaborations conventionally used the second method



Planck Collaboration [arXiv:1807.06209]



The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Before EFTofLSS...

There are **two main** historical ways of using LSS data:

1. From the linear perturbation theory:

$$P_g(z, k, \mu) \simeq [b_1(z) + f\mu^2]^2 P_m(z, k)$$

Lack of precision


Kaiser '87

b_1 : bias parameter, f : growth rate and $\mu = \hat{z} \cdot \hat{k}$


2. BAO angles + Redshift Space Distortion (RSD) information: $\text{BAO}/f\sigma_8$

LSS collaborations conventionally used the second method

Lack of information




Planck Collaboration [arXiv:1807.06209]




The Effective Field Theory of Large-Scale Structures (EFTofLSS)


Before EFTofLSS...



DESI Collaboration [arXiv:2411.12021]



DESI Collaboration [arXiv:2503.14738]



DESI Collaboration [arXiv:2411.12021]

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Main ingredients



EFTofLSS

Carrasco++ [arXiv:1206.2926]


Baumann++ [arXiv:1004.2488]

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Main ingredients

Solve cosmological equations only for **large-scale physics**

$$\begin{aligned}\delta(\mathbf{k}) &= \delta_l(\mathbf{k}) + \delta_s(\mathbf{k}), \\ \delta_l(\mathbf{k}) &= \delta(\|\mathbf{k}\| < \Lambda^{-1})\end{aligned}$$



Carrasco++ [arXiv:1206.2926]

Baumann++ [arXiv:1004.2488]

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Main ingredients

Solve cosmological equations only for **large-scale physics**

$$\begin{aligned}\delta(\mathbf{k}) &= \delta_l(\mathbf{k}) + \delta_s(\mathbf{k}), \\ \delta_l(\mathbf{k}) &= \delta(\|\mathbf{k}\| < \Lambda^{-1})\end{aligned}$$

Dark matter: the Vlasov system

$$\begin{aligned}\dot{\delta}_l + \theta_l &= -\delta\theta_l - v^j \partial_j \delta_l, \\ \dot{\theta}_l + aH\theta_l + \nabla^2 \psi_l &= -v^j \partial_j \theta_l - \partial_i v_l^j \cdot \partial_j v_l^i - \partial_j \left(\frac{1}{\rho_l} \partial_i [\tau^{ij}]_\Lambda \right)\end{aligned}$$

EFTofLSS

Carrasco++ [arXiv:1206.2926]

Baumann++ [arXiv:1004.2488]

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Main ingredients

Solve cosmological equations only for **large-scale physics**

$$\delta(\mathbf{k}) = \delta_l(\mathbf{k}) + \delta_s(\mathbf{k}),$$
$$\delta_l(\mathbf{k}) = \delta(\|\mathbf{k}\| < \Lambda^{-1})$$

Dark matter: the Vlasov system

$$\dot{\delta}_l + \theta_l = -\delta\theta_l - v^j \partial_j \delta_l,$$
$$\dot{\theta}_l + aH\theta_l + \nabla^2 \psi_l = -v^j \partial_j \theta_l - \partial_i v_l^j \cdot \partial_j v_l^i - \partial_j \left(\frac{1}{\rho_l} \partial_i [\tau^{ij}]_{\Lambda} \right)$$

EFTofLSS

$$\nabla^2 \psi_l = \frac{3}{2} \Omega_m(a) (aH)^2 \delta_l$$

Gravity: the Poisson equation

Carrasco++ [arXiv:1206.2926]

Baumann++ [arXiv:1004.2488]

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Main ingredients

Solve cosmological equations only for **large-scale physics**

$$\delta(\mathbf{k}) = \delta_l(\mathbf{k}) + \delta_s(\mathbf{k}),$$
$$\delta_l(\mathbf{k}) = \delta(\|\mathbf{k}\| < \Lambda^{-1})$$

Dark matter: the Vlasov system

$$\dot{\delta}_l + \theta_l = -\delta\theta_l - v^j \partial_j \delta_l,$$
$$\dot{\theta}_l + aH\theta_l + \nabla^2 \psi_l = -v^j \partial_j \theta_l - \partial_i v_l^j \cdot \partial_j v_l^i - \partial_j \left(\frac{1}{\rho_l} \partial_i [\tau^{ij}]_\Lambda \right)$$

EFTofLSS

$$\nabla^2 \psi_l = \frac{3}{2} \Omega_m(a) (aH)^2 \delta_l$$

Gravity: the Poisson equation

$$\vec{x} \rightarrow \vec{x} + \vec{a}$$
$$\vec{v} \rightarrow \vec{v} + \partial_t \vec{a}$$

Symmetries: Galilean invariance

Carrasco++ [arXiv:1206.2926]

Baumann++ [arXiv:1004.2488]

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Main steps

Step by Step...

1- **Solve** dark matter equations **perturbatively**: $\delta_l(\vec{x}, t) = \delta_l^{(1)}(\vec{x}, t) + \delta_l^{(2)}(\vec{x}, t) + \dots + \delta_l^{(n)}(\vec{x}, t)$ *Bernardeau++ '01*

2- Obtain the **mildly non-linear matter power spectrum**:

Carrasco++ [arXiv:1206.2926]

$$P_m(k, \tau) = P_{11}(k, \tau) + P_{22}^\Lambda(k, \tau) + 2P_{13}^\Lambda(k, \tau) + 2P_{c_{\text{comb}}}^\Lambda(k, \tau)$$

Senatore [arXiv:1406.7843]

Mirbabayi++ [arXiv:1412.5169]

3- Write down **all possible operators in the galaxy bias expansion**: $\delta_g = b_1 \delta_l^{(1)} + b_2 \delta_l^{(2)} + R_*^2 \partial^2 \delta_l^{(1)} + \dots$

4- Take into account the **redshift-space distortion** (RSD) effect (to subtract the contribution of the peculiar velocity of the galaxies to the cosmological redshift) *Senatore++ [arXiv:1409.1225]*

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Main steps

Step by Step...

1- **Solve** dark matter equations **perturbatively**: $\delta_l(\vec{x}, t) = \delta_l^{(1)}(\vec{x}, t) + \delta_l^{(2)}(\vec{x}, t) + \dots + \delta_l^{(n)}(\vec{x}, t)$ *Bernardeau++ '01*

2- Obtain the **mildly non-linear matter power spectrum**:

Carrasco++ [arXiv:1206.2926]

$$P_m(k, \tau) = P_{11}(k, \tau) + P_{22}^\Lambda(k, \tau) + 2P_{13}^\Lambda(k, \tau) + 2P_{c2}^\Lambda(k, \tau)$$

Tree-level

One-loop level

Counterterm

Senatore [arXiv:1406.7843]

Mirbabayi++ [arXiv:1412.5169]

3- Write down **all possible operators in the galaxy bias expansion**: $\delta_g = b_1 \delta_l^{(1)} + b_2 \delta_l^{(2)} + R_*^2 \partial^2 \delta_l^{(1)} + \dots$

4- Take into account the **redshift-space distortion** (RSD) effect (to subtract the contribution of the peculiar velocity of the galaxies to the cosmological redshift) *Senatore++ [arXiv:1409.1225]*

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Galaxy power spectrum

The **galaxy power spectrum** in the EFTofLSS framework:

$$P_g(k, \mu) \simeq [b_1 + f\mu^2]^2 P_m(k) = Z_1(\mu)^2 P_m(k)$$



We go from 1 to 10 free parameters

$$\begin{aligned} P_g(k, \mu) = & Z_1(\mu)^2 P_{11}(k) + 2Z_1(\mu) P_{11}(k) \left(c_{\text{ct}} \frac{k^2}{k_{\text{M}}^2} + c_{r,1} \mu^2 \frac{k^2}{k_{\text{R}}^2} + c_{r,2} \mu^4 \frac{k^2}{k_{\text{R}}^2} \right) \\ & + 2 \int \frac{d^3 q}{(2\pi)^3} Z_2(\mathbf{q}, \mathbf{k} - \mathbf{q}, \mu)^2 P_{11}(|\mathbf{k} - \mathbf{q}|) P_{11}(q) + 6Z_1(\mu) P_{11}(k) \int \frac{d^3 q}{(2\pi)^3} Z_3(\mathbf{q}, -\mathbf{q}, \mathbf{k}, \mu) P_{11}(q) \\ & + \frac{1}{\bar{n}_g} \left(c_{\epsilon,0} + c_{\epsilon}^{\text{mono}} \frac{k^2}{k_{\text{M}}^2} + 3c_{\epsilon}^{\text{quad}} \left(\mu^2 - \frac{1}{3} \right) \frac{k^2}{k_{\text{M}}^2} \right), \end{aligned}$$

Carrasco++ [arXiv:1206.2926] ; Baumann++ [arXiv:1004.2488]

Senatore [arXiv:1406.7843] ; Perko++ [arXiv:1610.09321] ; ...

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Galaxy power spectrum

The **galaxy power spectrum** in the EFTofLSS framework:

$$P_g(k, \mu) \simeq [b_1 + f\mu^2]^2 P_m(k) = Z_1(\mu)^2 P_m(k)$$



We go from 1 to 10 free parameters

$$\begin{aligned} P_g(k, \mu) = & Z_1(\mu)^2 P_{11}(k) + 2Z_1(\mu) P_{11}(k) \left(c_{\text{ct}} \frac{k^2}{k_M^2} + c_{r,1} \mu^2 \frac{k^2}{k_R^2} + c_{r,2} \mu^4 \frac{k^2}{k_R^2} \right) \\ & + 2 \int \frac{d^3 q}{(2\pi)^3} Z_2(\mathbf{q}, \mathbf{k} - \mathbf{q}, \mu) P_{11}(|\mathbf{k} - \mathbf{q}|) P_{11}(q) + 6Z_1(\mu) P_{11}(k) \int \frac{d^3 q}{(2\pi)^3} Z_3(\mathbf{q}, -\mathbf{q}, \mathbf{k}, \mu) P_{11}(q) \\ & + \frac{1}{\bar{n}_g} \left(c_{\epsilon,0} + c_{\epsilon}^{\text{mono}} \frac{k^2}{k_M^2} + 3c_{\epsilon}^{\text{quad}} \left(\mu^2 - \frac{1}{3} \right) \frac{k^2}{k_M^2} \right), \end{aligned}$$

Carrasco++ [arXiv:1206.2926] ; Baumann++ [arXiv:1004.2488]

Senatore [arXiv:1406.7843] ; Perko++ [arXiv:1610.09321] ; ...

$P_g(k, \mu)$ can be determined directly
from $P_{11}(k) = P_m^{\text{lin}}(k)$

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Galaxy power spectrum

The **galaxy power spectrum** in the EFTofLSS framework:

$$P_g(k, \mu) \simeq [b_1 + f\mu^2]^2 P_m(k) = Z_1(\mu)^2 P_m(k)$$



We go from 1 to 10 free parameters

P_{11}

$$\begin{aligned} P_g(k, \mu) = & \underbrace{Z_1(\mu)^2 P_{11}(k)} + 2Z_1(\mu)P_{11}(k) \left(c_{\text{ct}} \frac{k^2}{k_{\text{M}}^2} + c_{r,1} \mu^2 \frac{k^2}{k_{\text{R}}^2} + c_{r,2} \mu^4 \frac{k^2}{k_{\text{R}}^2} \right) \\ & + 2 \int \frac{d^3 q}{(2\pi)^3} Z_2(\mathbf{q}, \mathbf{k} - \mathbf{q}, \mu)^2 P_{11}(|\mathbf{k} - \mathbf{q}|) P_{11}(q) + 6Z_1(\mu)P_{11}(k) \int \frac{d^3 q}{(2\pi)^3} Z_3(\mathbf{q}, -\mathbf{q}, \mathbf{k}, \mu) P_{11}(q) \\ & + \frac{1}{\bar{n}_g} \left(c_{\epsilon,0} + c_{\epsilon}^{\text{mono}} \frac{k^2}{k_{\text{M}}^2} + 3c_{\epsilon}^{\text{quad}} \left(\mu^2 - \frac{1}{3} \right) \frac{k^2}{k_{\text{M}}^2} \right), \end{aligned}$$

Carrasco++ [arXiv:1206.2926] ; Baumann++ [arXiv:1004.2488]

Senatore [arXiv:1406.7843] ; Perko++ [arXiv:1610.09321] ; ...

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Galaxy power spectrum

The **galaxy power spectrum** in the EFTofLSS framework:

$$P_g(k, \mu) \simeq [b_1 + f\mu^2]^2 P_m(k) = Z_1(\mu)^2 P_m(k)$$



We go from 1 to 10 free parameters

$$\begin{aligned}
 P_g(k, \mu) = & Z_1(\mu)^2 P_{11}(k) + 2Z_1(\mu)P_{11}(k) \left(c_{\text{ct}} \frac{k^2}{k_M^2} + c_{r,1} \mu^2 \frac{k^2}{k_R^2} + c_{r,2} \mu^4 \frac{k^2}{k_R^2} \right) \\
 & + 2 \int \frac{d^3 q}{(2\pi)^3} Z_2(\mathbf{q}, \mathbf{k} - \mathbf{q}, \mu)^2 P_{11}(|\mathbf{k} - \mathbf{q}|) P_{11}(q) + 6Z_1(\mu)P_{11}(k) \int \frac{d^3 q}{(2\pi)^3} Z_3(\mathbf{q}, -\mathbf{q}, \mathbf{k}, \mu) P_{11}(q) \\
 & + \frac{1}{\bar{n}_g} \left(c_{\epsilon,0} + c_{\epsilon}^{\text{mono}} \frac{k^2}{k_M^2} + 3c_{\epsilon}^{\text{quad}} \left(\mu^2 - \frac{1}{3} \right) \frac{k^2}{k_M^2} \right),
 \end{aligned}$$

Carrasco++ [arXiv:1206.2926] ; Baumann++ [arXiv:1004.2488]

Senatore [arXiv:1406.7843] ; Perko++ [arXiv:1610.09321] ; ...

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Galaxy power spectrum

The **galaxy power spectrum** in the EFTofLSS framework:

$$P_g(k, \mu) \simeq [b_1 + f\mu^2]^2 P_m(k) = Z_1(\mu)^2 P_m(k)$$



We go from 1 to 10 free parameters

$$\begin{aligned}
 P_g(k, \mu) = & \overset{P_{11}}{Z_1(\mu)^2 P_{11}(k)} + 2Z_1(\mu)P_{11}(k) \left(c_{\text{ct}} \frac{k^2}{k_{\text{M}}^2} + c_{r,1} \mu^2 \frac{k^2}{k_{\text{R}}^2} + c_{r,2} \mu^4 \frac{k^2}{k_{\text{R}}^2} \right) \\
 & \overset{P_{22}}{+ 2 \int \frac{d^3 q}{(2\pi)^3} Z_2(\mathbf{q}, \mathbf{k} - \mathbf{q}, \mu)^2 P_{11}(|\mathbf{k} - \mathbf{q}|) P_{11}(q)} + \overset{P_{13}}{6Z_1(\mu)P_{11}(k) \int \frac{d^3 q}{(2\pi)^3} Z_3(\mathbf{q}, -\mathbf{q}, \mathbf{k}, \mu) P_{11}(q)} \\
 & + \frac{1}{\bar{n}_g} \left(c_{\epsilon,0} + c_{\epsilon}^{\text{mono}} \frac{k^2}{k_{\text{M}}^2} + 3c_{\epsilon}^{\text{quad}} \left(\mu^2 - \frac{1}{3} \right) \frac{k^2}{k_{\text{M}}^2} \right),
 \end{aligned}$$

Carrasco++ [arXiv:1206.2926] ; Baumann++ [arXiv:1004.2488]

Senatore [arXiv:1406.7843] ; Perko++ [arXiv:1610.09321] ; ...

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Galaxy power spectrum

The **galaxy power spectrum** in the EFTofLSS framework:

$$P_g(k, \mu) \simeq [b_1 + f\mu^2]^2 P_m(k) = Z_1(\mu)^2 P_m(k)$$



We go from 1 to 10 free parameters

$$\begin{aligned}
 P_g(k, \mu) = & \overset{P_{11}}{Z_1(\mu)^2 P_{11}(k)} + \overset{P_{13}^{\text{ctr}}}{2Z_1(\mu)P_{11}(k) \left(c_{\text{ct}} \frac{k^2}{k_{\text{M}}^2} + c_{r,1} \mu^2 \frac{k^2}{k_{\text{R}}^2} + c_{r,2} \mu^4 \frac{k^2}{k_{\text{R}}^2} \right)} \\
 & \overset{P_{22}}{+ 2 \int \frac{d^3 q}{(2\pi)^3} Z_2(\mathbf{q}, \mathbf{k} - \mathbf{q}, \mu)^2 P_{11}(|\mathbf{k} - \mathbf{q}|) P_{11}(q)} + \overset{P_{13}}{6Z_1(\mu)P_{11}(k) \int \frac{d^3 q}{(2\pi)^3} Z_3(\mathbf{q}, -\mathbf{q}, \mathbf{k}, \mu) P_{11}(q)} \\
 & + \frac{1}{\bar{n}_g} \left(c_{\epsilon,0} + c_{\epsilon}^{\text{mono}} \frac{k^2}{k_{\text{M}}^2} + 3c_{\epsilon}^{\text{quad}} \left(\mu^2 - \frac{1}{3} \right) \frac{k^2}{k_{\text{M}}^2} \right),
 \end{aligned}$$

Carrasco++ [arXiv:1206.2926] ; Baumann++ [arXiv:1004.2488]

Senatore [arXiv:1406.7843] ; Perko++ [arXiv:1610.09321] ; ...

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Galaxy power spectrum

The **galaxy power spectrum** in the EFTofLSS framework:

$$P_g(k, \mu) \simeq [b_1 + f\mu^2]^2 P_m(k) = Z_1(\mu)^2 P_m(k)$$

We go from 1 to 10 free parameters

$$\begin{aligned}
 P_g(k, \mu) = & Z_1(\mu)^2 P_{11}(k) + 2Z_1(\mu) P_{11}(k) \left(c_{\text{ct}} \frac{k^2}{k_M^2} + c_{r,1} \mu^2 \frac{k^2}{k_R^2} + c_{r,2} \mu^4 \frac{k^2}{k_R^2} \right) \\
 & + 2 \int \frac{d^3 q}{(2\pi)^3} Z_2(q, k - q, \mu)^2 P_{11}(|k - q|) P_{11}(q) + 6Z_1(\mu) P_{11}(k) \int \frac{d^3 q}{(2\pi)^3} Z_3(q, -q, k, \mu) P_{11}(q) \\
 & + \frac{1}{\bar{n}_q} \left(c_{\epsilon,0} + c_{\epsilon}^{\text{mono}} \frac{k^2}{k_M^2} + 3c_{\epsilon}^{\text{quad}} \left(\mu^2 - \frac{1}{3} \right) \frac{k^2}{k_M^2} \right) P_{22}^{\text{ctr}}
 \end{aligned}$$

Carrasco++ [arXiv:1206.2926] ; Baumann++ [arXiv:1004.2488]

Senatore [arXiv:1406.7843] ; Perko++ [arXiv:1610.09321] ; ...

The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Galaxy power spectrum

The **galaxy power spectrum** in the EFTofLSS framework:

$$P_g(k, \mu) \simeq [b_1 + f\mu^2]^2 P_m(k) = Z_1(\mu)^2 P_m(k)$$

We go from 1 to 10 free parameters

$$\begin{aligned}
 P_g(k, \mu) &= Z_1(\mu)^2 P_{11}(k) + (2Z_1(\mu)P_{11}(k) \left(c_{ct} \frac{k^2}{k_M^2} - c_{r,1} \mu^2 \frac{k^2}{k_R^2} - c_{r,2} \mu^4 \frac{k^2}{k_R^2} \right) \\
 &+ 2 \int \frac{d^3 q}{(2\pi)^3} Z_2(q, k-q, \mu)^2 P_{11}(|k-q|) P_{11}(q) + 6Z_1(\mu) P_{11}(k) \int \frac{d^3 q}{(2\pi)^3} Z_3(q, -q, k, \mu) P_{11}(q) \\
 &+ \frac{1}{\bar{n}_q} \left(c_{\epsilon,0} - c_{\epsilon}^{\text{mono}} \frac{k^2}{k_M^2} + 5c_{\epsilon}^{\text{quad}} \left(\mu^2 - \frac{1}{3} \right) \frac{k^2}{k_M^2} \right) P_{22}^{\text{ctr}}
 \end{aligned}$$

Carrasco++ [arXiv:1206.2926] ; Baumann++ [arXiv:1004.2488]

Senatore [arXiv:1406.7843] ; Perko++ [arXiv:1610.09321] ; ...

10 EFT parameters

○ **4 parameters** b_i ($i = 1, 2, 3, 4$) to describe the **galaxy bias** which arises from the one-loop contributions

○ **3 parameters** corresponding to **counterterms** (c_{ct} linear combination of a higher derivative bias and the dark matter sound speed, while $c_{r,1}$ and $c_{r,2}$ are the redshift-space counterterms)

○ **3 parameters** which describe **stochastic terms**



The Effective Field Theory of Large-Scale Structures (EFTofLSS)

Application to BOSS data

Multipoles of the galaxy power spectrum, obtained through a **Legendre** polynomials ($\mathcal{L}_\ell$) decomposition:

$$P_g(z, k, \mu) = \sum_{\ell \text{ even}} \mathcal{L}_\ell(\mu) P_\ell(z, k)$$

→ the two main contributions to $P_g(z, k, \mu)$ are the **monopole** ($\ell = 0$) and the **quadrupole** ($\ell = 2$)



*D'Amico++ [arXiv:1909.05271] ; Colas++ [arXiv:1909.07951]
Philcox++ [arXiv:2002.04035] ; Ivanov++ [arXiv:1909.05277]*

The Effective Field Theory of Large-Scale Structures (EFTofLSS)



Application to BOSS data

Multipoles of the galaxy power spectrum, obtained through a **Legendre** polynomials ($\mathcal{L}_\ell$) decomposition:

$$P_g(z, k, \mu) = \sum_{\ell \text{ even}} \mathcal{L}_\ell(\mu) P_\ell(z, k)$$


→ the two main contributions to $P_g(z, k, \mu)$ are the **monopole** ($\ell = 0$) and the **quadrupole** ($\ell = 2$)

Improvement in precision!




D'Amico++ [arXiv:1909.05271] ; Colas++ [arXiv:1909.07951]
Philcox++ [arXiv:2002.04035] ; Ivanov++ [arXiv:1909.05277]

BAO/ $f\sigma_8$ vs EFTofLSS



- For **eBOSS**, the error bars of Ω_m and σ_8 are reduced by a factor ~ 2.0 and ~ 1.3
- For **BOSS**, the error bars of Ω_m and h are reduced by a factor ~ 5.4 and ~ 3.2

LSS data vs Planck




ext-BAO: 6dF & MGS (SDSS) data

- The combination of eBOSS + BOSS allows to determine Ω_m and h at a **precision similar to Planck**
- The combination of LSS data remains consistent with Planck → **we can combine them!**

TS++ [arXiv:2210.14931]

LSS data combined with Planck




LSS: eBOSS + BOSS + ext-BAO + Pantheon

(Uncalibrated Supernovae)

- **Compared to Planck alone**, the constraints on Ω_m and h are **improved by $\sim 30\%$**
- σ_8 and A_s are not significantly impacted

TS++ [arXiv:2210.14931]

SDSS data vs DESI




DESI Collaboration [arXiv:2411.12021]

- **Good agreement between DESI and SDSS**
- The current constraints of DESI are similar to those of SDSS
- **Slight discrepancy between DESI and Planck** for Ω_m and h

Beyond the two-point correlation function


Power spectrum (2PCF)

$$\langle \delta_{g,r}(\vec{k}) \delta_{g,r}(\vec{k}') \rangle = (2\pi)^3 \delta_D(\vec{k} + \vec{k}') P(k, \mu)$$



Bispectrum (3PCF)


$$\langle \delta_{g,r}(\vec{k}_1) \delta_{g,r}(\vec{k}_2) \delta_{g,r}(\vec{k}_3) \rangle = (2\pi)^3 \delta_D(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) B(k_1, k_2, k_3, \mu, \phi)$$



Beyond the two-point correlation function

Power spectrum (2PCF)

$$\langle \delta_{g,r}(\vec{k}) \delta_{g,r}(\vec{k}') \rangle = (2\pi)^3 \delta_D(\vec{k} + \vec{k}') P(k, \mu)$$



$$P = P_{11} + (P_{13} + P_{13}^{ct}) + (P_{22} + P_{22}^{\epsilon})$$


10 EFT parameters

EFTofLSS computation
(Bias tracers, redshift space)

D'Amico++ [arXiv:2211.17130]
Philcox++ [arXiv:2206.02800]

Bispectrum (3PCF)

$$\langle \delta_{g,r}(\vec{k}_1) \delta_{g,r}(\vec{k}_2) \delta_{g,r}(\vec{k}_3) \rangle = (2\pi)^3 \delta_D(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) B(k_1, k_2, k_3, \mu, \phi)$$




$$B = B_{211} + (B_{321}^{(II)} + B_{321}^{(II),ct}) + (B_{411} + B_{411}^{ct}) \\ + (B_{222} + B_{222}^{\epsilon}) + (B_{321}^{(I)} + B_{321}^{(I),\epsilon})$$

41 EFT parameters


Beyond the two-point correlation function

Analysis of BOSS (LRG) with $P + B$, for $0.01 < k < 0.23 \text{ [} h\text{Mpc}^{-1}\text{]}$



D'Amico++ [arXiv:2206.08327]

- **Good agreement** between Planck and BOSS (again)
- 13 % improvement on Ω_m , 18 % on h and 30 % on σ_8 thanks to the bispectrum
- **30 % of improvement on w**



(Base = Planck + PanPlus + ext-BAO)

Spaar++ [arXiv:2312.15164]

II- Constraining dark energy history with the EFTofLSS

Based on:

- Z. Lu, **TS** and P. Zhang [arXiv:2503.04602]


Preference for evolving dark energy in light of the galaxy bispectrum

General context: preference for evolving dark energy


Extension of the Λ CDM model, where the dark energy equation-of-state is parametrized by

$$w_{\text{DE}} = \frac{P(z)}{\rho(z)} = w_0 + w_a(1 - a)$$

- **Preference for evolving dark energy over Λ CDM at 3.1σ for DESI BAO DR2 + CMB**
- 2.8σ , 3.8σ and 4.2σ when we add Pantheon+, Union3 or DES Y5
- Same level of preference for the DESI Y1 full-shape analysis (EFTofLSS)




DESI Collaboration [arXiv:2503.14738]



DESI Collaboration [arXiv:2411.12021]


Main results



Dataset	w_0	w_a	$p > \Lambda$	$\Delta\chi^2$
<i>Planck</i> + PanPlus + DESI-BAO	-0.821 ± 0.063	-0.77 ± 0.27	2.5σ	-8.9
<i>Planck</i> + PanPlus + ext-BAO + EFTBOSS	$-0.809^{+0.053}_{-0.061}$	$-0.72^{+0.28}_{-0.25}$	2.8σ	-10.3
<i>Planck</i> + PanPlus + DESI-BAO + EFTBOSS/ r_s^{marg}	-0.789 ± 0.050	$-0.80^{+0.25}_{-0.22}$	3.7σ	-16.9
<i>Planck</i> + Union3 + DESI-BAO + EFTBOSS/ r_s^{marg}	-0.677 ± 0.075	$-1.14^{+0.33}_{-0.29}$	3.8σ	-17.7
<i>Planck</i> + DESY5 + DESI-BAO + EFTBOSS/ r_s^{marg}	-0.726 ± 0.060	$-1.00^{+0.30}_{-0.25}$	4.4σ	-22.9

- We obtain a **2.8σ preference** for the w_0w_a CDM model with pre-DESI data, namely *Planck* + PanPlus + EFTBOSS
- **The preference increases to $3.6\sigma / 3.9\sigma$** when considering Union3 / DES Y5
- When we add DESI BAO Y1 on top, we have a $3.7\sigma - 4.4\sigma$



Breaking degeneracies between SN1a and galaxy clustering



- The combinations of *Planck* with either PanPlus or EFTBOSS **do not show any preference** for evolving dark energy
- We we combine PanPlus with EFTBOSS we **break some degeneracies** between $h - w_0/w_a$ and $\Omega_m - w_0/w_a$
- This is the first time that we have preference for evolving dark energy without data from DESI!!



Impact of the bispectrum contribution

- Without the bispectrum, **we do not get a preference** over Λ (1.5σ)
- With the bispectrum, **we get a $2.8 - 3.9\sigma$ preference** over Λ (shift of $\sim 0.5\sigma$ + reduction of 30 % of $w_0 - w_a$)
- Deviations from Λ , together with improvement in the constraints, **start to be significant when** $k_{\max} = 0.14 \, h\text{Mpc}^{-1}$, where $k_{\max}$ is the highest Fourier mode included in the analysis of the bispectrum




Impact of the bispectrum contribution

- Without the bispectrum, **we do not get a preference** over Λ (1.5σ)
- With the bispectrum, **we get a $2.8 - 3.9\sigma$ preference** over Λ (shift of $\sim 0.5\sigma$ + reduction of 30 % of $w_0 - w_a$)
- Deviations from Λ , together with improvement in the constraints, **start to be significant when** $k_{\max} = 0.14 \, h\text{Mpc}^{-1}$, where $k_{\max}$ is the highest Fourier mode included in the analysis of the bispectrum



Impact of the bispectrum contribution

- Without the bispectrum, **we do not get a preference** over Λ (1.5σ)
- With the bispectrum, **we get a $2.8 - 3.9\sigma$ preference** over Λ (shift of $\sim 0.5\sigma$ + reduction of 30 % of $w_0 - w_a$)
- Deviations from Λ , together with improvement in the constraints, **start to be significant when** $k_{\max} = 0.14 \, h\text{Mpc}^{-1}$, where $k_{\max}$ is the highest Fourier mode included in the analysis of the bispectrum




The preference for $w_0 w_a$ comes from the one-loop bispectrum contribution!

III- Constraining the initial conditions with the EFTofLSS

Based on:

- R. Calderon, **TS**, A. Shafieloo and D. K. Hazra [arXiv:2504.06183]
Primordial Features in light of the Effective Field Theory of Large Scale Structure

EFTofLSS meets the standard initial conditions




$$P_{\phi,0} = \frac{2\pi^2 A_s}{k^3} \left(\frac{k}{k_*} \right)^{n_s-1}$$

- The *Planck* constraints on $\ln(10^{10}A_s)$ and n_s are respectively **8 and 10 times better** than those of BOSS + eBOSS (LSS)
- **Question:** What is the sensitivity of the current large-scale structure data on inflationary dynamics beyond the standard “slow-roll” regime?

ETFofLSS meets primordial global oscillations


Linear oscillations

$$P_{\phi,0} \rightarrow P_{\phi,0} \left[1 + A_{\text{lin}} \sin(\omega_{\text{lin}} \frac{k}{k_*} + \phi_{\text{lin}}) \right]$$



Logarithmic oscillations


$$P_{\phi,0} \rightarrow P_{\phi,0} \left[1 + A_{\text{log}} \sin(\omega_{\text{log}} \ln \left(\frac{k}{k_*} \right) + \phi_{\text{log}}) \right]$$



ETFofLSS meets primordial global oscillations

Linear oscillations

$$P_{\phi,0} \rightarrow P_{\phi,0} \left[1 + A_{\text{lin}} \sin\left(\omega_{\text{lin}} \frac{k}{k_*} + \phi_{\text{lin}}\right) \right]$$




Logarithmic oscillations

- **Very good consistency** between Nbody simulations and EFTofLSS prediction
- Non-linearities introduce a **damping** of the oscillatory features
- The damping is **enhanced for small redshifts**
- (e)BOSS data allow to explore the regime $k \sim 0.1 - 0.2 h/\text{Mpc}$, **which is not accessible by Planck**

ETFofLSS meets primordial global oscillations

Linear oscillations

$$P_{\phi,0} \rightarrow P_{\phi,0} \left[1 + A_{\text{lin}} \sin\left(\omega_{\text{lin}} \frac{k}{k_*} + \phi_{\text{lin}}\right) \right]$$




$$A_{\text{lin}}^{\text{LSS}} < 0.031$$
$$A_{\text{lin}}^{\text{CMB}} < 0.019$$

- The LSS constraint on A_{lin} is **only 1.6 times weaker** than that of the CMB
- When we combine LSS with CMB, **we improve the constraints by 25 %** compared to CMB
- ϕ_{lin} and ω_{lin} **are not (yet?) constrained** by CMB and LSS data

ETFofLSS meets primordial global oscillations

Logarithmic oscillations

$$P_{\phi,0} \rightarrow P_{\phi,0} \left[1 + A_{\log} \sin(\omega_{\log} \ln \left(\frac{k}{k_*} \right) + \phi_{\log}) \right]$$



$$A_{\log}^{\text{LSS}} < 0.024$$
$$A_{\text{lin}}^{\text{CMB}} < 0.024$$


- We obtain **the same constraint** on $A_{\log}$ between CMB and LSS!
- When we combine LSS with CMB, **we improve the constraints by 50 %** compared to CMB
- $\phi_{\log}$ and $\omega_{\log}$ **are not (yet?) constrained** by CMB and LSS data

ETFofLSS meets primordial local oscillations

One Spectrum

$$P_{\phi,0} \rightarrow P_{\phi,0} \left[1 + \alpha \frac{\sin(\omega(k - k_0))}{1 + \beta(k - k_0)^4} \right]$$

Hazra++ [arXiv:2201.12000]



$$\alpha^{\text{CMB}} < 0.094$$

$$\alpha^{\text{LSS}} < 0.034$$

$$\alpha^{\text{CMB+LSS}} < 0.035$$

- LSS data are **3 times more constraining** than CMB data!
- EFTofLSS analysis applied to (e)BOSS data is able to **disfavor localized primordial oscillations** that are favored by *Planck*

General conclusion

Part I: *How to obtain the most robust and optimised constraints on Λ CDM cosmological parameters from large-scale structure data?*

- EFTofLSS allows **much larger constraints** than the BAO/ $f\sigma_8$ information on the Λ CDM model and its extensions
- Constraints from LSS data are **competitive with CMB data**
- One-loop bispectrum analysis (up to $k_{\text{max}} = 0.23 \, h\text{Mpc}^{-1}$) **adds constraining power** (up to 30 %)

General conclusion

Part I: *How to obtain the most robust and optimised constraints on Λ CDM cosmological parameters from large-scale structure data?*

- EFTofLSS allows **much larger constraints** than the BAO/ $f\sigma_8$ information on the Λ CDM model and its extensions
- Constraints from LSS data are **competitive with CMB data**
- One-loop bispectrum analysis (up to $k_{\text{max}} = 0.23 \, h\text{Mpc}^{-1}$) **adds constraining power** (up to 30 %)

Part II: *How can clustering data be used to constrain the nature of dark energy and the initial conditions?*

- For *Planck* + SN1a + EFTBOSS **we obtain a $2.8 - 3.9\sigma$ preference for w_0w_a over Λ** (good check of the DESI results)
- This preference **disappears when we do not include the one-loop bispectrum** (it is important to consider it!)
- EFTofLSS offers a **robust and fast way to put strong constraints on primordial features** with galaxy clustering

General conclusion

Part I: *How to obtain the most robust and optimised constraints on Λ CDM cosmological parameters from large-scale structure data?*


- EFTofLSS allows **much larger constraints** than the BAO/ $f\sigma_8$ information on the Λ CDM model and its extensions
- Constraints from LSS data are **competitive with CMB data**
- One-loop bispectrum analysis (up to $k_{\text{max}} = 0.23 \, h\text{Mpc}^{-1}$) **adds constraining power** (up to 30 %)

Part II: *How can clustering data be used to constrain the nature of dark energy and the initial conditions?*

- For *Planck* + SN1a + EFTBOSS **we obtain a $2.8 - 3.9\sigma$ preference for w_0w_a over Λ** (good check of the DESI results)
- This preference **disappears when we do not include the one-loop bispectrum** (it is important to consider it!)
- EFTofLSS offers a **robust and fast way to put strong constraints on primordial features** with galaxy clustering

Thanks for your attention!

LSS data vs Planck



TS++ [arXiv:2210.14931]

- **eBOSS, BOSS and Planck are consistent** at $\lesssim 1.8\sigma$ on all cosmological parameters
 - The h and σ_8 Planck values are **in-between** those of BOSS and eBOSS
- **there is no tension between Planck and BOSS/eBOSS**

On the consistency of EFTofLSS

Presentation of the problem

There are **several codes** in the literature with **different parametrizations**:

→ **CLASS PT** with the **EC** parametrization *Chudaykin++ [arXiv:2004.10607]*


→ **PyBird** with the **WC** parametrization *D'Amico++ [arXiv:2003.07956]*

(+ **Velocileptors** + **CLASS-OneLoop**)

Chen++ [arXiv:2005.00523] ; Linde++ [arXiv:2402.09778]

→ these codes use **different sets of priors** on EFT parameters

D'Amico++ [arXiv:1909.05271]
Philcox++ [arXiv:2002.04035]



TS++ [arXiv:2208.05929]

On the consistency of EFTofLSS

Presentation of the problem

There are **several codes** in the literature with **different parametrizations**:

→ **CLASS PT** with the **EC** parametrization *Chudaykin++ [arXiv:2004.10607]*


→ **PyBird** with the **WC** parametrization *D'Amico++ [arXiv:2003.07956]*

(+ **Velocileptors** + **CLASS-OneLoop**)

Chen++ [arXiv:2005.00523] ; Linde++ [arXiv:2402.09778]

→ these codes use **different sets of priors** on EFT parameters

D'Amico++ [arXiv:1909.05271]
Philcox++ [arXiv:2002.04035]




TS++ [arXiv:2208.05929]

Data, theoretical **parametrizations** and **codes** are supposed to be **equivalent**: what is going on?

On the consistency of EFTofLSS


The EFT prior issue



TS++ [arXiv:2208.05929]

On the consistency of EFTofLSS

The EFT prior issue



TS++ [arXiv:2208.05929]

Prior effects

- **The prior weight effect:** if the region allowed by the prior is far from the true value of a parameter, then the posterior and bestfit will be shifted from the true value
- **The prior volume effect:** a posterior depends on the volume enclosed by the priors $\implies$ large parameter regions are emphasized compared to smaller regions

Bayes' theorem:


$$P \propto \mathcal{L} \times p$$

On the consistency of EFTofLSS


Profile likelihood

Advantage: frequentist analysis is **independent of priors** and therefore of projection effects

Disadvantage: the data prefers several EFT parameters to take on **extreme values**, possibly breaking the perturbative nature of the theory



Brinch, Herold, TS++ [arXiv:2309.04468]




On the consistency of EFTofLSS

Profile likelihood


Advantage: frequentist analysis is **independent of priors** and therefore of projection effects

Disadvantage: the data prefers several EFT parameters to take on **extreme values**, possibly breaking the perturbative nature of the theory



The low value of σ_8 is due to prior effects!



Brinch, Herold, TS++ [arXiv:2309.04468]



EFTofLSS applied to eBOSS QSO data

- **343 708 quasars** selected in the redshift range $0.8 < z < 2.2$
- $z_{\text{eff}} = 1.5$
- 2 skycuts: NGC and SGC

eBOSS Collaboration
[arXiv:2007.08991]




TS++ [arXiv:2210.14931]

EFTofLSS applied to eBOSS QSO data


- **343 708 quasars** selected in the redshift range $0.8 < z < 2.2$
- $z_{\text{eff}} = 1.5$
- 2 skycuts: NGC and SGC

eBOSS Collaboration
[arXiv:2007.08991]



cut-off scale $k_{\text{max}} = 0.24 h \cdot \text{Mpc}^{-1}$

TS++ [arXiv:2210.14931]



Determination of the cut-off scale $k_{\max}$ of the one-loop prediction

The next-to-next-to-leading order (NNLO) terms

At **one-loop order**, the galaxy power spectrum reads:

$$P_g(k, \mu) = Z_1(\mu)^2 P_{11}(k) + 2Z_1(\mu) P_{11}(k) \left(c_{\text{ct}} \frac{k^2}{k_{\text{M}}^2} + c_{r,1} \mu^2 \frac{k^2}{k_{\text{M}}^2} + c_{r,2} \mu^4 \frac{k^2}{k_{\text{M}}^2} \right) \\ + 2 \int \frac{d^3 q}{(2\pi)^3} Z_2(\mathbf{q}, \mathbf{k} - \mathbf{q}, \mu)^2 P_{11}(|\mathbf{k} - \mathbf{q}|) P_{11}(q) + 6Z_1(\mu) P_{11}(k) \int \frac{d^3 q}{(2\pi)^3} Z_3(\mathbf{q}, -\mathbf{q}, \mathbf{k}, \mu) P_{11}(q) \\ + \frac{1}{\bar{n}_g} \left(c_{\epsilon,0} + c_{\epsilon,1} \frac{k^2}{k_{\text{M}}^2} + c_{\epsilon,2} f \mu^2 \frac{k^2}{k_{\text{M}}^2} \right),$$

One can add the **NNLO terms** (i.e., the dominant two-loop terms):

$$P_{\text{NNLO}}(k, \mu) = \frac{1}{4} b_1 (c_{r,4} b_1 + c_{r,6} \mu^2) \mu^4 \frac{k^4}{k_{\text{R}}^4} P_{11}(k)$$

Zhang++ [arXiv:2110.07539]

If the contribution of $P_{\text{NNLO}}(k, \mu)$ becomes **too large**, the one-loop prediction is **not accurate enough** → this determines the **cut-off scale** $k_{\max}$ of the prediction

Determination of the cut-off scale k_{max} of the one-loop prediction

The next-to-next-to-leading order (NNLO) terms

At **one-loop order**, the galaxy power spectrum reads:

$$P_g(k, \mu) = Z_1(\mu)^2 P_{11}(k) + 2Z_1(\mu) P_{11}(k) \left(c_{\text{ct}} \frac{k^2}{k_{\text{M}}^2} + c_{r,1} \mu^2 \frac{k^2}{k_{\text{M}}^2} + c_{r,2} \mu^4 \frac{k^2}{k_{\text{M}}^2} \right) \\ + 2 \int \frac{d^3 q}{(2\pi)^3} Z_2(\mathbf{q}, \mathbf{k} - \mathbf{q}, \mu)^2 P_{11}(|\mathbf{k} - \mathbf{q}|) P_{11}(q) + 6Z_1(\mu) P_{11}(k) \int \frac{d^3 q}{(2\pi)^3} Z_3(\mathbf{q}, -\mathbf{q}, \mathbf{k}, \mu) P_{11}(q) \\ + \frac{1}{\bar{n}_g} \left(c_{\epsilon,0} + c_{\epsilon,1} \frac{k^2}{k_{\text{M}}^2} + c_{\epsilon,2} f \mu^2 \frac{k^2}{k_{\text{M}}^2} \right),$$

One can add the **NNLO terms** (i.e., the dominant two-loop terms):

$$P_{\text{NNLO}}(k, \mu) = \frac{1}{4} b_1 (c_{r,4} b_1 + c_{r,6} \mu^2) \mu^4 \frac{k^4}{k_{\text{R}}^4} P_{11}(k)$$

Zhang++ [arXiv:2110.07539]

2 new EFT parameters

If the contribution of $P_{\text{NNLO}}(k, \mu)$ becomes **too large**, the one-loop prediction is **not accurate enough** → this determines the **cut-off scale** k_{max} of the prediction

Determination of the cut-off scale k_{max} of the one-loop prediction

The next-to-next-to-leading order (NNLO) terms

At **one-loop order**, the galaxy power spectrum reads:

$$P_g(k, \mu) = Z$$
$$+ 2 \int \frac{d^3 q}{(2\pi)^3}$$
$$+ \frac{1}{\bar{n}_g} \left(c_{\epsilon,0} + \right)$$

One can add the **NNLO**

If the contribution of
accurate enough →

	$\Delta_{\text{NNLO}}^{\text{shift}} / \sigma_{\text{stat}}^{\text{eBOSS}}$			
$k_{\text{max}} [h \text{ Mpc}^{-1}]$	0.21	0.24	0.27	0.30
Ω_{m}	0.09	-0.01	0.09	-0.49
h	-0.09	-0.05	0.09	0.21
σ_8	-0.01	-0.02	0.22	-0.02
b_1^{N}	0.02	-0.01	0.01	-0.10
c_2^{N}	0.07	0.07	0.62	0.38
b_1^{S}	0.05	-0.01	0.05	-0.01
c_2^{S}	0.06	-0.07	0.49	0.35


$$P_g(k, \mu) P_{11}(q)$$

[arXiv:2110.07539]

ction is **not**
on

Determination of the cut-off scale $k_{\max}$ of the one-loop prediction

The EZmock



TS++ [arXiv:2210.14931]

- **EZmock**: mocks that are built to simulate eBOSS observational characteristics

Chuang++ [arXiv:1409.1124]


- Up to $k_{\max} = 0.24h \text{ Mpc}^{-1}$, the best-fit values of the cosmological parameters are shifted with respect to the truth of the simulations by $\lesssim 1/3 \cdot \sigma$

Extensions to Λ CDM: total neutrino mass $\sum m_\nu$

- The LSS constraint derived in this work is **only** $\sim 10\%$ **weaker than the Planck constraint** ($\sum m_\nu < 0.241\text{eV}$)
- The EFT analysis **significantly improves the constraints** on $\sum m_\nu$ (by a factor of ~ 18) over the conventional BAO/ $f\sigma_8$ analysis ($\sum m_\nu < 4.84\text{eV}$)
Palanque-Delabrouille++ [arXiv:1911.09073]
- This analysis **disfavors the inverse hierarchy** at $\sim 2.2\sigma$ & is **competitive to the Lyman- α constraints**

LSS:
 $\sum m_\nu < 0.274\text{eV}$

LSS+Planck:
 $\sum m_\nu < 0.093\text{eV}$




Extensions to Λ CDM: total neutrino mass $\sum m_\nu$

- The LSS constraint derived in this work is **only** $\sim 10\%$ **weaker than the Planck constraint** ($\sum m_\nu < 0.241\text{eV}$)
- The EFT analysis **significantly improves the constraints** on $\sum m_\nu$ (by a factor of ~ 18) over the conventional BAO/ $f\sigma_8$ analysis ($\sum m_\nu < 4.84\text{eV}$)
Palanque-Delabrouille++ [arXiv:1911.09073]
- This analysis **disfavors the inverse hierarchy** at $\sim 2.2\sigma$ & is **competitive to the Lyman- α constraints**

DESI++ [arXiv:2411.12022]

LSS:
 $\sum m_\nu < 0.274\text{eV}$
 $\sum m_\nu < 0.300\text{eV}$
LSS+Planck:
 $\sum m_\nu < 0.093\text{eV}$
 $\sum m_\nu < 0.071\text{eV}$



Extensions to Λ CDM: curvature density fraction Ω_k


- With LSS data only, we find Ω_k **compatible with zero curvature** at 1.3σ
- The EFT analysis **significantly improves the constraints** on Ω_k by $\sim 50\%$ compared to the conventional BAO/ $f\sigma_8$ analysis
- The combination of LSS and Planck leads to a **strong constraint** and excludes the (slightly favored) negative values of Ω_k

LSS:

$$\Omega_k = -0.039^{+0.028}_{-0.029}$$

LSS+Planck:

$$\Omega_k = 0.0008^{+0.0018}_{-0.0017}$$




Extensions to Λ CDM: dark energy equation of state w_0

- With the LSS data only, we find **no evidence for a universe with $w_0 \neq -1$**
- The EFT analysis **improves the constraints** on w_0 by $\sim 20\%$ compared to the conventional BAO/ $f\sigma_8$ analysis
- The addition of LSS data select values of w_0 close to -1 , located in the 2σ region reconstructed from Planck data

LSS:
 $w_0 = -1.038 \pm 0.041$

LSS+Planck:
 $w_0 = -1.039 \pm 0.029$




Extensions to Λ CDM: effective number of relativistic species N_{eff}

- The value of ΔN_{eff} is **compatible with the standard model**
- Unlike EFTofLSS, the **conventional BAO/ $f\sigma_8$ analysis is unable to constrain this parameter**
- The addition of the LSS data **improves** the results of Planck alone by $\sim 25\%$

LSS:
 $\Delta N_{\text{eff}} = 0.40^{+0.44}_{-0.91}$

LSS+Planck:
 $\Delta N_{\text{eff}} = -0.07^{+0.15}_{-0.16}$




Dark energy equation of state $w_0 \geq -1$

- One can see that this new prior shifts the 2D posteriors inferred from the LSS data in a non-negligible way, while it remains globally stable for the LSS + Planck
- For these analyses, $\Delta\chi^2 = 0$ with respect to Λ CDM, since we obtain best-fit values of $w_0 = -1$

LSS:
 $w_0 < -0.932$


LSS+Planck:
 $w_0 < -0.965$




General context: preference for evolving dark energy

The w_0w_a CDM model allows


- to reduce the **tension between Planck and DESI BAO** in the $H_0r_d - \Omega_m$ plane
- to provide a **better fit of the low- z distance modulus** data from supernovae
- to provide of **the bump at $z \sim 0.7$ in the DESI DR2** (isotropic) BAO



DESI Collaboration [arXiv:2503.14738]



DESI Collaboration [arXiv:2503.24343]



DESI Collaboration [arXiv:2503.14743]

Dark energy perturbations

In this work, we consider two parametrizations of the dark energy perturbations

Smooth quintessence ($c_s^2 \rightarrow 1$)

No perturbation (like DESI results)

Clustering quintessence ($c_s^2 \rightarrow 0$)

Dark energy clusters exactly like dark matter

Creminelli++ [arXiv:0811.0827]

Lewandowski++ [arXiv:1611.07966]

D'Amico++ [arXiv:2012.07554]

Dark energy perturbations

In this work, we consider two parametrizations of the dark energy perturbations

Smooth quintessence ($c_s^2 \rightarrow 1$)

No perturbation (like DESI results)

$$\partial^2 \Phi = \frac{\bar{\rho}_m a^2}{2M_{\text{pl}}^2} \delta_m$$

Poisson

Clustering quintessence ($c_s^2 \rightarrow 0$)

Dark energy clusters exactly like dark matter

$$\partial^2 \Phi = \frac{\bar{\rho}_m a^2}{2M_{\text{pl}}^2} \delta_A$$

Creminelli++ [arXiv:0811.0827]

Lewandowski++ [arXiv:1611.07966]

$$\delta_A = \delta_m + \delta_d$$

$$\delta_d = \frac{1+w}{c_s^2} \frac{\bar{\rho}_d}{\bar{\rho}_m} (\dot{\pi} - \Phi)$$

D'Amico++ [arXiv:2012.07554]

Dark energy perturbations

In this work, we consider two parametrizations of the dark energy perturbations

Smooth quintessence ($c_s^2 \rightarrow 1$)

No perturbation (like DESI results)

$$\partial^2 \Phi = \frac{\bar{\rho}_m a^2}{2M_{\text{pl}}^2} \delta_m$$

$$\dot{\delta}_m + \frac{1}{a} \theta = -\frac{1}{a} \partial_i (\delta v^i)$$

Poisson

Continuity

Clustering quintessence ($c_s^2 \rightarrow 0$)

Dark energy clusters exactly like dark matter

$$\partial^2 \Phi = \frac{\bar{\rho}_m a^2}{2M_{\text{pl}}^2} \underline{\delta_A}$$

$$\underline{\dot{\delta}_A} + \frac{1}{a} \underline{C(a)} \theta = -\frac{1}{a} \partial_i (\underline{\delta_A} v^i)$$

Creminelli++ [arXiv:0811.0827]

Lewandowski++ [arXiv:1611.07966]

$$\delta_A = \delta_m + \delta_d$$

$$\delta_d = \frac{1+w}{c_s^2} \frac{\bar{\rho}_d}{\bar{\rho}_m} (\dot{\pi} - \Phi)$$

$$C(a) = 1 + (1+w) \frac{\bar{\rho}_m}{\bar{\rho}_d}$$

D'Amico++ [arXiv:2012.07554]

Dark energy perturbations

In this work, we consider two parametrizations of the dark energy perturbations

Smooth quintessence ($c_s^2 \rightarrow 1$)

No perturbation (like DESI results)

$$\partial^2 \Phi = \frac{\bar{\rho}_m a^2}{2M_{\text{pl}}^2} \delta_m$$

$$\dot{\delta}_m + \frac{1}{a} \theta = -\frac{1}{a} \partial_i (\delta v^i)$$

$$\dot{\theta} + H\theta + \frac{1}{a} \partial^2 \Phi = -\frac{1}{a} \partial_i (v^j \partial_j v^i)$$

Poisson

Continuity

Euler

Clustering quintessence ($c_s^2 \rightarrow 0$)

Dark energy clusters exactly like dark matter

$$\partial^2 \Phi = \frac{\bar{\rho}_m a^2}{2M_{\text{pl}}^2} \underline{\delta_A}$$

$$\underline{\dot{\delta}_A} + \frac{1}{a} \underline{C(a)} \theta = -\frac{1}{a} \partial_i (\underline{\delta_A} v^i)$$

$$\dot{\theta} + H\theta + \frac{1}{a} \partial^2 \Phi = -\frac{1}{a} \partial_i (v^j \partial_j v^i)$$

Creminelli++ [arXiv:0811.0827]

Lewandowski++ [arXiv:1611.07966]


$$\delta_A = \delta_m + \delta_d$$

$$\delta_d = \frac{1+w}{c_s^2} \frac{\bar{\rho}_d}{\bar{\rho}_m} (\dot{\pi} - \Phi)$$

$$C(a) = 1 + (1+w) \frac{\bar{\rho}_m}{\bar{\rho}_d}$$


D'Amico++ [arXiv:2012.07554]

Dark energy perturbation



- The evidence for $w_0 w_a$ is 2.6σ for **smooth quintessence** and 2.8σ for **clustering quintessence**
- Constraints on (w_0, w_a) depend not only on the background but also **on the propagation of dark energy perturbations**, affecting clustering differently

EFT prior effects



Our analysis is not affected by EFT prior effects:

→ the constraints in the $w_0 - w_a$ plane are unchanged the EFT priors are multiplied by 2

→ the relative shifts between the maximum a posteriori and the posterior means are $\lesssim 0.2\sigma$

Prior effects

→ **The prior weight effect:** if the region allowed by the prior is far from the true value of a parameter, then the posterior and bestfit will be shifted from the true value

→ **The prior volume effect:** a posterior depends on the volume enclosed by the priors $\implies$ large parameter regions are emphasized compared to smaller regions

Bayes' theorem:
 $P \propto \mathcal{L} \times p$


TS++ [arXiv:2208.05929]

Dark energy history

Where does the information to constrain $w(z)$ come from?

General observations

- $w(z)$ is **most constrained** at $z \sim 0.25$
- The **preference for $w_0 > -1$** is driven by w_1^z , while **the preference for $w_a < 0$** is driven by the bins spanning from $z \sim 0.6$ to $z \sim 2.3$
- **Beyond $z \sim 2.3$, the data has no constraining power on w_0 and w_a** (since $\Omega_{\text{DE}}(z)/\Omega_{\text{m}}(z) \rightarrow 0$)




Model independent reconstruction of $w(z)$ based on a general piecewise parametrization $w_i^z, i \in [1,9]$

Dark energy history

Where does the information to constrain $w(z)$ come from?


Conclusions

- So **we should not consider** the (w_0, w_a) parameterization after $z \sim 2.3$
- From the model-independent reconstruction, we find a **1.7σ deviation from Λ**
- **The phantom crossing occurs at $z \sim 0.4$** (which is below $z \sim 2.3$!), and we find a **2.3σ preference** for a phantom phase within the $w_0 w_a$



Model independent reconstruction of $w(z)$ based on a general piecewise parametrization $w_i^z, i \in [1,9]$


Variation of the supernovae data



Evolving dark energy and curvature


$$\Omega_k = -0.0014 \pm 0.0027$$

$$\Omega_k = -0.0027 \pm 0.0025$$




- The constraints on $w_0 - w_a$ are weakened because of the degeneracy with Ω_k
- Yet, the preference over Λ remains similar as in the flat case

Exact-time dependance





Dark energy pivot scale




- We define $w_* = w_0 + (1 - a_*)w_a$, such that $w(a) = w_* + (a_* - a)w_a$
- The pivot epoch a_* is chosen such that w_* has minimal variance (w_* is the principal component of $w_0 - w_a$)
- We obtain $z_* = 0.25$

Stability condition



- *K-essence model* (in orange): $w < -1$ is prohibited. It remains consistent with Λ
- $c_s^2 \rightarrow 1$ *limit*: stable for $w < -1$ when additional EFTofDE parameters stabilizing the gradient are present. This is the case with α_B , the braiding parameter, for which we find a 2.9σ preference over Λ (α_B is detected at 2σ)
- $c_s^2 \rightarrow 0$ *limit*: stable if $w > -1$ when stabilized using higher-order operators



D'Amico++ [arXiv:2012.07554]