




Cosmology with CMB and galaxy surveys from space




Giulio Fabbian

Institute d'Astrophysique Spatiale




Open questions in cosmology as of 2025



- What is dark matter? Do **massive neutrinos** contribute?
- What is the nature of **dark energy**?
- Which mechanism driving primordial **inflation**?


Cosmological probes: CMB, large-scale structures

Cosmic Microwave Background (CMB)




Planck
collaboration
(2018)


Intensity and polarization



- Constituents and composition of the Universe from early times.



- Imprint of structures at different times



Galaxy surveys




Desi
Collaboration

Galaxy position, distances, shapes







- Perturbations growth across-time and spatial distribution.
- More information on dark energy/neutrinos.



Is CMB “the” CMB?

- CMB is a snapshot of the universe at $z \sim 1100$... plus lots of other things!
 - Imprint of astrophysical objects / late time physics (CIB, ISW, SZ).
 - Gravitational lensing.



- ICM (g)astrophysics, cosmology (SZ)
- Star formation, cosmology (CIB)
- Extragalactic astronomy, Galaxy evolution

The CMB lensing potential



- Unbiased tracers of the whole integrated matter distribution along the line of sight.

$$\phi(\boldsymbol{\theta}) = -2 \int_0^{\chi_s} \frac{D_A(\chi_s - \chi')}{D_A(\chi_s) D_A(\chi')} \Psi(\boldsymbol{\theta}, \chi') d\chi' \quad \mathbf{d} = \nabla \phi$$

Notation Warning!


$$\kappa \propto -\nabla^2 \phi \rightarrow \delta$$

- Sensitive to total matter distribution $\sigma_8 \Omega_m^{0.25}$ at $z \sim 0.6-5$ on mildly non-linear scales.



CMB probes and cross-correlation

- CMB lensing: integrated, 2D projected, unbiased probe of matter.
- Integrated = overlaps with all the matter in the universe.
- Synergies with any other probe of matter e.g. LSS surveys.
- Reduces systematics



$$\kappa_{gal,obs} = \kappa_{gal} + s_{gal} \quad \kappa_{CMB,obs} = \kappa_{CMB} + s_{CMB}$$

$$\langle \kappa_{gal,obs} \kappa_{gal,obs} \rangle = \langle \kappa_{gal} \kappa_{gal} \rangle + \langle s_{gal} s_{gal} \rangle + \langle \kappa_{gal} s_{gal} \rangle$$

$$\langle \kappa_{CMB,obs} \kappa_{CMB,obs} \rangle = \langle \kappa_{CMB} \kappa_{CMB} \rangle + \langle s_{CMB} s_{CMB} \rangle + \langle \kappa_{CMB} s_{CMB} \rangle$$

$$\langle \kappa_{CMB,obs} \kappa_{gal,obs} \rangle = \langle \kappa_{CMB} \kappa_{gal} \rangle + \cancel{\langle s_{CMB} s_{gal} \rangle} + \cancel{\langle \kappa_{CMB} s_{gal} \rangle} + \cancel{\langle \kappa_{gal} s_{CMB} \rangle}$$



Cross-correlation basis and examples

$$C_L^{AB} \approx \int \frac{d\chi}{\chi^2} W_A(\chi) W_B(\chi) P_\delta \left(k = \frac{L + 1/2}{\chi}, z(\chi) \right)$$

CMB lensing



Photometric
Galaxy clustering

Galaxy lensing

$$W_g(\chi) = b_g(z) \frac{1}{n} \frac{dn}{dz} \frac{dz}{d\chi}$$


$$W_\kappa(\chi, \chi_s) = \gamma(\chi) \chi^2 \left(\frac{1}{\chi} - \frac{1}{\chi_s} \right) \Theta(\chi_s - \chi)$$

$$D_G \approx (C_\ell^{\kappa g})^2 / C_\ell^{gg} \approx \hat{\theta} \in \{C_\ell^{\kappa g}, C_\ell^{gg}\}$$



Cosmological probes and cross-correlations


- Different probes test different scales and redshift (and different systematics).
- Need to assess if growth tension is real or effect of non-linear physics.



CMB cross-correlations for cosmology




- Current data:
 - marginalizing over systematics at limited cost
 - limited additional power!
- Situation bound to change very soon:
 - New CMB data on sky!
 - New galaxy surveys on sky!



AOB: Simons Observatory first light achieved!

Small Aperture Telescopes (SATs), on sky!



- B-modes $r \sim 0.003$
- 3 there, more soon!
- Commissioning data!

Simons Observatory collaboration: science goals and forecasts (2019)




Large Aperture Telescope (LAT) in Feb 2025!

- Small-scale E/B-modes
- Lensing and SZ
- $N_{\text{eff}} \sim 0.07$
- $\sum m_\nu \sim 50 \text{ meV}$




Sneak preview

- Just ~2 weeks of science commissioning data....
- Includes foreground hardening




EUCLID

RESULTS SO FAR



VISTA VMC J-band




2MASS J-band


Credit: Ralf Bender & Ross Collins using public data from Euclid and VISTA

EUCLID

RESULTS SO FAR



VISTA VMC J-band



Euclid NISP Test Image

Credit: Ralf Bender & Ross Collins using public data from Euclid and VISTA

What to expect: synergies between surveys.




Euclid Early Release Observation image

Euclid Q1 release





Rubin 1st wide image




DESI Legacy Imaging Survey

Romanowsky+(2025)



In the meantime... Gaia!

- Photometry, astrometry, slitless spectroscopy with $30 \leq \lambda/\Delta\lambda \leq 100$ resolution
- DR3 released 6.6 million quasar candidates!





Bailer-Jones+2022

- Complete, low purity, as many are stars masquerading as QSO :/ but... we can improve it

Gaia: more than astrometry


- Space-quality data means e.g. no seeing, airmass, stable observing conditions...
 - larger volume, cleaner selection function than any existing sample (full sky).
 - Synergies with external data allowed to clean this sample further: Quaia



Quaia: the cosmological QSO sample of Gaia


- unWISE + SDSS DR18 + color and proper motion cuts = better purity, low systematics

Storey-Fisher+
(2023)



The selection function

- A handful of templates + Gaussian processes corrects for scanning pattern, stars, extinction...
- Some further updates ongoing, new Gaia DR also soon... (Williams, Storey-Fisher+ in prep.)



How it compares?

Quaia

SDSS DR16Q





Credits Kate Storey-Fisher

	N	f_{sky}	$\bar{n}$, deg^{-2}	V_{span} , $(h^{-1} \text{ Gpc})^3$	V_{eff} , $(h^{-1} \text{ Gpc})^3$	z_{med}	$f(\delta z < 0.01)$	$f(\delta z < 0.1)$
Quaia	1,209,833	0.71	41.50	138.46	7.01	1.47	0.61	0.83
<i>Gaia</i> Purer	1,261,884	0.71	43.29	138.44	6.43	1.61	0.62	0.70
WISE-PS1	1,106,010	0.54	49.81	105.45	7.21	1.43	0.12	0.76
eBOSS	190,194	0.14	34.03	26.55	1.01	1.49		


What Quiaia can do for cosmology?

- High redshift
- Linear or quasi-linear scales.



What Quiaia can do for cosmology?


- Large sky coverage means capability to measure large angular scales.
- Towards non-degenerate σ_8, Ω_m measurements




Alonso, GF, Storey-Fisher+(2023)

Cosmology from Quaia x Planck CMB lensing

- C_ℓ^{gg}, C_ℓ^{kg} with Planck PR4 CMB lensing in 2 redshift bins.
- Results competitive with current LSS surveys with fewer objects and/or worse redshifts.



Alonso, GF, Storey-Fisher+(2023)





No tension
with Planck!

Cosmology from Quaia x CMB lensing

- Tomographic $C_{\ell}^{gg}, C_{\ell}^{kg}$ with Planck lensing
 - New high- z σ_8 constraints
 - Results competitive with current LSS surveys with fewer objects and/or worse redshifts.
- Tomographic 3x2pt $C_{\ell}^{gg}, C_{\ell}^{kg}, C_{\ell}^{kk}$ with ACT CMB lensing
 - Constraints on $\sigma_8(z = 5.1)$

$$P_{\text{lin}}^{\text{new}}(k, z) = P_{\text{lin}}^{\text{input}}(k, z) A(z)$$


$$= P_{\text{lin}}^{\text{input}}(k, z) \begin{cases} A_0 & 0 \leq z < z_1 \\ A_1 & z_1 \leq z < z_2 \\ A_2 & z_2 \leq z \end{cases}$$



Non-Gaussianities and scale dependent bias


$$\Phi(\mathbf{x})_{\text{NG}} = \Phi(\mathbf{x}) + f_{\text{NL}} (\Phi^2(\mathbf{x}) - \langle \Phi^2 \rangle)$$

$$\Delta b_1 \propto \frac{f_{\text{NL}} b_1 b_\phi}{k^2 T(k)}$$




- Huge volume and high bias tracers is required to enhance detection.
- Quasars: detectable at high z thanks to bright AGNs, $\sim$ linear scales, ideal for f_{NL} studies.
- Galactic extinction, stellar contamination, seeing, survey depth inhomogeneity cause large scale variations / power excess in clustering measurements...

Signature of f_{NL} for projected statistics



Fabbian, Alonso, Storey-Fisher (2025)


Constraints on f_{NL}



- Cross-correlation can reach the largest scales!

- $\sigma(f_{\text{NL}})$: worse than expectations due to residual systematics $\Delta C_\ell^{kg} \propto \sqrt{C_\ell^{kk} C_\ell^{gg}}$



Robustness tests and perspectives




- Sensitivity comparable to spectroscopic galaxy surveys in 3D
- Limited by systematics on auto correlation on large scales but highly robust to systematics.
- Newer data releases from Gaia could improve this results $\sigma(f_{NL}) \sim 9$ (more sources, different source separation, new data...)

An example of Euclid synergies

- Large number of objects detected as dropouts in Euclid bands...
- AGN/QSO cosmological sample can be constructed, work in progress...
 - Build from Quiaia work to construct / improve samples.
- Robust constraints can be achieved for non-Gaussianities $\sigma(f_{NL}) \sim 2$



Towards Euclid DR3




expected in Oct 2031
covering 15 k deg²

Overlap with **other LSS survey**:

- ▶ with with DESI over $\sim 9000 \text{ deg}^2$
- ▶ with both BOSS and DESI over $\sim 6000 \text{ deg}^2$



- ▶ **SO** has the largest overlap with Euclid (65% overlap)
- ▶ SPT3G-WIDE is the second best (45%)
- ▶ ACT: similar SO overlap but with lower sensitivity



Naidoo et al, 2023 (Euclid collab paper)

Forecast for 3x2pt analysis

- Assumes the pessimistic Euclid survey in ~ 5 tomographic redshift bins for GC-phometric and WL, $z_{\text{max}}=1$





Euclid collab. (Blanchard+19)

Euclid collab. w/ CMBX SWG (Ilic+21)

Forecast for 6x2pt analysis

- Pessimistic Euclid survey in ~ 5 tomographic redshift bins and WL, $z_{\text{max}}=1$
- Assumes SO-like baseline sensitivity for CMB data (conservative): 4-10x improvements in dark energy / modified gravity parameters.




Euclid collab. (Blanchard+19)

Euclid collab. w/ CMBX SWG (Ilic+21)

Towards DRI: the QI data release

Quick Data
Release this
March



expected
internally
VERY SOON




- Preliminary release with 1st quarter of data (QI) included $\sim 60 \text{ deg}^2$ with “pre-launch” pipeline.

- Ideal for cross-correlation!


Fabbian, Legrand, Pagano, Lembo, Piccirilli,
Kou+CMBX SWG (in prep.)

- DRI analysis: Planck for the north, both ACT and SPT in the south patch
- QI analysis: Planck for the north, ACT for the EDFF and EDFs patches


Towards DRI: the QI data release



- Magnitude cut in $\text{VIS} < 24.5$
- Properties consistent across fields
- Agreement with Flagship simulation predictions “uncalibrated”!!!!




Towards DRI: the QI data release



- Two different sample selection based on purity of galaxy classification or star classification rejection respectively.
- The two cuts deliver samples with: ~ 11 gal/arcmin, ~ 21 gal/arcmin, with different redshift limits ($\sim 1.2, \sim 2.5$)
- Systematics deprojection needed due to lack of deep external data and no full visibility processing (minimal effects)


Towards DRI: the QI data release




- Two different sample selection based on purity of galaxy classification or star classification rejection respectively.
- The two cuts deliver samples with: ~ 11 gal/arcmin, ~ 21 gal/arcmin, with different redshift limits (~ 1.2 , ~ 2.5)
- Systematics deprojection needed due to lack of deep external data and no full visibility processing (minimal effects)

Preliminary results: single bin

w vs /wo systematics deproj



Different sample selections





- **WARNING: theory lines are not fits!**
 - Planck 2018 + estimated $N(z)$ + fiducial Flagship calibrated bias model.
 - TLDR: great PTEs (w/ RSD, magnification bias...)
- Cross-correlation extremely stable and robust to any systematics deprojection!

Preliminary results: 3 tomographic bins!

w vs /wo systematics deproj

Different sample selections



- WARNING: theory lines are not fits!


Fabbian+(in prep.)

- Planck 2018 + estimated $N(z)$ + fiducial Flagship calibrated PT bias model.
- Great PTEs!
- $k_{\max} \sim 0.5 \text{ h/Mpc}$ (but good to smaller scales)

Some preliminary findings

- CMB lensing cross-correlation is ready for fine bin tomography.
 - Well modeled across different fields and fine redshift binning
- Auto-correlation: it depends....
 - We are seeing inhomogeneous redshift quality (as expected)
 - Limited tomographic capabilities.
 - Star galaxy separation...
- We can recover constraining power with cross-correlation.


It's a long way to go



- Dense sample allows to fit redshift errors from data

$$p(z) \rightarrow p(z + (z-z_0)w + \Delta z)$$

- We retrieve feature of photo-z estimation.



Tucci+(2025, Euclid coll.)

Conclusions

- **Exciting prospects from new CMB and space-based galaxy surveys data...**
 - Quia allowed for non-degenerate σ_8 , Ω_m , S_8 consistent with Planck (and independent H_0 ...)
 - Robust and competitive f_{NL} from cross-correlation with CMB lensing.
 - More is expected from SO and Euclid (e.g. delensing B-modes from inflation constraints).
- **Euclid is working beautifully!**
 - QI allowed us to benchmark the data for cosmology and all looks extremely promising so far.
 - But not without challenges: i.e. previous results with a pinch of salt...
 - We are catching up with the shear analysis...
- **Results coming soon, stay tuned!**