



Dipole Excitations in Nuclei: Bridging Femtometer Physics with Cosmic-Scale Questions


PANDORA Project

Photo-Absorption of Nuclei and Decay Observation for Reactions in Astrophysics



2

Kamila Sieja



THEORY TEAM: ACTIVITIES

- **High-energy (particle) physics**

1. Effective field theories, Lie algebras

2. Physics beyond the Standard Model: from formal aspects to phenomenology




- **Low-energy (nuclear) physics**

1. Few-body systems:

- Weakly-bound systems, resonances, physics of antimatter

2. Many-body systems:


- Exotic nuclei, weak processes, applications in astrophysics
- Exotic symmetries, superheavy systems



HIGH-ENERGY (PARTICLE) PHYSICS

PHYSICS SUBJECTS:

- Supersymmetry, supergravity
- Lie groups
- Quantum field theory : effective field theories for particle physics, scattering amplitudes, symmetries
- Particle physics beyond-the-Standard-Model : strong CP problem, axions, flavor
- Early-universe cosmology : dark matter, cosmic strings



PhD thesis:

S. Beaudoin (PhD ongoing)

« Axions, cordes cosmiques et monopoles magnétiques »

V. Saulquin (PhD ongoing)


« Nouveaux modèles de physique des particules inspirés par le problème CP fort et les symétries de saveur »



Q. Bonnefoy



**M. Rausch
de Trautenberg**



J. Polonyi (emeritus)



S. Beaudoin




V. Saulquin

FEW-BODY SYSTEMS

PHYSICS SUBJECTS:

- 3-5 particle systems atomic, molecular, nuclear
- Weakly-bound systems, resonances
- Physics of antimatter, interaction between matter and antimatter
- Experiments GBAR and PUMA @CERN



ANR
FBCUBES



M. Dufour



R. Lazauskas

M. Schäfer
(post-doc)

MANY-BODY SYSTEMS (1)

PHYSICS SUBJECTS:

- K-isomers, shape isomers, Gallagher-Moszkowski splitting in odd-odd nuclei, fission
- High-rank symmetries
- Multidisciplinary applications: hyperfine structure



J. Bartel
(emeritus)

J. Dudek
(emeritus)



H. Molique

MANY-BODY SYSTEMS (2): valence-space approaches

PHYSICS SUBJECTS:

- Exotic nuclei: spectroscopy, shell evolution, shape changes
- Dipole excitations
- Weak-decays (beta and duple-beta)
- Reactions of interest for astrophysics (r-process, UHECR)

TOOLS:





- Effective interactions
- Codes CI-SM and DNO-SM (PGCM):
- ANTOINE, NATHAN, CARINA



D. D. Dao



F. Nowacki



K. Sieja



O. Le Noan



**Dipole Excitations in Atomic Nuclei:
Theoretical Models and Applications**
(PhD O. Le Noan+ QMAT project 2024-2026)

The Collective Response of the Nucleus: Giant Resonances



Isoscalar (In phase)
 $\Delta T = 0$

Isvector (Out of phase)
 $\Delta T = 1$



Monopole
 $\Delta L = 0$
(GMR)



Dipole
 $\Delta L = 1$
(GDR)





Quadrupole
 $\Delta L = 2$
(GQR)



Nuclear Dipole Excitations: from Academic to Industrial Applications

- Fundamental mode of nuclear excitations
- Dominated by E1 mode (with a smaller contribution of the M1)
- Basic input to nuclear reaction codes



generated by AI

$$\sigma_{(n,\gamma)}^{\mu\nu}(E_i, n) = \frac{\pi \hbar^2}{2M_{i,n} E_{i,n}} \frac{1}{(2J_i^\mu + 1)(2J_n + 1)} \sum_{J,\pi} (2J + 1) \frac{T_n^\mu T_\gamma^\nu}{T_n^\mu + T_\gamma^\nu}$$

for $E_n \sim \text{keV}$ $T_n^\mu \gg T_\gamma^\nu \rightarrow \sigma^{\mu\nu} \sim T_\gamma^\nu$

For a given multipolarity

$$T_{XL}(E, J, \pi, E^\nu, J^\nu, \pi^\nu) = 2\pi E_\gamma^{2L+1} f_{XL}(E, E_\gamma)$$

f_{XL} = Photon Strength Function (PSF)


TALYS code: Modeling of Nuclear Reactions

A. Koning, S. Hilaire, S. Gorioli et al. *Eur. Phys. J A* 59 (2023) 131


Nuclear Physics Meets the Sources of the Ultra-High Energy Cosmic Rays

After >50 years of studying UHECR open questions remain :


- Where do they come from?
- How they are accelerated?




PANDORA Project
Photo-Absorption of Nuclei and Decay Observation for Reactions in Astrophysics
A. Tamii et al, Eur. Phys. J.A59 (2023) 208




PANDORA
Photo-Absorption of Nuclei and Decay Observation for Reactions in Astrophysics



Boncioli, D., Fedynitch, A. & Winter, W., Sci Rep 7, 4882 (2017).



Many-body methods to compute dipole response




$$H_{\text{MF}} = \sum_i [T_i + U(r_i)] \longleftrightarrow H = \sum_i T_i + \sum_{i < j} V_{ij} + \dots$$

Quasiparticle Random Phase Approximation (QRPA)

$$i\partial_t \mathcal{R}(t) = [\mathcal{H}(t) + \mathcal{F}(t), \mathcal{R}(t)],$$

$$\begin{aligned} (E_\mu + E_\nu - \omega) X_{\mu\nu}(\omega) + \delta H^{20}(\omega) &= -F^{20}(\omega) \\ (E_\mu + E_\nu + \omega) Y_{\mu\nu}(\omega) + \delta H^{02}(\omega) &= -F^{02}(\omega) \end{aligned}$$



Configuration Interaction Shell Model (CI-SM)

$$|\Psi_0\rangle = C_0|\Phi_0\rangle + \sum_{i\alpha} C_{i\alpha}|\Phi_{i\alpha}\rangle + C_{ij\alpha\beta}|\Phi_{ij\alpha\beta}\rangle + \dots$$

$$E = \langle \Psi_0 | \hat{H} | \Psi_0 \rangle$$

Projected Generator Coordinate Method (PGCM)

$$|\Psi_{GCM}^\sigma\rangle = \int f_\sigma(\vec{q}) |\Phi(\vec{q})\rangle d\vec{q}$$



$$\begin{aligned} E_\sigma[|\Psi_{GCM}^\sigma\rangle] &= \frac{\langle \Psi_{GCM}^\sigma | \hat{H} | \Psi_{GCM}^\sigma \rangle}{\langle \Psi_{GCM}^\sigma | \Psi_{GCM}^\sigma \rangle} = \\ &= \frac{\int \langle \Phi(\vec{q}) | \hat{H} | \Phi(\vec{q}') \rangle f_\sigma^*(\vec{q}) f_\sigma(\vec{q}') d\vec{q} d\vec{q}'}{\int \langle \Phi(\vec{q}) | \Phi(\vec{q}') \rangle f_\sigma^*(\vec{q}) f_\sigma(\vec{q}') d\vec{q} d\vec{q}'} \end{aligned}$$

Benchmark of many-body methods:

1. M. Frosini, W. Ryssens and K. Sieja, *Phys. Rev. C* **110**, 014307 (2024)
2. S. Bofos, J. Martinez, B. Bally, T. Duquet, M. Frosini, T. Rodriguez and K. Sieja, *Phys. Rev. C* **112**, 064312 (2025)



Impact of Photonuclear Reaction Models on Propagation Ultra-High Energy Cosmic Rays

R. Fearick, **O. Le Noan**, H. Matsubara, P. von Neumann-Cosel, **K. Sieja** and A. Tamii, in preparation.



O. Le Noan and K. Sieja, PRC111 (2025) 064308.

CI-SM PSF
database
for TALYS




$$\begin{aligned} \frac{dN_{Z,A}}{dt} = & N_{Z+1,A} \lambda_{\beta}^{Z+1,A} + N_{Z-1,A} \lambda_{\beta}^{Z-1,A} \\ & + N_{Z,A+1} \lambda_{\gamma,n}^{Z,A+1} + N_{Z+1,A+1} \lambda_{\gamma,p}^{Z+1,A+1} \\ & + N_{Z+2,A+4} \lambda_{\gamma,\alpha}^{Z+2,A+4} + N_{Z,A+2} \lambda_{\gamma,2n}^{Z,A+2} \\ & + N_{Z+2,A+2} \lambda_{\gamma,2p}^{Z+2,A+2} + N_{Z+4,A+8} \lambda_{\gamma,2\alpha}^{Z+4,A+8} \\ & + N_{Z+1,A+2} \lambda_{\gamma,np}^{Z+1,A+2} + N_{Z+2,A+5} \lambda_{\gamma,n\alpha}^{Z+2,A+5} \\ & + N_{Z+3,A+5} \lambda_{\gamma,p\alpha}^{Z+3,A+5} \\ & - N_{Z,A} \left[\lambda_{\beta}^{Z,A} + \sum_x \lambda_{\gamma,x}^{Z,A} \right], \end{aligned} \quad (9)$$






O. Le Noan, S. Goriely, E. Khan and **K. Sieja**, to be submitted.

New Insights into the Nature of the Pygmy Dipole Resonance

O. Le Noan and K. Sieja, PRC111 (2025) 064308.




$$\delta\rho_{\nu 0}(\vec{r}) = \langle \nu | \sum_k \delta(\vec{r} - \vec{r}_k) | 0 \rangle,$$



Motivations to study PDR:

- PDR enhances probability of (n,γ) CS
=> influence on astrophysical r-process
- PDR relates to neutron-skin thickness
=> relation to neutron stars



P. von Neumann Cosel et al. Physics (2023)

Open questions:

- Evolution of PDR with angular momenta and deformation
- Isospin mixing of PDR (IS vs IV)
- Is PDR a toroidal mode?

Thank You !

Thanks to:

IPHC: O. Le Noan

CEA Saclay: T. Duguet

CEA Cadarache: S. Bofos, M. Frosini, A. Pastore

CEA DAM: S. Hilaire

IJCLab: E. Khan

TU Darmstadt: B. Bally

Universidad de Sevilla: T. R. Rodriguez

Universidad Autonoma de Madrid: J. Martinez-Larraz

Université Libre de Bruxelles: S. Goriely, W. Ryssens

University of Zagreb: N. Paar

