

UNIONS

cosmological results with 2D weak lensing



Martin Kilbinger

on behalf of the UNIONS CosmoStat Weak Lensing Team

cea


 CosmoStat

Deep CosmoStat days, Feb 12, 2026

Outline of the talk

- The UNIONS survey
- Weak lensing with UNIONS
- First shear cosmology with UNIONS:
 - 2-point correlation functions
 - Image simulations & calibration
 - Systematics: PSF leakage & B-modes
 - Photometric redshift estimation & blinding
 - Inference & covariance
 - Cosmological constraints

UNIONS: A combination of 3 Hawai'ian telescopes




UNIONS: Ultra-violet Near-Infrared Optical Northern Survey


Goal: 6,200 deg² in 5 bands;
 u, r (CFIS: Canada-France Imaging Survey)
 i, z (Pan-STARRS)
 g, z (HSC).

P.I.: Jean-Charles Cuillandre (CEA Paris-Saclay) & Alain McConnachie (Victoria/Canada)

- Optical bands for Euclid for photometric redshifts
- Weak lensing
- Milky Way dynamics
- Large-scale structure
- Galaxy evolution





UNIONS multi-band data sky coverage




UNIONS image quality

Best wide-field imager on CFHT ever.


Improvements (2011 - 2014)



Dome venting




40 CCDs + Fast readout



New "square" filters

UNIONS in the broader survey landscape

Final UNIONS will be a
“Stage 3.5” survey—
roughly equivalent in
depth & area to LSST Y1.



ShapePipe & the UNIONS shape catalog

<https://github.com/cosmostat/shapepipe>

Farrens et al., 2022, [A&A, 664, 141](#)

ShapePipe

ShapePipe is a galaxy shape measurement pipeline developed within the CosmoStat lab at CEA Paris-Saclay.

See the [documentation](#) for details on how to install and run ShapePipe.

https://github.com/cosmostat/sp_validation/ for post-processing



ShapePipe philosophy

Goals



- Modular
- Easy
- Fast (enough)
- Robust

Code installation



- Conda
- Docker [allows for cloud computing]
- CD/CI

Three components

Pipeline



- Arguments & config
- I/O
- Job handling (MPI, SMP)
- Errors & logging

Modules



- WL data processing
- Book-keeping
- Post-processing

Utilities



- Scripts
- Tools
- Survey-specific content

Can process 10k+ images, create catalogues with ~ 500 million objects, 150 Tb.



ShapePipe WL image processing

github/shapepipe
Farrens et al., 2022, [A&A, 664, 141](#)

Input images are pre-processed (calibrated for astrometry and photometry)

Main processing

- Mask
- Detect objects
- Star candidates on single exposures
- Galaxy candidates on stacks
- Select stars
- Create PSF model (PSFEx, Bertin et al. 2011; MCCD, Liaudat et al. 2021)
- Interpolate PSF model to galaxy positions
- Validate PSF model
- Measure galaxy shapes including calibration information [ngmix + metacalibration]



Post-processing

- Galaxy selection
- Apply calibration
- Systematic checks and validation





Image simulations & shear calibration

Simulated and real CFHT exposure.

Simulations:

- Realistic galaxy morphologies (Sérsic profiles matched to COSMOS)
- Matching survey properties: camera geometry, dither pattern, noise
- Matching observed galaxy distribution: ellipticity, size, SNR, PSF, ...
- 80 deg² for pipeline validation and calibration





Image simulations & shear calibration

Shape/shear measurement: metacalibration = **self-calibration** using data.

Residual shear bias mainly from blends, expected 1-2%. Calibrated with **image simulations**. Final value TBD.

UNIONS vs. DES:

- Seeing 0.69" vs. 0.95", less sensitive to blends:
 - **2X smaller** amplitude
 - At **2X smaller distance** between galaxies





Image simulations & shear calibration


Additional residual shear bias m .

Simulations quantify m due to


- **Software bug**
(pixel vs. arcsec units)
- **Galaxy selection**
(removing objects as
blended is correlated with
ellipticity and shear)



2-point correlation functions



2-point shear correlation function
↔ variance of convergence σ_{κ}^2
↔ κ power spectrum
= projection(δ power spectrum)



Linear combinations

$$\xi_+(\vartheta) = \langle \gamma_t \gamma_t \rangle(\vartheta) + \langle \gamma_x \gamma_x \rangle(\vartheta)$$

$$\xi_-(\vartheta) = \langle \gamma_t \gamma_t \rangle(\vartheta) - \langle \gamma_x \gamma_x \rangle(\vartheta)$$

First shear cosmology with UNIONS

- Traditional cosmic shear analysis à la KiDS, DES, HSC.
- 2D: single redshift bin since more time is required to estimate & validate tomographic redshifts.
- Blinded analysis; blinding performed on the redshift distribution.
- 2-point correlation function (2PCF) used as data vector for cosmological inference.

The 2D cosmic shear team

Core members in alphabetical order (many others have contributed as well):



Cail Daley
(B-mode systematics)



Lisa Goh
(inference / covariance)



Sacha Guerrini
(inference / PSF systematics)



Calum Murray
(validation, IA)



Fabian Hervas-Peters
(image simulations)



Martin Kilbinger
(shape catalog)





Anna Wittje
(redshift estimation)



Antonin Corinaldi
(intrinsic alignment)

PSF Systematics

- Two modes:
 - PSF leakage: PSF imprint in galaxy shapes.
UNIONS: 1-2% leakage; corrected on galaxy basis.
 - PSF modelling errors: residuals can mimic shear correlations.
- Use star-galaxy correlations to quantify [Guerrini et al. 2025]




B-mode systematics

Spin-2 shear fields can be decomposed into **E-modes** containing the vast majority of lensing information and **B-modes** which are a probe of systematics at UNIONS noise levels.

In the presence of masking, some **ambiguous** modes cannot be cleanly attributed to E or B, although recent **purified** estimators can separate out these modes.


We use three B-mode approaches: pure correlation functions, COSEBIS, and bandpowers.




BICEP2 Collaboration

Pure B-mode correlation functions

B-modes on small scales ($\sim 4'$ in ξ_+ and $\sim 30'$ in ξ_-); large scales are ok.



Pure B-mode correlation functions




Testing masking and galaxy selections.

B-mode scale cuts

Pure-B correlation functions and COSEBIS tell a similar story:


- ξ_+ PTE is acceptable above $\theta \sim 3-4$
- B_\square (dominated by ξ_+) PTE is acceptable above $\theta \sim 3-4$
- ξ_- (probing $\lesssim 10$ x smaller scales) PTE is acceptable above $\theta \sim 20-30$.



B-mode COSEBIs

- Complete Orthogonal Sets of E/B-mode Integrals [Schneider, Eifler & Krause 2010].
- Transform shear correlation function into discrete, pure E- and B-modes.
- First few modes have most SNR

ℓ





B-mode band powers

Harmonic-space band power spectra.

- New direct catalogue-based estimation [Wolz et al. 2025], implemented in NaMaster
- Accounts for noise and mask mixing matrix, estimates covariance
- B-mode passes null test with $l_{\min} = 300$


Conclusion: consistent scale cuts in configuration space (12' - 83') and harmonic space (300 - 1200).



Redshift Distributions

Redshift distribution estimated using self-organizing maps (SOMs).


Three blinded redshift distributions produced—allows us to run the full inference pipeline on the data without risking confirmation bias.




Covariance & Inference

Covariance estimated with CosmoCov and validated against data-driven jackknife.

Other parameters marginalized over in inference: intrinsic alignment [Hervas Peters 2024 direct measurement], multiplicative bias, PSF systematics, $n(z)$ bias.




Current Best-fit Theory & Systematics



Cutting two $5' < \theta < 10'$ data points in ξ_+ improves χ^2 by 13..


Best fit in harmonic space



Blinded Cosmological Contours

Constraints on S_8 : ± 0.06


- ~ 0.015 comes from conservative $\theta > 10'$ scale cut
- Non-tomographic analysis significantly reduces constraining power as well.
- Shift between configuration and harmonic space at 1.5σ



Blinded constraints


Constraints on S_8 : ± 0.06

- ~2-3x larger than best constraints from DES, HSC, & KiDS
- Somewhat larger; consistent with Planck and KiDS-Legacy
- Insensitive against further scale cuts, non-linear model, covariance.



Configuration vs. harmonic space

- Shift quantified with log-normal mocks.
- Skewed distribution?
- Similarly observed in KiDS.




Summary & Next Steps

- UNIONS is a unique dataset for weak lensing:
 - excellent image quality (Mauna Kea)
 - homogeneous survey depth (adaptive observing strategy).
 - Large area ($> 6,000 \text{ deg}^2$)
 - Very good photo-z's (u -band @ CFHT)
- Competitive in the Euclid and Rubin era, in particular for cross-correlations with SDSS and DESI: lensing by galaxies, groups, clusters, voids; 3x2pt.
- First UNIONS cosmic shear results are imminent!
 - Analysis is in its final stages, tracking down potential scale-dependent systematics before unblinding.
 - Error bar on S_8 forecasted to be ~ 0.06 , and may improve if we can reduce systematics on small scales.
- Up next: simulation-based inference, tomographic analysis, and much more!

Backup slides

Conf vs. harmonic space



UNIONS weak-lensing publications

Published/finished

Void lensing
UNIONS overview paper
Galaxy-galaxy lensing of mergers
Cluster lensing of mergers
Cluster lensing
Intrinsic galaxy alignment
PSF systematics and diagnostics
PSF diagnostics for galaxy-galaxy lensing
Black-hole-mass - halo-mass relation
Peak counts
UNIONS first weak-lensing analysis
Group & cluster masses
Dark-matter halo shapes
Multi-CCD PSF model

Martin et al. 2026, [MNRAS in press](#)
Gwyn et al. 2025, [ApJ, 170, 6, 324](#)
Cheng et al. 2025, [ApJ, 992, 2, 171](#)
Ahad et al., 2026, [submitted](#)
Mpetha et al. 2025, [MNRAS, 543, 2, 1393](#)
Hervas Peters et al. 2025, [A&A, 699, A201](#)
Guerrini et al. 2024, [A&A in press](#)
Zhang et al. 2024, [A&A 691, A75](#)
Li et al. 2024, [ApJ, 969, 2, L25](#)
Ayçoberry et al., 2023, [A&A, 671, A17](#)
Guinot et al., 2022, A&A, 666, A1
Spitzer et al., 2022, submitted to MNRAS
Robison et al., 2022, [arXiv:2209.09088](#)
Lioudat et al., 2021, [A&A, 646, A27](#)

In progress

2D cosmic shear catalogues
2D cosmic shear validation & B-modes
2D cosmic shear in configuration space
2D cosmic shear in harmonic space
2D cosmic shear calibration & image simulations
Simulation-based inference
3x2pt cosmology
3D intrinsic alignment
Intrinsic alignment at formation time
Intrinsic alignment multipole measurements

Hervas Peters et al. in prep.
Daley et al in prep.
Goh et al. in prep.
Guerrini et al. in prep.
Hervas Peters et al. in prep.
Guerrini, Maupas in prep.
Hervas Peters et al. in prep.
Corinaldi et al in prep.
Murray et al. in prep.
Paviot et al. in prep.

PSF Systematics

Three quantities can be used to form six ρ -statistics and three τ -statistics:

- Ellipticity
(model & galaxy)
- Ellipticity errors
(model evaluated at star locations)
- Size errors
(model evaluated at star locations)

$$\delta \mathbf{e}_{\text{model}}^{\text{sys}} = \alpha \underbrace{\mathbf{e}_{\text{model}}}_{\text{Leakage}} + \beta \underbrace{(\mathbf{e}_* - \mathbf{e}_{\text{model}})}_{\text{Ellipticity error}} + \eta \underbrace{\left(\mathbf{e}_* \frac{T_* - T_{\text{model}}}{T_*} \right)}_{\text{Size error}}$$


$$\begin{pmatrix} \tau_{0,1} \\ \tau_{2,1} \\ \tau_{5,1} \\ \vdots \\ \tau_{0,n} \\ \tau_{2,n} \\ \tau_{5,n} \end{pmatrix} = \begin{pmatrix} \rho_{0,1} & \rho_{2,1} & \rho_{5,1} \\ \rho_{2,1} & \rho_{1,1} & \rho_{4,1} \\ \rho_{5,1} & \rho_{4,1} & \rho_{3,1} \\ \ddots & & \\ \rho_{0,n} & \rho_{2,n} & \rho_{5,n} \\ \rho_{2,n} & \rho_{1,n} & \rho_{4,n} \\ \rho_{5,n} & \rho_{4,n} & \rho_{3,n} \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \\ \eta \end{pmatrix},$$

Solve system of linear equations to get
leakage contribution to 2PCF:


$$\begin{aligned} \xi_{\text{PSF,sys}}(\vartheta) &= \alpha^2 \rho_0(\vartheta) + \beta^2 \rho_1(\vartheta) + \eta^2 \rho_3(\vartheta) \\ &\quad + 2\alpha\beta \rho_2(\vartheta) + 2\alpha\eta \rho_5(\vartheta) + 2\beta\eta \rho_4(\vartheta) \end{aligned}$$

Photo-z's


WISHES area





PanSTARRS area



Tomography




Simulation-based inference




Mathis Maupas, Sacha Guerrini

Simulation-based inference




UNIONS-like simulations



Mathis Maupas, Sacha Guerrini

UNIONS extension, $\delta < 30^\circ$



Ground-based coverage of the 16 Kdeg² Euclid Wide Survey Region of Interest [origin/bands/overlap/calendar] [Mollweide Celestial]

DES (Blanco), griz : 4.8 Kdeg² overlap since 2019

LSST Wide-Fast-Deep (Rubin), ugriz : 10.2 Kdeg² overlap by 2026

UNIONS (CFHT/Pan-STARRS/Subaru), ugriz : 4.5 Kdeg² by 2025

UNIONS extended, ugriz : 1.4 Kdeg² by 2027

Euclid Region of Interest : 16.2 Kdeg²

Euclid Deep Fields [53 deg²]



Background image: Euclid Consortium / Planck Collaboration / A. Heilinger

27 accepted peer reviewed publications so far:

< 2024

Resolved Stellar Populations Galaxy evolution Weak lensing

27. Robison, B., et al., 2023, *in press*, "The shape of dark matter haloes: results from weak lensing in the Ultraviolet-Near Infrared Optical Northern Survey (UNIONS)"
26. Lim, S., et al., 2023, *MNRAS*, *in press*, "Constraints on galaxy formation from the cosmic-infrared-background / optical-imaging cross-correlation using Herschel and UNIONS"
25. Smith, S., et al., 2023, *ApJ*, *in press*, "Discovery of a new Local Group galaxy candidate in UNIONS: Bo'otes V"
24. Chu, A., et al., 2023, *A&A*, *in press*, A UNIONS view of the brightest central galaxies of candidate fossil groups
23. Bickley, R., et al., 2023, *MNRAS*, 519, 6149, "AGN in post mergers from the Ultraviolet Near Infrared Optical Northern Survey"
22. Ayc,coberny, E., et al., 2023, *A&A*, 671, 17, "UNIONS : impact of systematic errors on weak-lensing peak counts"
21. Savary, E., et al., 2022, *A&A*, 666, 1 "A search for galaxy-scale strong gravitational lenses in UNIONS"
20. Chan, J. H. H., et al. 2022, *A&A*, 659, 140 "Discovery of Strongly Lensed Quasars in UNIONS"
19. Wilkinson, S., et al., 2022, *MNRAS*, 516, 4354, "The merger fraction of post-starburst galaxies in UNIONS"
18. Ellison, S., et al., *MNRAS*, 517, L92, "Galaxy mergers can rapidly shut down star formation"
17. Bickley, R., et al., 2022, *MNRAS*, 514, 3294, "Star formation characteristics of CNN-identified post-mergers in the Ultraviolet Near Infrared Optical Northern Survey (UNIONS)"
16. Farrans, S., et al., 2022, *A&A*, 664, A141, "A modular weak lensing processing and analysis pipeline"
15. Guinot, A., et al., 2022, *A&A*, 666, 162, "ShapePipe: a new shape measurement pipeline and weak-lensing application to UNIONS/CFIS data"
14. Sola, E., et al., 2022, *A&A*, 662, 124, "Characterization of LSB structures in annotated deep images"
13. Roberts, I., et al., 2022, *MNRAS*, 509, 1342, "Ram Pressure Candidates in UNIONS"
12. Jensen, J., et al., 2021, *MNRAS*, 507, 1923, "Uncovering fossils of the distant Galaxy with UNIONS: NGC 5466 and its stellar stream"
11. Bickley, R., et al., 2021, *MNRAS*, 504, 372, "Convolutional neural network identification of galaxy post-mergers in UNIONS using IllustrisTNG"
10. Fantin, N., et al., 2021, *ApJ*, 913, 30, "The Mass and Age Distribution of Halo White Dwarf Candidates in the Canada-France Imaging Survey"
9. Lioudat, T., et al., 2021, *A&A*, A27, "Multi-CCD modelling of the point spread function"
8. Thomas, G., et al., 2020, *ApJ*, 902, 89, "The Hidden Past of M92: Detection and Characterization of a Newly Formed 17 \circ Long Stellar Stream Using the Canada-France Imaging Survey"
7. Fantin N., et al., 2019, *ApJ*, 877, 148, "The Canada France Imaging Survey: Reconstructing the Milky Way from its white dwarf population"
6. Thomas, G., et al., 2019, *ApJ*, 866, 10, "Dwarfs or giants? Stellar metallicities and distances from ugrizG multi-band photometry"
5. Ellison, S., et al., 2019, *MNRAS*, 487, 2491, "A definitive merger-AGN connection at z=0 with CFIS: mergers have an excess of AGN and AGN hosts are more frequently disturbed"
4. Thomas, G. et al. 2019, *MNRAS*, 483, 3, "A-type stars in the Canada-France Imaging Survey - II. Tracing the height of the disc at large distances with Blue Stragglers"
3. Thomas, G. et al., 2018, *MNRAS*, 481, 4, "A-type stars in the Canada-France Imaging Survey I. The stellar halo of the Milky Way traced to large radius by blue horizontal branch stars"
2. Ibata, R., et al., 2017, *ApJ*, 848, 2, 129, "Chemical Mapping of the Milky Way with The Canada-France Imaging Survey: A Non-parametric Metallicity-Distance Decomposition of the Galaxy"
1. Ibata, R., et al., 2017, *ApJ*, 848, 2, 128, "The Canada-France Imaging Survey: First Results from the u-Band Component"

Visual inspection!

