


Dark matter & axion detection

Maurizio Giannotti,
University of Zaragoza, CAPA
& Barry University

Contact: mgiannotti@unizar.es



GraSPA 2026
23 July, 2026
ANNECY, FRANCE

ID Card

Name: Maurizio Giannotti,
Citizenship: Italian, American
Profession: Astroparticle physics

Contact → mgiannotti@unizar.es



Interest:

Astrophysical methods to
study Fundamental Physics

Fundamental interactions

New particles




Particle Physics beyond the
Standard Model (SM)

Be ready
to Travel






Universidad de Zaragoza (Spain)

The Circle of Life



The Circle of Life



Summary

- *Dark Matter*
- *Axions as a fundamental QCD ingredient*
- *The search for axions*


Disclaimer: *Far from complete. Please contact me if you want to get more info or want to know about groups involved in some of this research*

Part 1:

Dark Matter

The Dark Matter problem

What we understand about the universe



The Dark Matter problem



Most of the matter is dark

Ordinary matter

~ 18 %

Dark matter

~ 82 %





Dark Matter?

- **Lord Kelvin** discussed in the appendix of a book based on a series of lectures given in 1884 the hypothesis of “dark bodies”. In discussing Kelvin’s work, in 1906 **Henri Poincaré** used the French term [matière obscure] ("dark matter")
- Dutch astronomer **Jacobus Kapteyn** suggest the existence of dark matter using stellar velocities in 1922
- **Oort** found that the mass in the galactic plane must be greater than what was observed, but this measurement was later determined to be incorrect
- In 1933, Swiss astrophysicist **Fritz Zwicky** applied the virial theorem to the Coma Cluster and obtained evidence of unseen mass he called dunkle Materie ('dark matter'). Zwicky estimated its mass based on the motions of galaxies near its edge and compared that to an estimate based on its brightness and number of galaxies.
- One of the observations that served as evidence for the existence of galactic halos of dark matter were those of **Vera Rubin** and **Kent Ford** about the shape of galaxy rotation curves (1970s).

Rotation curves of spiral galaxies

Measuring the Doppler shift of atomic lines in spiral galaxies → circular velocity of stars as a function of their distance from the galactic center

rotation curve.



$$m \frac{v_{\text{circ}}^2(r)}{r} = \frac{Gm\mathcal{M}(r)}{r^2}$$

$$\Rightarrow v_{\text{circ}}(r) = \sqrt{\frac{G\mathcal{M}(r)}{r}}$$


Yu: 2024 TASI Lectures: A Dark Matter Primer

Rotation curves of spiral galaxies

Measuring the Doppler shift of atomic lines in spiral galaxies → circular velocity of stars as a function of their distance from the galactic center

Rubin and Ford in 1970: first precise measurement of the rotation curve of the Andromeda galaxy (M31)

They determined the curve to be rather flat out to ~22 kpc



[Yu: 2024 TASI Lectures: A Dark Matter Primer](#)

X-rays probe of DM in Galaxy Clusters

Hydrostatic equilibrium for a gas in a galaxy cluster

$$\frac{1}{\rho_{\text{gas}}(r)} \frac{dP_{\text{gas}}}{dr} = - \frac{d\phi}{dr} = - \frac{GM(r)}{r^2}$$

Ideal gas:

$$P_{\text{gas}} = \frac{\rho_{\text{gas}} k T_{\text{gas}}}{\mu m_p},$$

($\mu \approx 0.6$ = mean molecular weight of an admixture of $\sim 75\%$ H and $\sim 25\%$ He)

Density and temperature can be measured from the intensity and the spectrum of the X-ray emission, thereby allowing to reconstruct the total cluster mass as well as its profile using HE equation

The total gravitating mass inferred from HE greatly exceeds the combined mass of the hot intracluster gas, stars, and other observed baryonic components

DM halos around galaxies

The existence of massive DM halos around galaxies can also be proven via galaxy-galaxy lensing




Figure credit [Bodo Schwabe](#)


DM halos around galaxies

The existence of massive DM halos around galaxies can also be proven via galaxy-galaxy lensing

This is the distortion of the images of background galaxies induced by the gravitational lensing effect of the foreground galaxies.

Lensed images of the same galaxy

In the case in figure the lens is the galaxy cluster 0024+1654



Hubble image. Source: science.nasa.gov

The Bullet Cluster

One of the most remarkable empirical evidence of DM


System consists of two colliding galaxy clusters.

Lensing allows to map the total mass (including DM) while X-ray observations can map the hot baryon gas.

Evidently, the majority of the mass is not impacted by the collision, while the baryonic gas is slowed down.


⇒ Strong constraints on DM self-interaction

One of the most detailed maps of the bullet cluster (include JWST observations) in
→ S. Cha et al. (2025) [arXiv:2503.21870](https://arxiv.org/abs/2503.21870)



Mass in green contours and hot gas in color scale. Fig. from the original study: [D. Clowe et al. \(2006\)](#)


Galaxy Formation



Source: [Alexander Knebe lecture](#)

Galaxy distribution

two-degree-Field Galaxy
Redshift Survey



The universe is homogenous at scales above $\sim 100h^{-1}\text{Mpc}$.

On smaller scales, galaxies form clusters which are themselves not distributed uniformly, but their positions are correlated, grouped together in superclusters.


The Inhomogeneous Universe and DM

For structures to collapse, gravity should dominate over pressure.

For baryons this can happen only after decoupling, $z \approx 1100$. For DM it can happen earlier, at $z_{eq} \approx 3300$

After decoupling, baryons behave approximately as pressureless matter. At last they feel only gravity, so they fall into the existing CDM potential wells and start to grow, chasing the blue curve (DM)

Present day inhomogeneity observations require DM.



The Inhomogeneous Universe and DM


DM tends to be roughly spherically distributed in gravitationally bound systems. Normal baryonic matter then collapses in the gravitational well, during which it dissipates energy and cools down, resulting in the spinning disks exhibited by many galaxies

N-body simulations suggest the **Navarro-Frenk-White (NFW)** profile of individual halos

$$\rho_{\text{NFW}}(r) = \rho_s \frac{r_s}{r} \left(1 + \frac{r}{r_s} \right)^{-2}$$

with $d \ln \rho / d \ln r$ changing from -3 to -1 around r_s

DM halos are **not necessarily spherical!** In addition, simulations reveal the presence of numerous **smaller sub-halos orbiting within and around main halos.**



DM density near the sun.

Current studies point to


$$\rho_{\odot} \approx 0.40 \text{ GeV/cm}^3$$

$$\approx 0.0106 M_{\odot}/\text{pc}^3$$

That is considerably less than the estimated total matter density in a averaged over a neighborhood of a few 100 pc around the location of the Sun

Authors	Date	$\rho_{\odot}$ in GeV/cm^3	Notes
Turner	1986	0.28	'uncertainty of about a factor of 2'
Flores	1988	$0.3 \leftrightarrow 0.43$	
Kuijken & Gilmore	1991	$0.42 (\pm 20\%)$	
Widrow et al.	2008	0.304 ± 0.053	Einasto NFW isothermal stellar tracers non-isothermal stellar tracers
Catena & Ullio	2009	0.385 ± 0.027 0.389 ± 0.025	
Weber & de Boer	2009	$0.2 \leftrightarrow 0.4$	
Salucci et al.	2010	$0.43 \pm 0.11 \pm 0.10$	
McMillan	2011	0.40 ± 0.04	
Garbari et al.	2011	$0.11^{+0.34}_{-0.27}$ $1.25^{+0.30}_{-0.34}$	
Iocco, Pato & Bertone	2011	$0.2 \rightarrow 0.56$	
Bovy & Tremaine	2012	0.3 ± 0.1	
Zhang et al.	2012	0.28 ± 0.08	
Piffl et al.	2014	$0.59 (\pm 15\%)$	
Pato, Iocco & Bertone	2015	0.420 ± 0.025	
McKee et al.	2015	0.49 ± 0.13	
McMillan	2016	0.40 ± 0.04	
Sivertsson et al.	2017	$0.46^{+0.07}_{-0.09}$	
Buch et al.	2018	0.608 ± 0.380	result depends on chosen tracers stat and sys errors own determination average of recent results inferred 2σ range from simulations
Eilers et al.	2018	0.30 ± 0.03	
Evans et al.	2018	0.55 ± 0.17	
Karukes et al.	2019	$0.43 \pm 0.02 \pm 0.01$	
Cautun et al.	2020	0.33 ± 0.02	
Sofue	2020	0.359 ± 0.017 0.39 ± 0.09	
Salomon et al.	2020	$0.42 \leftrightarrow 0.53$	
Hattori et al.	2020	0.342 ± 0.007	
Benito, Iocco & Cuoco	2021	$0.4 \leftrightarrow 0.7$	
Lim, Putney, Buckley, Shih	2023	0.446 ± 0.054	
Staudt et al.	2024	0.42 ± 0.06	

Dark Matter Hunting



Alternative to DM


- The evidence for DM is, at the moment, purely gravitational.
- Could just that mean that we don't have an adequate theory of gravity to describe very large scales?
- Modified gravity theories like **MOND** (Modified Newtonian Dynamics) have progressed a lot but still fail to provide a simple model valid on all scales.
- MOND does very well on galactic scales, but it fails on large scales (clusters, CMB)
- the Skordis & Złośnik (2021) "AeST" theory can now fit the CMB and matter power spectrum — but at the price of extra dark-sector-like fields, which reduces the appeal of these models

Alternative to DM

- In MOND there is no separate mass component to adjust. So the **radial field** (traced by the rotation curve) and the **vertical restoring force** perpendicular to the disk are locked to the same baryonic distribution. Measure both independently and modified gravity is over-determined; a nearly spherical dark halo, by contrast, dominates the radial field at large R while leaving the local vertical field near the Sun essentially untouched, so it can satisfy both at once. It is a self-consistency test of gravity carried out within a single galaxy
- Recently, *Zheng-long Wang et al* (12 May 2026), Milky Way Dynamics Favor Dark Matter over Modified Gravity Models, combined Milky Way rotation curves + Gaia data to reconstruct the Galactic gravitational field and show that all modified gravity theories are disfavored.

What is Dark Matter?


See → [M. Cirelli, A. Strumia and J. Zupan \(2024\)](#)



90 orders of magnitude!


What is Dark Matter?

See → [M. Cirelli, A. Strumia and J. Zupan \(2024\)](#)




What is Dark Matter?

See → [M. Cirelli, A. Strumia and J. Zupan \(2024\)](#)



What is Dark Matter?

See → [M. Cirelli, A. Strumia and J. Zupan \(2024\)](#)




Here we focus especially on axions. For a general and updated review
See → [M. Cirelli, A. Strumia and J. Zupan \(2024\)](#) .

For a more WIMP oriented talk, plus other candidates as well,
See → [M. Cirelli, IDM Zaragoza 2026](#)

Macroscopic DM

See → [Jacobs, Starkman, Lynn \(2015\)](#) and → [M. Cirelli, A. Strumia and J. Zupan \(2024\)](#)

Bounds on macroscopic DM



Monitoring of a 1m² area
inside the Skylab space station


cross section on matter
 $\sigma = \pi R^2$

Bounds don't assume
specific production mechanisms. Examples
of cosmological prod. in
→ [Cirelli et al. \(2024\)](#)

Collision of DM with WD
could ignite thermonuclear
runaways fusion and ignite a
type Ia supernova
→ [Peter W. Graham et al \(2018\)](#)

MACHOs

MACHOs = Massive Astrophysical Compact Halo Objects



MACHOs hint
(Unconfirmed by later
survey)


Baryonic MACHOs can
be only a small fraction
of DM (because of BBN
constraints)

Macroscopic DM

Astrophysical objects that consist of baryonic matter but have been created before the BBN are not subject to the BBN constraints since the material that they are made of gets subtracted from the baryonic budget very early on.

Primordial Black Holes

Another example of macroscopic DM which avoids the BBN are PBH



PBHs can evade the BBN bound on baryonic matter because they form before BBN from the collapse of overdense regions of the radiation-dominated Universe. Their mass is therefore not associated with the cosmic baryon abundance, and after formation they behave as cold, pressureless matter.

The formation of PBH is not trivial. We should assure that there is gravitational collapse during radiation dominated era. So, the over density must be efficiently boosted to large values.

Key check: → The PBH should not have evaporated by now.

Stefano Profumo (16 June 2026), [Are Primordial Black Holes a Natural Dark Matter Candidate?](#).



dismisses that the PBH dark matter requires fine-tuning.

Primordial Black Holes


Searched through a series of methods:

- Gravitational microlensing,
- Compact-binary GWs,
- Hawking-radiation bursts (not seen by gamma ray telescopes), etc.

PBH bounds in red



Recent and controversial bounds are shown as dashed lines



Details in → [B. Kavanagh PBHbounds \(Github\)](#)



Weakly Interacting Massive Particles (WIMPs)

The evolution of the **WIMP** number density n is governed by the Boltzmann equation

$$\frac{dn}{dt} + 3Hn = -\langle\sigma v\rangle\left(n^2 - \left(n_{\text{eq}}\right)^2\right)$$

Freeze out when the interaction rate falls below the Hubble rate


$$\Gamma = n_{\text{eq}}\langle\sigma v\rangle \lesssim H$$



Weakly Interacting Massive Particles (WIMPs)


The evolution of the **WIMP** number density n is governed by the Boltzmann equation

- $z \ll 1 : T \gg M$, WIMPs are relativistic and abundant.
- $z \sim 1 : T \sim M$, WIMPs become non-relativistic.
- $z \sim 20 - 30$: typical freeze-out region.
- $z \gg 1$: abundance $Y = n/s$ becomes constant: the relic density is fixed.




The WIMP Miracle

The precise numerical result for the special value σv , which reproduces the cosmological DM abundance turns out to ***miraculously correspond*** to a typical weak annihilation cross section.




Hunting WIMPs


→ fig 4.6 of [M. Cirelli, A. Strumia and J. Zupan \(2024\)](#)



Recoil events in nuclei or electrons in a highly shielded and closely monitored underground detector




DM annihilations/decays in astrophysical environments, by searching for signatures in cosmic rays, photons or neutrinos arriving at Earth.



Production of DM in a controlled environment (e.g., pp collisions at LHC, e^+e^- collisions at lower energy machines or in beam dump experiments). Presence of DM deduced via missing energy.

Could also happen in astrophysical environment but with tiny rates


Hunting WIMPs



WIMPs can be detected through the tiny recoil energy produced when a WIMP collides with an atomic nucleus in a detector buried underground to shield from background noise


Latest estimates in →
F. Calore et al. (2025)

Incomplete sample of
experimental searches.
For more see, e.g.,
→ [M. Cirelli, A. Strumia
and J. Zupan \(2024\)](#)



Hunting WIMPs: Sensitivity Curve

The reach is worst at low and high mass, best in between — set by two competing effects:



Low mass →

Recoil energy falls below the detector's energy threshold — signals vanish.

$$E_R \simeq \frac{2 \mu^2 v^2}{m_N}$$

Sweet spot →

Energy transfer peaks when the WIMP mass $\approx$ the nucleus mass — tens of GeV.


High mass →

The number density drops as $1/\text{mass}$, so the rate falls; the limit weakens $\propto \text{mass}$.

$$n_\chi = \frac{\rho_\chi}{m_\chi}$$

Two knobs set the reach: the **recoil threshold** (low mass) and the **WIMP flux** (high mass).

Hunting WIMPs




→ [M. Cirelli, A. Strumia and J. Zupan \(2024\)](#)

Direct detection bounds for spin-independent DM–nucleon scattering, assuming DM mass at around the Z boson mass.

Part 2: Axioms



Axions

Axions are among the most studied DM candidates



World-wide effort to detect it.

Axions



The θ -term

QCD Lagrangian

$$L = \underbrace{-\frac{1}{4}G^2}_{\text{Gluon kinetic term}} + \underbrace{\bar{q}(i\gamma^\mu D_\mu - m_q)q}_{\text{Quark kinetic and mass term, plus quark gluon interactions}} + \underbrace{\theta \frac{1}{32\pi^2} G_{\mu\nu}^a \tilde{G}^{a\mu\nu}}_{\text{\(\theta\)-term: Violates CP}}$$

The $\theta \frac{\alpha_s}{8\pi} G_{\mu\nu}^a \tilde{G}^{a\mu\nu}$ term is a total derivative. However, there are configurations which contribute to the action integral. These are called **instantons** $\Rightarrow$ this term cannot be thrown away and has phenomenological consequences.

One consequence is to give mass to the η' meson. In fact, all the terms which explicitly break the symmetry (so, both mass term and anomaly term) contribute to the mass of the Goldstone boson.

The θ -term

See, e.g., Schwartz, Ch. 30.5.2

It is hard to calculate the η' mass. One method is lattice.

Veneziano and Witten (1979) showed that the η' mass is related to the topological susceptibility $\chi_t \equiv \langle (G\tilde{G})(G\tilde{G}) \rangle$ as

$$\chi_t = \frac{f_\pi^2}{12} (m_\eta^2 + m_{\eta'}^2 - 2m_K^2)$$



→ if $G\tilde{G}$ had no effect then $\chi_t = 0$ and the η' mass would be small. In particular, in the chiral limit $m_\eta, m_K \rightarrow 0$ one would find also $m_{\eta'} = 0$.

→ The experimental evidence that η' is not a Goldstone boson (it is “heavy”) suggests that the instantons do play a role in QCD.

The θ -term

- The vacuum energy depends on θ
It is minimized for $\theta = 0$
- The nucleon masses depend on θ
 $\Rightarrow$ the neutron lifetime $\Rightarrow$ nucleosynthesis depends on θ .

$$m_n - m_p \simeq (1.29 + 0.21 \theta^2 + \mathcal{O}(\theta^4)) \text{ MeV}$$



C. Vafa, E. Witten, Phys. Rev. Lett. 53 (1984) 535

• L. Ubaldi, Phys. Rev. D81 (2010) 025011

• M. Dine, L. Stephenson Haskins, L. Ubaldi, D. Xu, JHEP 05 (2018) 171

• Lee, Meißner, Olive, Shifman, Vonk, Phys.Rev.Res. 2 (2020) 3, 033392 (2020)


See Nick Houston talk at
 $\rightarrow$ Axions beyond boundaries, [GGI-2023](#)

The θ -term and Strong CP

The θ -term generates a neutron EDM

$$\rightarrow d_n \approx 2.4(1.0) \times 10^{-16} \theta e \cdot \text{cm}$$

M. Pospelov, A. Ritz (2020)



This gives the strongest constraint

See PhD dissertation by Drew Backhouse, University of Oxford (2021), [arXiv:2108.04285](https://arxiv.org/abs/2108.04285), for a pedagogical introduction

The latest experimental search found

$$|d_n^{\text{exp}}| < 1.8 \cdot 10^{-26} e \text{ cm} \quad (90 \% \text{ CL}) \quad \Rightarrow \quad \theta \lesssim 10^{-10}$$

Abel et al., Phys.Rev.Lett. 124 (2020) 8, 081803

Strong CP Problem

Peccei-Quinn Solution

Peccei and Quinn solution:

→ θ is dynamical → Settles to 0 (Vafa-Witten theorem)

Peccei, Quinn (1977)



Very similar to the $m_u = 0$ solution.
However, the new axial symmetry
is added *by hand*

The Axion

$$L = -\frac{1}{4}G_{\mu\nu}^a G^{a\mu\nu} + \bar{q}(i\gamma^\mu D_\mu - \bar{q}M)q - \theta \frac{1}{32\pi^2} G_{\mu\nu}^a \tilde{G}^{a\mu\nu} + \frac{1}{2}(\partial_\mu a)^2 + \frac{a}{f_a} \frac{g_s^2}{32\pi^2} G\tilde{G} + \dots$$

$$\theta \rightarrow \theta + a/f_a$$

Axion

Note the need for a new energy scale, f_a , from a pure dimensional argument.

This is related (but not necessarily equal to) the scale at which $U(1)_{PQ}$ is spontaneously broken.


S. Weinberg Phys. Rev. Lett. 40 (1978) 223-226;

F. Wilczek Phys. Rev. Lett. 40 (1978) 279-282



Some facts about QCD Axions

1- The axion potential can be calculated in QCD



Axions can get mass only from terms that break explicitly the PQ symmetry. The only thing that does that is the anomaly. So, the axion mass can only come from $G\tilde{G}$.


Not surprisingly, the resulting mass term is related to the topological susceptibility χ_T , just like in the case of the η' .

$$m_a \simeq 5.7 \left(\frac{10^{12} \text{ GeV}}{f_a} \right) \mu\text{eV}$$

Grilli di Cortona et al., JHEP 1601 (2016)

Some facts about QCD Axions

1- The axion potential can be calculated in QCD



Relaxing the $m_a - f_a$ relation requires substantial changes to QCD, for example the addition of a new QCD sector. This is all theoretically possible. However, here we will disregard this possibility for now. See sec. 6.6.2 of L. Di Luzio, M.G., Nardi, Visinelli, Phys.Rept. 870 (2020) for some options.

$$m_a \simeq 5.7 \left(\frac{10^{12} \text{ GeV}}{f_a} \right) \mu\text{eV}$$

Grilli di Cortona et al., JHEP 1601 (2016)

Some facts about QCD Axions

1- The axion potential can be calculated in QCD

2- The axion couplings are model dependent.


There are 2 sources for the axion couplings:

- The model independent $aG\tilde{G}$ coupling; it generates couplings to quarks and photons. Not to electrons (at tree level)
- Model dependent contributions from the specific UV completion

Some facts about QCD Axions

1- The axion potential can be calculated in QCD

2- The axion couplings are model dependent.

$$L_{int} = C_{af} \frac{\partial_\mu a}{2f_a} \bar{f} \gamma_5 \gamma_\mu f + \frac{1}{4} C_{a\gamma} \frac{\alpha}{2\pi f_a} a F \tilde{F}$$


Fermions (electrons, nucleons)

Photons

Spin-density coupling with matter

Two photon coupling of the form $a \mathbf{E} \cdot \mathbf{B}$, from electromagnetic anomaly

Some facts about QCD Axions

1- The axion potential can be calculated in QCD

2- The axion couplings are model dependent.

$$L_{int} = C_{af} \frac{\partial_\mu a}{2f_a} \bar{f} \gamma_5 \gamma_\mu f + \frac{1}{4} C_{a\gamma} \frac{\alpha}{2\pi f_a} a F \tilde{F}$$



Some facts about QCD Axions

1- The axion potential can be calculated in QCD

2- The axion couplings are model dependent.

$$L_{int} = C_{af} \frac{\partial_\mu a}{2f_a} \bar{f} \gamma_5 \gamma_\mu f + \frac{1}{4} C_{a\gamma} \frac{\alpha}{2\pi f_a} a F \tilde{F}$$

f_a suppression (Remember $m_a \propto 1/f_a$)

In general: light $\leftrightarrow$ weakly coupled

Some facts about QCD Axions

1- The axion potential can be calculated in QCD

2- The axion couplings are model dependent.

$$L_{int} = C_{af} \frac{\partial_\mu a}{2f_a} \bar{f} \gamma_5 \gamma_\mu f + \frac{1}{4} C_{a\gamma} \frac{\alpha}{2\pi f_a} a F \tilde{F}$$

QCD contribution to the couplings can be substantially changed according to the particular UV completion...

... in principle, the coupling to fermions can be significantly reduced: $C_{af} \rightarrow 0$.

However, difficult for C_{aN}

- Di Luzio, Mescia, Nardi, Panci, Ziegler, [Phys.Rev.Lett. 120 \(2018\)](#)
- M. Badziak, K. Harigaya, [arXiv:2301.09647](#) (2023);
- F. Takahashi, W. Yin, [arXiv:2301.10757](#) (2023)

Some facts about QCD Axions

1- The axion potential can be calculated in QCD

2- The axion couplings are model dependent.

$$L_{int} = C_{af} \frac{\partial_\mu a}{2f_a} \bar{f} \gamma_5 \gamma_\mu f + \frac{1}{4} C_{a\gamma} \frac{\alpha}{2\pi f_a} a F \tilde{F}$$

$$C_{a\gamma} = \frac{E}{N} - \frac{2}{3} \frac{4m_d + m_u}{m_u + m_d} = \frac{E}{N} - 1.92(4)$$

... The coupling to photons is also model dependent and can (in principle) be tuned to $C_{a\gamma} \ll 1$

Model dependent contribution


Model independent contribution

Some facts about QCD Axions

- 1- The axion potential can be calculated in QCD
- 2- The axion couplings are model dependent.
- 3- Axions contribute to Dark Matter .

- Preskill, Wise and Wilczek (1983)
- Abbott and Sikivie (1983)
- Dine and Fischler (1983)

The PQ solution is dynamical, not instantaneous




The universe has a finite age

The axion field is still oscillating today around the CP-conserving minimum of the potential

→ Axion is CDM

The axion/ALP parameter space



ALP =
Axion-Like
Particle

Part 3:

Axion Sources

and Detection

Natural Axion/ALP sources

Cosmological

[	Realignment	Non-thermal: CDM	$E_a \simeq m_a$
	Topological Defects		
	Thermal	HDM	

Astrophysical

[	Stellar	Thermal	$E_a \sim T_{\text{core}}, \text{ keV or 100 MeV (SN)}$
		Nuclear	$E_a \sim \text{keV, MeV}$
		Resonant $\gamma + B \rightarrow a + B$	$E_a \sim \omega_{pl} \text{ (eV in the sun)}$
	Galactic/ Extra-Galactic	Diffuse SN background	$\sim 100 \text{ MeV}$
		Conversion of photons in B_{ext}	High to very high energy

Natural Axion/ALP sources

Cosmological

[	Realignment	]	<u>Direct</u> : Haloscopes
	Topological Defects		
	Thermal		<u>Indirect</u> : cosmological probes, telescopes ($a \rightarrow \gamma\gamma$), ...

Astrophysical

[	Stellar	[	Thermal	]	<u>Direct</u> : Helioscopes $\rightarrow$ Sun, SN(?)
			Nuclear		
			Resonant $\gamma + B \rightarrow a + B$		<u>Indirect</u> : NuSTAR/Fermi ($a + B \rightarrow \gamma$)
[	Galactic/ Extra-Galactic	[	Diffuse SN background	]	Fermi: <u>Indirect</u> ($a + B \rightarrow \gamma$)
			Conversion of photons in B_{ext}		<u>Indirect</u> ($a + B \rightarrow \gamma$) Chandra, NuSTAR, Fermi...

Axions as Dark Matter

- In spite of being light, axions — and Axion Like Particles (ALPs) — can be **CDM**, thanks to non-thermal production mechanisms.

Very light DM behaves like waves.

Requires specific technologies. Extremely fruitful research field.

“Discovery of dark matter waves would provide a glimpse into the earliest moments in the origin of the universe and the laws of nature at ultrahigh energies, beyond what can be probed in colliders”

→ Kolb et al., <https://doi.org/10.2172/1659757>, 2018

Axions as Dark Matter

For an observed DM density of $\rho \sim 0.2 - 0.56 \text{ GeV/cm}^3$

Axion number density


$$n_a \sim \rho_a/m_a \sim 4 \times 10^{13} \left(\frac{10\mu\text{eV}}{m_a} \right) \text{ axions /cm}^3$$

Axion DB wavelength ($v \sim 10^{-3}$): $\lambda_{\text{dB}} = \frac{h}{p} = \frac{h}{mv} \simeq 120 \text{ m} \left(\frac{10\mu\text{eV}}{m_a} \right)$

So, the axion occupation number= number of axions in a reduced de Broglie volume, is $\sim 10^3$ for $m_a \sim 1 \text{ eV}$.

→ Very large. Axion DM behaves like waves.

Axion Dark matter




See C. O'Hare, [Axion cosmology](#) COST (2023) Training School

The axion emerges from the phase of the scalar field

$$\phi = \frac{1}{\sqrt{2}} (v_a + \rho) e^{ia/v_a}$$


Axion Dark matter



At low enough temperature the PQ symmetry is spontaneously broken.

→ the θ angle is randomly selected. The **axion** is the massless d.o.f. in the flat direction of the potential

Axion Dark matter




Around the QCD transition ($T \approx 1 \text{ GeV}$), axions acquire a mass

$$m_a(T)^2 = \begin{cases} m_a^2 \left(\frac{T}{T_{\text{QCD}}} \right)^{-n} & \text{for } T > T_{\text{QCD}} \\ m_a^2 & \text{for } T < T_{\text{QCD}} \end{cases}$$

with $T_{\text{QCD}} \approx 150 \text{ MeV}$ and $n \approx 8$.

Axion Dark matter



So, we have an axion field, which evolves according to the

→ Equation of motion

$$\ddot{a} + 3H\dot{a} + m_a^2 a = 0$$



Axion Dark matter

→ Equation of motion

$$\ddot{a} + 3H\dot{a} + m_a^2 a = 0$$

Numerical solution

See [Olivier Wantz, E.P.S. Shellard](#)



Axion Dark Matter: Some Considerations

For a spatially uniform real field $a(t)$ with potential $V = \frac{1}{2}m_a^2 a^2$,

$$\rho = \frac{1}{2}\dot{a}^2 + \frac{1}{2}m_a^2 a^2, \quad P = \frac{1}{2}\dot{a}^2 - \frac{1}{2}m_a^2 a^2, \quad w \equiv \frac{P}{\rho}.$$


If $m_a \gg H$, the friction can be ignored and the axion oscillates approximately as

$$a(t) = A(t)\cos(m_a t + \theta) \quad \text{Thus,}$$

$$\Rightarrow \langle \dot{a}^2 \rangle = m_a^2 A^2 \langle \sin^2 \rangle = \frac{1}{2}m_a^2 A^2, \quad \langle a^2 \rangle = \frac{1}{2}A^2$$

$$\Rightarrow \langle P \rangle = \frac{1}{2} \langle \dot{a}^2 - m_a^2 a^2 \rangle = 0, \quad \langle \rho \rangle = \frac{1}{2}m_a^2 A^2.$$

$$\text{Also: } A(t) \propto R(t)^{-3/2} \quad \Rightarrow \quad \langle \rho_a \rangle \propto R(t)^{-3},$$




Cold dark matter!

$$\Omega_a h^2 = 0.12 \left(\frac{\theta_i}{2.155} \right)^2 \left(\frac{28 \mu\text{eV}}{m_a} \right)^{1.16}$$

Axion Dark matter: Pre-inflationary scenario

Scenario 1: Pre-inflationary axions



Calculable axion abundance
but unknown initial conditions

Axion Dark matter: Post-inflationary scenario

Predictable initial angle.

Axion abundance depends also on production from topological defects

Estimating contribution from topological defects is very difficult. Numerical simulations still make very different predictions.

Important numerical advances thanks to Adaptive Mesh Refinement

M. Buschmann et al., *Nature Commun.* 13 (2022) 1, 1049.

Still controversial. More work required.

- M. Gorghetto, E. Hardy, [arXiv:2212.13263](https://arxiv.org/abs/2212.13263)
- O'Hare, Pierobon, Redondo, Wong, [Phys.Rev.D 105 \(2022\)](https://arxiv.org/abs/2201.00009)
- M. Gorghetto, E. Hardy, G. Villadoro, *SciPost Phys.* 10, 050 (2021)





Figure Credits: C. O'Hare (2021)

<https://cajohare.files.wordpress.com/2021/10/axions.pdf>

Dark Matter Mass Predictions



From [C. O'Hare](#)

Predictions in post-inflation point at $m_a \gtrsim \mu\text{eV}$

Hunting down the elusive axions


Our focus: $m_a \lesssim 1\text{eV}$.

Specific experimental challenges, and combination of specific know-hows

→ cross-disciplinary technology transfer:

E.g., high-field magnets, super-conduction, RF techniques, X-ray optics, low background detection, low radioactivity techniques, quantum sensors, atomic physics, etc.

Most (but not all) detection strategies make use of the axion-photon coupling $g_{a\gamma}$
 $a \vec{E} \cdot \vec{B}$



Sikivie (1983)


The Cavity Haloscope

Ideal, cylindrical cavity, with perfectly metallic surfaces

Standing EM waves in a cavity

$$\begin{aligned}\epsilon \nabla \cdot \mathbf{E} &= \rho_o, \\ \nabla \cdot \mathbf{B} &= 0, \\ \nabla \times \mathbf{E} + \dot{\mathbf{B}} &= 0, \\ \mu^{-1} \nabla \times \mathbf{B} - \epsilon \dot{\mathbf{E}} &= \mathbf{J}_o.\end{aligned}$$

Sources



Two kinds of modes TE_{nmp} and TM_{nmp}

Symbol	Meaning	Allowed values
m	azimuthal order (number of field variations in ϕ)	$0, 1, 2, \dots$
n	radial order (n-th root of Bessel or its derivative)	$1, 2, \dots$
p	axial order (number of half-wavelengths along z)	$0, 1, 2, \dots$


The Cavity Haloscope

Add axions + Uniform $\mathbf{B}_{\text{Lab}}$ $L_{a\gamma} = -\frac{g_{a\gamma\gamma}}{4} a F_{\mu\nu} \tilde{F}^{\mu\nu} \propto a \mathbf{B} \cdot \mathbf{E}$

Standing EM waves in a cavity

$$\begin{aligned} \epsilon \nabla \cdot \mathbf{E} &= \cancel{\rho_o} + \rho_a, \\ \nabla \cdot \mathbf{B} &= 0, \\ \nabla \times \mathbf{E} + \dot{\mathbf{B}} &= 0, \\ \mu^{-1} \nabla \times \mathbf{B} - \epsilon \dot{\mathbf{E}} &= \cancel{\mathbf{J}_o} + \mathbf{J}_a. \end{aligned}$$

Sources



$$\rho_a = g_{a\gamma\gamma} \nabla a \cdot \mathbf{B}_0$$

$$\mathbf{J}_a = -g_{a\gamma\gamma} (\dot{\mathbf{B}} + (\nabla a) \times \mathbf{E})$$


The Cavity Haloscope

Add axions + Uniform $\mathbf{B}_{\text{Lab}}$ $L_{a\gamma} = -\frac{g_{a\gamma\gamma}}{4} a F_{\mu\nu} \tilde{F}^{\mu\nu} \propto a \mathbf{B} \cdot \mathbf{E}$

Standing EM waves in a cavity

$$\begin{aligned} \epsilon \nabla \cdot \mathbf{E} &= \cancel{\rho_o} + \rho_a, \\ \nabla \cdot \mathbf{B} &= 0, \\ \nabla \times \mathbf{E} + \dot{\mathbf{B}} &= 0, \\ \mu^{-1} \nabla \times \mathbf{B} - \epsilon \dot{\mathbf{E}} &= \cancel{\mathbf{J}_o} + \mathbf{J}_a. \end{aligned}$$

Sources



$$\rho_a = g_{a\gamma\gamma} \cancel{\nabla a} \cdot \mathbf{B}_0$$

Uniform background of CDM axions



$$\mathbf{J}_a = -g_{a\gamma\gamma} (\dot{\mathbf{B}} + (\cancel{\nabla a} \times \mathbf{E}))$$

The Cavity Haloscope

Standing EM waves in a cavity

$$L_{a\gamma} \propto a \mathbf{B} \cdot \mathbf{E}$$

$\approx \mathbf{B}_{\text{lab}}$ Cavity mode $\mathbf{E}_{\text{cav}}$



Only TM_{nmp} modes $\rightarrow$ lowest mode TM_{010} .

The Cavity Haloscope: the axion as a drive

- Expand the cavity field in its eigenmodes, each with a frequency set by the geometry.
- Every mode amplitude then obeys a driven, damped harmonic oscillator.
- Losses (walls + readout) enter as ω/Q , with $Q \sim 10^4\text{--}10^5$

Eigenmode expansion:

$$\mathbf{A}(\mathbf{r}, t) = \sum_n \psi_n(t) \mathbf{e}_n(\mathbf{r})$$

Each amplitude obeys a driven, damped oscillator — the axion current is the drive:

$$\ddot{\psi}_n + \frac{\omega_n}{Q} \dot{\psi}_n + \omega_n^2 \psi_n = g_{a\gamma\gamma} \dot{a} \frac{\int_V \mathbf{B}_0 \cdot \mathbf{e}_n dV}{\int_V |\mathbf{e}_n|^2 dV}$$

Driven oscillator: the axion is the **drive**, **Q** sets the sharpness.


The Cavity Haloscope: resonance & tuning

- The response is resonantly enhanced when the cavity frequency matches the axion frequency.
- On resonance the amplitude grows by about Q ; a little off resonance, the signal collapses.
- Tune the cavity size to sweep across the unknown axion mass.

Steady-state amplitude:

$$\psi_n \propto \frac{g_{a\gamma\gamma} \int_V \mathbf{B}_0 \cdot \mathbf{e}_n dV}{\omega_n^2 - \omega_a^2 - i\gamma_n \omega_a}$$

$$\omega_a \simeq \omega_n \implies |\psi_n| \sim Q$$



Signal peaks sharply at $\omega_a = \omega_n$; width $\sim \omega_n / Q$.

Tuning — cylinder TM_{010} mode:


$$f_{TM_{010}} \simeq \frac{2.405 c}{2\pi R}$$

Scanning = **stepping the cavity frequency** across candidate axion masses.

The Cavity Haloscope: resonance & tuning

- The response is resonantly enhanced when the cavity frequency matches the axion frequency
- On resonance, little off resonance response
- Tune the cavity to axion mass

Large B_0 and large volume help



Steady-state amplitude:

$$\psi_n \propto \frac{g_{a\gamma\gamma} \int_V \mathbf{B}_0 \cdot \mathbf{e}_n dV}{\omega_n^2 - \omega_a^2 - i\gamma_n \omega_a}$$

$$\omega_a \simeq \omega_n \implies |\psi_n| \sim Q$$

Signal peaks sharply at $\omega_a = \omega_n$; width $\sim \omega_n / Q$.

Tuning — cylinder TM_{010} mode:

$$f_{TM_{010}} \simeq \frac{2.405 c}{2\pi R}$$

Scanning = **stepping the cavity frequency** across candidate axion masses.

The Cavity Haloscope: Signal Power

- The standard haloscope power formula (time-averaged and calculated at resonance) is

$$P = \left\langle \int_V \mathbf{J}_a \cdot \mathbf{E} dV \right\rangle_t$$

$$P = g_{a\gamma\gamma}^2 \frac{\rho_a}{m_a} B_0^2 V C Q$$

- coupling & local dark-matter density $\rightarrow$ particle physics
- field B_0 and volume V $\rightarrow$ build strong and big
- form factor C (mode overlap) and quality factor Q

Signal power is of order **10^{-23} W**


The Cavity Haloscope: Noise

Practically:

- **thermal noise** attributed to the blackbody radiation by the physical temperature of the cavity
- noise added by the RF readout chain


D. Kim et al. [JCAP 03 \(2020\) 066](#)

$$\text{SNR} \equiv \frac{P_{\text{signal}}}{\delta P_{\text{noise}}}$$



Dicke's radiometer equation

$$\frac{S}{N} = \frac{P_s}{T_{\text{sys}}} \sqrt{\frac{\Delta t}{\Delta \nu}}$$



Wish list:

- long scanning time
- narrow band
- low T_{sys}


The Cavity Haloscope: Noise

Practically:

- **thermal noise** attributed to the blackbody radiation by the physical temperature of the cavity
- noise added by the RF readout chain

D. Kim et al. [JCAP 03 \(2020\) 066](#)


$$\text{SNR} \equiv \frac{P_{\text{signal}}}{\delta P_{\text{noise}}}$$



Dicke's radiometer equation

$$\frac{S}{N} = \frac{P_s}{T_{\text{sys}}} \sqrt{\frac{\Delta t}{\Delta \nu}}$$

This is what controls the scanning time!



$$T_{\text{sys}} = T_{\text{phy}} + T_{\text{add}}$$

System noise

Amplifier

Physical temperature

Wish list:


- long scanning time
- narrow band
- low T_{sys}

The Cavity Haloscope: Noise

Modern approaches include

- low-temperature refrigerators
- Quantum Sensing to increase the S/N and reduce the scanning time

*Recent ERC grant awarded to
explore quantum technology
in axion cavity detection*



DarkQuantum



Cavity Experiments

Micro eV mass range:

Most experience.

- ADMX: proven sensitivity to **few μeV**

High Mass:

Difficult: higher masses requires smaller volume $\rightarrow$ lower sensitivity. Possible solutions:

- lower noise
- matching more cavities or new multicavity designs
- More powerful magnets
- Dielectric Haloscopes
- ...


Many projects: HAYSTAC, CAPP, MADMAX,

Low Mass:


Technologically simpler. However, expensive. Needs large magnets

- FLASH (concept)
- BabyIAXO (to be built)
- IAXO

Current Haloscope Experiments




Current Haloscope Experiments



See → [C. O'Hare, AxionLimits](#) (updated to July 2026)


Future Haloscope Experiments

Many proposals to probe the higher mass region



From FIP white paper (2022)


Opportunities: Axion Telescopes?



Stellar Axion Flux,
 $g_{a\gamma} = 10^{-11} \text{ GeV}^{-1}$, $g_{ai} = 0$, $m_a = 0$


→ P. Carenza, M.G., J. Isern, A. Mirizzi, O. Straniero, *Phys.Rept.* 1117 (2025) 1-102

Opportunities: Axion Telescopes?



→ P. Carenza, M.G., J. Isern, A. Mirizzi, O. Straniero, *Phys.Rept.* 1117 (2025) 1-102

Opportunities: Axion Telescopes?




Stellar Axion Flux,
 $g_{a\gamma} = 10^{-11} \text{ GeV}^{-1}$, $g_{ai} = 0$, $m_a = 0$

→ P. Carenza, M.G., J. Isern, A. Mirizzi, O. Straniero, *Phys.Rept.* 1117 (2025) 1-102

Opportunities: Axion Telescopes?


- Telescope for Super Giant stars.
- Find exact evolutionary stage in post He-burning.
- Pin down SN explosion time.
- Space X-ray detector. Needs $m_a < \text{neV}$



→ *P. Carenza, M.G., J. Isern, A. Mirizzi, O. Straniero, [Phys.Rept. 1117 \(2025\) 1-102](#)*


Opportunities: Axion Telescopes?

- Could reveal temperature, density and pions abundance.
- Best options with Fermi LAT or similar for $m_a \lesssim$ a few neV.
- Possible with helioscopes, for $m_a \lesssim$ eV but much less statistics.




→ P. Carenza, M.G., J. Isern, A. Mirizzi, O. Straniero, *Phys.Rept.* 1117 (2025) 1-102

Solar Axions

Coupling	Process	Energy
$g_{a\gamma}$	Primakoff (E)	$\sim (3 - 4) \text{ keV}$
	Primakoff (B) 	$\sim (10 - 200) \text{ eV (LP)}$ $\lesssim 1 \text{ keV (TP)}$
g_{ae}	ABC e . g . , $e + Z_e \rightarrow Ze + e + a$	$\sim 1 \text{ keV}$
g_{aN}	nuclear reactions	
	$p + d \rightarrow {}^3\text{He} + a$	5.5 MeV
	Nuclear de-excitation	
	${}^{57}\text{Fe}^* \rightarrow {}^{57}\text{Fe} + a$	14.4 keV
	${}^7\text{Li}^* \rightarrow {}^7\text{Li} + a$	0.478 MeV
	${}^{83}\text{Kr}^* \rightarrow {}^{83}\text{Kr} + a$	9.4 keV

Hunting Solar Axions: Sikivie Helioscope

P. Sikivie PRL 51:1415 (1983)



Rescalable: increasing collecting area, length, and B.


Sensitivity

$$P_{a\gamma} = \left(\frac{g_{a\gamma} BL}{2} \right)^2 \frac{\sin^2(qL/2)}{(qL/2)^2}$$

B = magnetic field

L = magnet length

q = momentum transfer



$$q \simeq \frac{m_a^2 - m_\gamma^2}{2\omega}$$

Sensitivity


$$P_{a\gamma} = \left(\frac{g_{a\gamma} B L}{2} \right)^2 \frac{\sin^2(qL/2)}{(qL/2)^2}$$

B = magnetic field

L = magnet length

q = momentum transfer

$$q \simeq \frac{m_a^2 - m_\gamma^2}{2\omega}$$




Sensitivity

$$P_{a\gamma} = \left(\frac{g_{a\gamma} BL}{2} \right)^2 \frac{\sin^2(qL/2)}{(qL/2)^2}$$

B = magnetic field

L = magnet length

q = momentum transfer




$$q \simeq \frac{m_a^2 - m_\gamma^2}{2\omega}$$

Van Bibber et al.
Phys.Rev. D 39:2089 (1989)

The CERN Axion Solar Telescope (CAST)

Reached the HB bound for the first time

V. Anastassopoulos, et al., Nature Phys. 13 (2017)



Decommissioned LHC test magnet,
 $B=9\text{T}$, $D=43\text{ mm}$, $L=9.3\text{ m}$

$\sim 2\text{ h}$ tracking/day


X-ray optics

The International AXion Observatory

Irastorza et al., JCAP 06 (2011)


E. Armengaud et al., JINST 9 (2014)

- Large toroidal 8-coil magnet $L = \sim 20$ m
- 8 bores: 60 cm diameter each
- 8 x-ray telescopes + 8 detection systems
- Rotating platform with services



IAXO will consist of a superconducting toroid magnet with eight custom x-ray telescopes that focus the reconverted photons onto ultra-low background detectors.


The International AXion Observatory



Strong potential to
probe the high
($>\text{meV}$) mass region
(stellar window)

Physics potential of the International Axion
Observatory (IAXO) JCAP 1906 (2019) 047

BabylAXO




- Prototype: Intermediate experimental stage before IAXO
- Test & improve all systems. Risk mitigation for full IAXO.
- Physics: expected relevant physics outcome (~100 x CAST FOM)



2017 ERC advanced grant by I. Irastorza to support the development

DESY PRC endorsed BabylAXO in

Light Shining Through a Wall



Everything is under control. However,
signal suppressed as $g_{a\gamma}^4$

Relativistic axions:

$$P_{a\gamma} = \left(\frac{g_{a\gamma} B L}{2} \right)^2 \frac{\sin^2(qL/2)}{(qL/2)^2}$$

B = magnetic field

L = magnet length

q = momentum transferred

$$q \simeq \frac{m_a^2 - m_\gamma^2}{2\omega}$$

Lost of coherence for $qL \gtrsim 1$

Light Shining Through a Wall

→ **ALPS** @ DESY and the **OSQAR** @ CERN are active LSW experiments.

Use powerful accelerator dipole magnets, from HERA (ALPS) and LHC (OSQAR).

In both cases the sensitivity drops above $m_a \sim 10^{-4}$ eV .

→ **CROWS** experiment @ CERN

Uses a different wavelength: microwaves!

Large scale MW LSW studied and proposed in the literature → **STAX**

LSW at X-rays also explored in the past (not large power)

→ Next Gen: **JURIA**. Concept being discussed at the Physics Beyond Colliders (PBC) group at CERN

- $L \sim 1\text{km}$, $B \sim 13\text{T}$, $P \sim 2.5\text{ MW}$,... Very challenging parameters...
- Physics case to be settled (it may depend if positive signal in other exps)

Relativistic axions:

$$P_{a\gamma} = \left(\frac{g_{a\gamma} B L}{2} \right)^2 \frac{\sin^2(qL/2)}{(qL/2)^2}$$

B = magnetic field

L = magnet length

q = momentum transferred


$$q \simeq \frac{m_a^2 - m_\gamma^2}{2\omega}$$

Lost of coherence for $qL \gtrsim 1$

Polarization experiments


Dichroism:

Production of real particles



Ellipticity:

Production of massive virtual particles




PVLAS experiment: study QED vacuum birefringence (standard effect), but also sensitivity to ALPs:

→ Under discussion at PBC: VMB@CERN

→ Most likely next step VMB LAB AT AEI. Fresh updates in [A. Ejlli, Corfu June 2026](#)

Axions Around the World



Conclusions and Comments

- A lot of research on Dark Matter and specifically on Axions
- Important progress in last years in the attempts to detect axions in all fronts:
 - ✓ Huge progress with cavity **haloscopes** + several new proposals
 - ✓ Major **LSW** experiment starting (ALPS II)
 - ✓ Major new **Helioscope** proposal under progress (IAXO)
 - ✓ Several **innovative ideas**
- Most interesting parameter space accessible to Next Gen Experiments
→ a **groundbreaking discovery is possible**
- **Axions** with couplings at the current thresholds would excellent **astrophysical messengers**.

To know more about Axions

- A soft and enjoyable introduction is the article by David J. E. Marsh:
Axions for amateurs [arXiv:2308.16003](https://arxiv.org/abs/2308.16003)
- More technical: L. Di Luzio, M.G., E. Nardi, L. Visinelli, *The landscape of QCD axion models*, [Phys.Rept. 870 \(2020\)](https://arxiv.org/abs/2007.13473)
- Especially good for theory:
 - *Villadoro Lectures* GGI 2015, GGI 2023
 - Anson Hook, *TASI Lectures*, [arXiv:1812.02669](https://arxiv.org/abs/1812.02669)
- For astrophysics:
 - P. Carenza, M.G., A. Mirizzi, J. Isern, O. Straniero, *Axion astrophysics*, [Phys.Rept. 1117 \(2025\)](https://arxiv.org/abs/2501.1117)
 - A. Caputo and G. Raffelt, *Astrophysical Axion Bounds: The 2024 Edition*, [PoS COSMICWISPers \(2024\) 041](https://arxiv.org/abs/2405.04101)
- For experiments: I. Irastorza, J. Redondo, *New experimental approaches in the search for axion-like particles*, [Prog.Part.Nucl.Phys. 102 \(2018\)](https://arxiv.org/abs/1708.07467)

... and in general, on Dark Matter

- Very recent, interesting and comprehensive review
→ M. Cirelli, A. Strumia and J. Zupan (2024)
- Also interesting the very recent short review Bozorgnia et al.
→ Dark Matter Candidates and Searches
- For Axion Cosmology, See C. O'Hare, Axion cosmology
COST (2023) Training School
- There is a Long evidence for Dark Matter.
See → A History of Dark Matter, by G. Bertone and D. Hooper (2018)

Cosmic WISPers in the Dark Universe

Join our network and working groups
→ [COST CA21106](#)

