



IBD selection and backgrounds in JUNO

Mingyuan Wang on behalf of **JUNO** Collaboration


Institute of High Energy Physics, CAS

mywang@ihep.ac.cn

Introduction

The Jiangmen Underground Neutrino Observatory (JUNO)

- Neutrino source
 - Mainly reactor neutrino $\bar{\nu}_e$
 - $\bar{\nu}_e + p \rightarrow e^+ + n$ (**Inverse beta decay**)
- Target substance
 - 20 kt liquid scintillator
 - 35.4-m-diameter sphere
- Physical goals
 - Neutrino mass ordering
 - Oscillation parameters
 - Other physics such as supernova neutrino




IBD selection strategy

1. Fiducial volume (FV)

- Exclude signals beyond the FV
- Suppress edge effects
 - Reject most of the external background
 - Better reconstruction performance
 - More subtle energy leakage
- Efficiency **~80.6%**


2. Veto

3. Pairing and multiplicity cut



IBD selection strategy



1. Fiducial volume
2. Veto
 1. Muon veto (5 ms)
Remove muon-induced secondary particles
Remove afterpulses
 2. Spallation neutron veto (4 m, 1.2 s)
Veto long-lived cosmogenic isotopes
(eg. ^9Li / ^8He)
- Livetime & Geometric Efficiency
~93.6%
3. Pairing and multiplicity cut





IBD selection strategy

1. Fiducial volume
2. Veto
3. Pairing and multiplicity cut
 - Exclude single events
 - Improve SNR
 - Efficiency $\sim 95.1\% \times 97.4\%$

Strategy




Result



Residual backgrounds


1. Cosmogenic isotope ${}^9\text{Li}/{}^8\text{He}$

- The most important background in current analysis, ~ 4.4 cpd
- Suppressed by **spallation neutron veto**
- Will be further suppressed by a track-based veto around reconstructed muon trajectories



2. ${}^{214}\text{Bi} - {}^{214}\text{Po}$ related

3.



Residual backgrounds

1. Cosmogenic isotope


2. $^{214}\text{Bi} - ^{214}\text{Po}$ related

- Energy region typically differs from IBD candidates
- Alpha long tail in LS, caused by α -p scattering, elevating the visible energy into the delayed signal window with probability $\sim 10^{-4}$

- Rate ~ 0.18 cpd in **JUNO**


- Also reported by **SNO+**

3.



Summary

- In the first measurement of reactor neutrino oscillations at JUNO, the overall selection efficiency is $69.9 \pm 1.6\%$
- **Fiducial volume**
 - Largest source of efficiency loss
 - Dominant to efficiency uncertainty
 - Will be improved by better vertex reconstruction
- Cosmogenic isotope ${}^9\text{Li}/{}^8\text{He}$
 - Primary background
 - Can be further suppressed by improved muon reconstruction



Thanks!