



Status of the RICOCHET experiment at ILL

RENCONTRES DE MORIOND – EW+U 2026 SESSION

SERRA Renaud on behalf of the RICOCHET collaboration

17/03/2026



Coherent Elastic Neutrino-Nucleus Scattering

- Theorized in 1974 by Daniel Z. Freedman
- First measurement in 2017 by COHERENT

Cross section predicted by the Standard Model:

$$\frac{d\sigma_{\nu/N}}{dE_R} = \frac{G_F^2}{4\pi} Q_W^2 m_N \left(1 - \frac{m_N E_R}{2 E_\nu^2} \right) F^2(E_R) \sim N^2$$





Diagram of CEνNS



Interactions cross-sections for various processes [L. Bailly-Salins, EPS-HEP 2025]

1. Standard Model: **Weinberg angle**
2. Sensitivity to **new physics**:
 - non-standard ν -quark interaction
 - neutrino magnetic moment μ_ν
 - new mediator boson Z'
3. Background for **dark matter**:
 - same signature and inevitability
 - experimental threshold





Illustration of theoretical interaction rates for the Ricochet experiment

[RICOCHET Collaboration, J Low Temp Phys 212 (2023), 127]



RICOCHET's shielding and cryostat


Only **8.8 m** to the reactor $\rightarrow \phi_{\nu} \sim 10^{12} \text{ cm}^{-2} \cdot \text{s}^{-1}$

Radioactive **background** divided into 3 categories

- **Cosmogenic**: from cosmic particles (μ , n , etc.)
- **Reactogenic**: isotopes activated by reactor neutrons
- **Radiogenic**: natural background of the environment

Four-part background **mitigation** strategy

- **Passive** shielding: borated PE, lead and soft iron
- **Active** shielding: muon veto
- **Discrimination** based on recoil type: **ER/NR**
- **ON-OFF** reactor cycle data subtraction



Planar detector operating diagram

Mini-CryoCube: 3 cryogenic bolometers

CryoCube: 2 × 3 mini-CryoCube

Full payload: 18 × 42 g ⇒ kg-scale

Technologies: PL & FID detectors

Strategy: ionization & heat signals

Discrimination: ionization yield or quenching $Q = \frac{E_{ion}}{E_{recoil}}$

- **Electronic Recoil (ER):** from γ & $e^- \Rightarrow Q = 1$
- **Nuclear Recoil (NR):** from ν & $n \Rightarrow Q \approx 0.2$
- **Low-Energy Excess (LEE) or Heat-Only (HO):** $Q = 0$




Illustration of ionization yield as a function of recoil energy

[Adapted from Phys. Rev. D 112 (2025), 112019]

Gamma background: electronic recoil band

Major contribution: $^{41}_{18}\text{Ar}$ decay during ON

Experimental result in region of interest*

- RUN014: $R_{ON-OFF}^{PL167} = (980 \pm 174) \text{ DRU}^\dagger$

GEANT4 simulations: in air and shielding[§]


- RUN014: $R_{GEANT4}^{PL167} = (839 \pm 69) \text{ DRU}$

- RUN015: $R_{GEANT4}^{PL167} = (117 \pm 14) \text{ DRU}$

86% reduction in event rate between RUNs

Due to improvements in **shielding sealing**

* from 2 to 7 keV & [†] 1 DRU = 1 event/day/kg/keV & [§] used in MCMC reconstruction



RICOCHET successfully operating at ILL


2024
2025
2026

- **RUN012:** cryogenic test at ILL
- **RUN013:** first test of PL detectors
- **RUN014:** background validation test
- **RUN015:** first test with FID detectors
- **RUN016:** first science run with full payload
- **RUN017:** second science run

Baseline resolutions RUN014: $\left\{ \begin{array}{l} 40 \text{ eV}_{ee} \text{ ionization} \\ 50\text{--}80 \text{ eV}_{ph} \text{ phonon} \end{array} \right.$

Development in progress

- **Q-Array:** superconducting crystals
- **Silicon detector:** complementary targets



CEvNS event rate spectra in Ge and Si crystals


[Julien Billard et al, JCAP11(2018), 016]

**Thank you for your
attention!**

Back-up


Neutrino-nucleus interactions

[M. Cadeddu et al, EPL 143 (2023), 34001]




Inelastic and incoherent :

$$\lambda_{Z^0} \ll 2R$$



Elastic and incoherent:

$$\lambda_{Z^0} \leq 2R$$



Elastic and coherent :

$$\lambda_{Z^0} \geq 2R$$

E_R : recoil energy

E_ν : neutrino energy

G_F : Fermi's constant

m_N : nucleus mass

$F(E_R)$: form factor

Q_w : nucleus weak charge

N : number of neutrons

Z : number of protons

θ_w : weak mixing angle



Cross section predicted by the Standard Model :

$$\frac{d\sigma_{\nu/N}}{dE_R} = \frac{G_F^2}{4\pi} Q_w^2 m_N \left(1 - \frac{m_N E_R}{2 E_\nu^2} \right) F^2(E_R)$$

With : $Q_w = N - (1 - 4 \sin^2 \theta_w) Z$

Unification of the electromagnetic force and the weak interaction :

- Represents the rotation caused by spontaneous symmetry breaking
- Proportion of weak and electromagnetic interactions in particle interactions



$\sin^2 \theta_w$ experimental measurements

Standard Model prediction :

$$\frac{d\sigma_{SM}}{dE_R} = \frac{G_F^2}{4\pi} Q_W^2 m_N \left(1 - \frac{m_N E_R}{2 E_\nu^2} \right) F^2(E_R)$$

Non-zero magnetic moment :

$$\frac{d\sigma_\mu}{dE_R} = \frac{\pi \alpha^2 \mu_\nu^2 Z^2}{m_e} \left(\frac{1}{E_R} - \frac{1}{E_\nu} + \frac{E_R}{4E_\nu^2} \right) F^2(E_R)$$

Scalar boson :


$$\frac{d\sigma_S}{dE_R} = \frac{Q_S^2}{4\pi} \frac{m_N^2 E_R}{E_\nu^2 (q^2 + m_S^2)^2} F^2(E_R) \quad Q_S = g_{\nu,S} g_{q,S} (15.1 Z + 14 N)$$

Tensor boson :




$$\frac{d\sigma_T}{dE_R} = \frac{Q_T^2}{2\pi} \frac{m_N (4 E_\nu^2 - m_N E_R)}{E_\nu^2 (q^2 + m_T^2)^2} F^2(E_R) \quad Q_T = g_{\nu,T} g_{q,T} (0.85 Z + 0,08 N)$$

Vector boson :



$$\frac{d\sigma_V}{dE_R} \propto \frac{d\sigma_{SM}}{dE_R} \quad Q_W \rightarrow Q_{tot} = Q_W - \frac{\sqrt{2}}{G_F q^2} \frac{g_{\nu,V} [Z (2 g_{u,V} + g_{d,V}) + N (g_{u,V} + 2 g_{d,V})]}{q^2 + m_V^2}$$



Gamma background sources





From top to bottom :
40-K and 60-Co




From left to right : $^{232}\text{-Th}$, $^{238}\text{-U}$ and $^{235}\text{-U}$ chains

Ionization spectra comparison






[RICOCHET Collaboration, EPJ C 84 (2024), 186]



Contents of the **RUN014/015** GEANT4 simulation :

- two-part lead shielding : save calculation time
 - transfer channel slab
 - Cable (electronics)
 - ambient air near cryostat
 - polyethylene shielding
 - N active detectors (only planar for the moment)
 - detector mass : $m_{Bolo} = 42.038 \text{ g}$
- } radiogenic contamination
- } ^{41}Ar contamination








Markov chain Monte Carlo algorithm :

$$\mathcal{L}(\vec{\theta}) \propto \exp\left(-\frac{1}{2}\chi^2\right) \Rightarrow \log[\mathcal{L}(\vec{\theta})] \propto -\frac{1}{2}\chi^2$$

$$\chi^2 = \sum_{i=1}^{N_{bin}} \left[w_i \times \frac{S_i^{exp.} - S_i^{sim.}(s_k^{mat.}, a_k^{mat.})}{\sigma_i} \right]^2$$

- $S_i^{exp.}$: experimental spectrum (RED167 & RED237)
- $S_i^{sim.}(s_k, a_k) = \sum_k a_k s_k$: simulated spectra model
- $\sigma_i = \sqrt{N_i}$: Poissonian errors on spectra
- w_i : weight function to constrain the model



Gelman-Rubin coefficient :

$$\hat{R}_1 = 1.0033 \approx 1$$

$$\hat{R}_2 = 1.0031 \approx 1$$

Gamma background reconstruction

