



Searches for new physics at high object masses



Andrea Malara


On behalf of the CMS Collaboration

March 2026



The Standard Model -- A triumph...

Precise description of particle interactions up to the electroweak scale!!





The image shows a chalkboard with the title "Standard Model" underlined at the top. Below the title, the Lagrangian is written in white chalk:

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + i\bar{\psi}\not{D}\psi + h.c. \\ + \psi_i y_{ij} \psi_j \phi + h.c. + |D_\mu\phi|^2 - V(\phi)$$



The Standard Model -- A triumph... with puzzles

Precise description of particle interactions up to the electro



Standard Model


$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + i\bar{\psi} \not{D} \psi + h.c. + \psi_i Y_{ij} \psi_j \phi + h.c. + |D_\mu \phi|^2 - V(\phi)$$



Hunting for New Physics at the Energy Frontier



► Beyond the Standard Model (BSM) theory models predict:

- Additional heavy bosons
- Dark matter candidates
- Extra-dimensions
-
- with properties largely unknown





► Broad range of signatures to analyse at the LHC:

- Hadronic final states
- Leptons
- missing transverse momentum






- CMS explores a wide range of models and experimental signatures
- **This talk highlights:**
 - Heavy vector resonances: Spin-1 combination: (**Run 2**)
 - New Run3 results on $W' \rightarrow l\nu$ searches (**Run 3**)
 - High-mass jet signatures: Dijet and pair of dijet searches (**Run 2 & Run 3**)


How can **New Physics (BSM)** appear ?




New Physics at high masses

How can New Physics (BSM) appear ?









> 2σ local excess in
 $W' \rightarrow tb$ at 3.4-4.4 TeV
for CMS-B2G-20-012



> 2σ local excess in
 $Z' \rightarrow \ell\ell$ at 2.7-3.4 TeV
for CMS-EXO-19-019




> 3σ local excess in
 $V' \rightarrow VV \rightarrow had$ at 2-3 TeV
for CMS-B2G-20-009




> 2σ local excess in
 $W' \rightarrow W(\ell\nu)V(qq)$ at 1TeV
for CMS-B2G-19-002



> 2σ local excess in
 $Z' \rightarrow Z(\nu\nu)H(bb)$ at 1TeV
for CMS-B2G-19-006





- ▶ Combination of 16 searches:
 - ▶ Maximal coverage of final states
 - ▶ Different sensitivity across the parameter space
- ▶ Interpretation under the HVT framework
 - ▶ Simplified framework for spin-1 resonances
 - ▶ Couplings controlled by 2 parameters





Channel	Prod. modes	$m_{V'}$ range [TeV]	Leptons	Jets	Reference
$Z' \rightarrow \tau^+ \tau^-$	DY	0.5–4.0	$\ell \tau_h$ or $2\tau_h$	—	[47]
$Z' \rightarrow \ell^+ \ell^-$	DY	0.8–5.5	2ℓ	—	[48]
$W' \rightarrow \ell \bar{\nu}$	DY	0.8–6.0	1ℓ	—	[49]
$W' \rightarrow \tau \bar{\nu}$	DY	0.8–6.0	$1\tau_h$	—	[50]
$V' \rightarrow q\bar{q}^{(\prime)}$	DY	1.8–8.7	0ℓ	jj	[51]
$W' \rightarrow t\bar{b}$ (0ℓ)	DY	1.0–4.0	0ℓ	$j^t j^b$	[52]
$W' \rightarrow t\bar{b}$ (1ℓ)	DY	2.0–6.0	1ℓ	$j^b j^b$	[53]
$Z' \rightarrow t\bar{t}$ (1ℓ)	DY	0.8–5.0	1ℓ	j^t or jjj^b	[54]
$Z' \rightarrow ZH$ ($2\ell 2b$)	DY + VBF	0.8–5.0	2ℓ	j^{bb}	[55]
$Z' \rightarrow ZH$ ($2\nu 2b$)	DY + VBF	0.8–5.0	0ℓ	j^{bb}	[55]
$Z' \rightarrow ZH$ ($2\ell 2q$)	DY	1.2–6.0	2ℓ	j^H	[56]
$Z' \rightarrow ZH$ ($2\nu 2q$)	DY	1.2–6.0	0ℓ	j^H	[56]
$W' \rightarrow WZ$ ($2q 2\ell$)	DY	0.6–2.0	2ℓ	j^V or jj	[57]
$W' \rightarrow WZ$ ($2q 2\nu$)	DY + VBF	1.0–4.5	0ℓ	j^V	[58]
$V' \rightarrow WV/WH$	DY + VBF	1.0–4.5	1ℓ	j^V	[59]
$V' \rightarrow VV/VH$	DY + VBF	1.3–6.0	0ℓ	$j^V j^V$	[20]





- ▶ $Z' \rightarrow \ell\ell$ and $W' \rightarrow \ell\nu$ as leading channels
- ▶ Previous 2.7–3.4 TeV fluctuation in $Z' \rightarrow \ell\ell$ reduced
- ▶ No significant deviations observed




- ▶ $V' \rightarrow VV/VH \rightarrow had$ as leading channel
- ▶ Run-2 tensions at 1–3 TeV are reduced
- ▶ No significant deviations observed



- Sensitivity depends on the relative couplings to fermions and bosons
- No significant deviation observed up to 6 TeV



- Beyond the assumption of fermion universality
- $g_{q_3} - g_H$ and $g_{q_3} - g_{l_3}$ entirely excluded





One of the most sensitive probes of heavy vector bosons

- ▶ Clean experimental signature
- ▶ High- p_T isolated lepton (e or μ)
- ▶ Large missing transverse momentum

- ▶ Transverse mass (m_T) used as the sensitive variable



$$m_T = \sqrt{2 \cdot p_T^l p_T^{\text{miss}} \cdot (1 - \cos \Delta\phi(p_T^l, p_T^{\text{miss}}))}$$

- ▶ Leading uncertainties:
 - ▶ Exp: Jet and lepton energy scale
 - ▶ Theo: PDF and high- m_T cross section



- ▶ No significant deviation from the SM
 - ▶ SSM W' mass excluded up to ~ 6 TeV
 - ▶ Model-independent interpretation




- ▶ Improved sensitivity compared to Run2
 - ▶ Higher center-of-mass energy
 - ▶ Lower trigger threshold for e-channel




Dataset	e channel Obs (exp) TeV	μ channel Obs (exp) TeV	e+ μ channel Obs (exp) TeV
Run2	5.4 (5.3)	5.6 (5.5)	5.7 (5.6)
2022+2023	5.7 (5.9)	5.6 (6.0)	5.9 (6.2)

New Physics at high masses


How can New Physics (BSM) appear ?



Nonresonant effects in dijet production


- ▶ Angular distribution (χ_{dijet}) as the sensitive variable
 - ▶ Scattering angle of the dijet system
 - ▶ Small dependence on theoretical uncertainties
- 
- ▶ Normalised differential cross sections:
 - ▶ Compared to NNLO QCD + EW predictions
 - ▶ Detector response accounted for via response smearing
 - ▶ Leading uncertainties:
 - ▶ Exp: Jet energy scale and resolution
 - ▶ Theo: Scale choice of μ_R/μ_F (updated wrt. previous results)


$$\chi_{dijet} = \exp(|y_1 - y_2|) \sim \frac{1 + \cos(\theta^*)}{1 - \cos(\theta^*)}$$



- ▶ No significant excess observed
- ▶ Several interpretations provided
 - ▶ Quark compositeness
 - ▶ DM models
 - ▶ Extra-dimensions
 - ▶ SMEFT

Model	Observed limit	Expected limit
CI $\Lambda_{LL/RR}^+$ (NLO)	17 TeV	18_{-2}^{+20} TeV
CI $\Lambda_{LL/RR}^-$ (NLO)	37 TeV	34_{-5}^{+8} TeV
CI Λ_{VV}^+ (NLO)	<19 and 27–31 TeV	41_{-22}^{+17} TeV
CI Λ_{VV}^- (NLO)	50 TeV	45_{-7}^{+11} TeV
CI Λ_{AA}^+ (NLO)	19 TeV	39_{-21}^{+16} TeV
CI Λ_{AA}^- (NLO)	49 TeV	44_{-7}^{+11} TeV
CI $\Lambda_{(V-A)}^+$ (NLO)	14.9 TeV	$14.0_{-1.4}^{+1.8}$ TeV
CI $\Lambda_{(V-A)}^-$ (NLO)	15.7 TeV	$14.6_{-1.6}^{+2.2}$ TeV
ADD Λ_T (GRW)	13.4 TeV	14.2 ± 1.0 TeV
ADD M_S (HLZ) $n_{ED} = 2$	15.0 TeV	16.0 ± 1.1 TeV
ADD M_S (HLZ) $n_{ED} = 3$	15.9 TeV	16.9 ± 1.2 TeV
ADD M_S (HLZ) $n_{ED} = 4$	13.4 TeV	14.2 ± 1.0 TeV
ADD M_S (HLZ) $n_{ED} = 5$	12.1 TeV	12.8 ± 0.9 TeV
ADD M_S (HLZ) $n_{ED} = 6$	11.3 TeV	11.9 ± 0.8 TeV
QBH M_{QBH} (ADD, $n_{ED} = 6$)	8.5 TeV	9.1 ± 0.3 TeV
QBH M_{QBH} (RS1)	6.3 TeV	6.8 ± 0.4 TeV
DM mediator m_{med}	6.2 TeV	6.2 ± 0.8 TeV
ALP c_g/f_a	0.42 TeV^{-1}	$0.38 \pm 0.03 \text{ TeV}^{-1}$
SMEFT C_G/Λ^2	0.0076 TeV^{-2}	$0.0081 \pm 0.0012 \text{ TeV}^{-2}$






Pair production of new particles X decaying to dijets

Resonant dijet pair production



- ▶ Two dijet resonances reconstructed with 4 jets
- ▶ Sensitive observables:
 - ▶ Four-jet mass m_{4j}
 - ▶ Average dijet mass \overline{m}_{2j}
- ▶ Highest mass events previously observed with Run2 data:
 - ▶ CMS: $m_{4j} = 8 \text{ TeV}$; $\overline{m}_{2j} = 2 \text{ TeV}$
- ▶ Current highest mass events with Run3 data:
 - ▶ CMS: $m_{4j} = 6.2 \text{ TeV}$; $\overline{m}_{2j} = 2 \text{ TeV}$





- ▶ Fit of the falling spectrum of m_{4j}
 - ▶ Performed in \overline{m}_{2j}/m_{4j} bins \rightarrow to minimise background sculpting
 - ▶ Discrete profiling is used for background modelling

▶ No significant excess observed


- ▶ Interpretation in diquark models (D2D) 90 fb⁻¹ (13.6 TeV)



- ▶ Intriguing excess from Run2 analysis not confirmed
- ▶ But more data still to be analysed





Pair-produced dijet resonances with b-jets



Pair production of new particles X decaying to dijets with 2 b-jets



Dijet pair production with 2 b jets

- ▶ Resonant:
 - ▶ m_{4j} and \overline{m}_{2j} as the sensitive variables
- ▶ Non-resonant:
 - ▶ \overline{m}_{2j} as the sensitive variable
- ▶ Current highest mass events with Run2 data:
 - ▶ CMS: $m_{4j} = 5.5 \text{ TeV}$; $\overline{m}_{2j} = 1.5 \text{ TeV}$



CMS Experiment at LHC, CERN
Run/Event : 320010 / 23731487
Lumi section: 18

- ▶ Fit of the falling spectrum of m_{4j}
 - ▶ $\overline{m_{2j}}$ for the non-resonant interpretation
- ▶ No significant excess observed
- ▶ Limits set on:
 - ▶ pair-produced diquarks (D2D)
 - ▶ Top squark (RPV)



Conclusions



CMS continues to probe new physics at the highest accessible energies

- ▶ Heavy vector resonances
 - ▶ Run2 legacy combination of 16 channels
 - ▶ Sensitivity up to 6 TeV
 - ▶ W' search showcases improved sensitivity using Run3 data
- ▶ High-mass searches
 - ▶ Dijet angular analysis probing non-resonant BSM effects
 - ▶ Pair-produced dijet resonances
- ▶ No significant deviation observed from the SM observed (yet)
 - ▶ Constantly improving techniques for high-mass signatures
 - ▶ Large Run3 dataset to be explored
 - ▶ Sensitivity will continue to grow



Thank you for your attention!

~~~ **Additional Material** ~~~



- ▶ Heavy Vector Triplet (HVT):
  - ▶ Simplified framework for heavy spin-1 resonances
  - ▶ Massive charged and neutral bosons (Z' and W')
  - ▶ Similarities with the SM EW sector
  - ▶ Couplings to fermions and bosons controlled by  $g_F$  and  $g_H$



$$\mathcal{L}_{\mathcal{W}}^{\text{int}} = -c_q \frac{g^2}{g_V} \mathcal{W}_{\mu}^a \bar{q}_k \gamma^{\mu} \frac{\sigma_a}{2} q_k - c_l \frac{g^2}{g_V} \mathcal{W}_{\mu}^a \bar{l}_k \gamma^{\mu} \frac{\sigma_a}{2} l_k - c_H g_V (\mathcal{W}_{\mu}^a H^{\dagger} \frac{\sigma_a}{2} i D^{\mu} H + \text{h.c.})$$





- ▶  $V' \rightarrow qq$  as leading channel
- ▶ Tension at 3.4-4.4 TeV from  $W' \rightarrow tb$  not confirmed
- ▶ No other significant deviations





- ▶  $V' \rightarrow VV/VH \rightarrow had$  as leading channel
- ▶ Tensions at 1 TeV in  $W' \rightarrow W(\ell\nu)V(qq)$  and  $Z' \rightarrow Z(\nu\nu)H(bb)$  reduced
- ▶ Tensions at 2-3 TeV in  $V' \rightarrow VV \rightarrow had$  reduced
- ▶ No other significant deviations

▶ Mixing of DY and VBF signals, when available!!



- ▶  $Z' \rightarrow \ell\ell$  and  $W' \rightarrow \ell\nu$  as leading channels
- ▶ Tension at 2.7-3.4 TeV from  $Z' \rightarrow \ell\ell$  reduced
- ▶ No other significant deviations



- ▶  $V' \rightarrow VV/VH \rightarrow had$  as leading channel<sup>1</sup>
- ▶ Tensions at 1 TeV in  $W' \rightarrow W(\ell\nu)V(qq)$  and  $Z' \rightarrow Z(\nu\nu)H(bb)$  reduced
- ▶ Tensions at 2-3 TeV in  $V' \rightarrow VV \rightarrow had$  reduced
- ▶ No other significant deviations

