

The Impact of Plasma Non-Linearities on Dark Matter Searches

Anson Hook

University of
Maryland

w / J. Huang
and M. Shalaby
2510.13956



1 Line

Heating A Plasma is Hard

Dark Photon Dark Matter

1) Well Motivated

- Common in any extra dimensional theory
- We've seen massive vectors before

2) Very restricted interactions - Kinetic Mixing

- Acts like a photon but with a smaller α

What does it look like?

Dark Photon Dark Matter

- **Dark Matter is non-relativistic (it's clumped up)**

$$v_{\text{now}} \sim 10^{-3}$$

- **A massive vector at rest is all Electric field**


$$\vec{E} = E_0 \cos(m_{DM} t) \hat{u}$$

What does it look like?

Looking for dark matter is as simple as looking for an oscillating electric field!


E.g. A leading experimental proposal is using a (very good) antenna to look for this E field

Early Universe Plasma



Early Universe Plasma

$$\vec{E} = E_0 \cos(m_{DM} t) \hat{u}$$



Early Universe Plasma

Famously plasmas oscillate at plasma frequency



Like a swing, if frequencies match the amplitude grows a lot!

Early Universe Plasma

Too much energy in plasma and model is excluded

$$\mu \text{ and } y \text{ distortions} - \Delta\rho \lesssim 10^{-4, -5} \rho_\gamma$$

If the plasma frequencies match, inject energy until the above occurs and you are excluded!

Early Universe Plasma

Too much energy in plasma and model is excluded

$$\mu \text{ and } y \text{ distortions} - \Delta\rho \lesssim 10^{-4,-5} \rho_\gamma$$

~~If the plasma frequencies match, inject energy until the above occurs and you are excluded!~~

Non-linearities push you off resonance when

$$\Delta\rho \sim \rho_{e^-}^{K.E.} \sim 10^{-10} \rho_\gamma$$

Early Universe Plasma

A swing

$$\partial_t^2 \phi + \omega^2 \sin \phi \approx \partial_t^2 \phi + \omega^2 \phi - \lambda \phi^3 = J \cos(\omega t)$$

ϕ grows linearly with time until it goes off resonance because of non-linear term


$$\omega_{\text{eff}}^2 = \omega^2 - 3\lambda\phi^2$$

Numerical Simulation


Theory is hard - Pictures/Videos are easy

Step 1 - Write Mathematica Code

Electric Field



Electron Phase Space



Numerical Simulation

Theory is hard - Pictures/Videos are easy

Step 1 - Write Mathematica Code

- **Grows as expected**
- **Eventually growth stops - off resonance**
- **Additional surprise - fragments!**

Numerical Simulation

Theory is hard - Pictures/Videos are easy

Step 1 - Write Mathematica Code

- **Grows as expected**
- **Eventually growth stops - off resonance**
- **Additional surprise - fragments!**

Step 2 - Write better C code


Step 3 - Realize why people do this for a living

Step 4 - Ask an expert for help

Numerical Simulation


Large Electric Field Limit

1. Initial instability from acoustic waves
2. Acoustic waves die off and a mess remains



Numerical Simulation


Large Electric Field Limit



Stops at $t \sim 1/\omega_{p,ion}$



Analytic E (no nonlinearities)

Numerical E



Numerical Simulation


Large Electric Field Limit



Numerical Simulation



Small Electric Field Limit

1. Starts to grow
2. Stops when acoustic waves (charge neutral) are important
3. Acoustic waves die off and a mess remains





Numerical Simulation

Small Electric Field Limit




Numerical Simulation

Small Electric Field Limit




Numerical Simulation

Even Smaller Electric Field Limit



Limits

Experimental/Lab-based bounds are now the world's leading constraints on dark photon dark matter



Conclusion

Plasma Physics is complicated!

- **Numerically - Energy deposition is capped at kinetic energy of electrons**
 - Annoying for energy deposition constraints
 - Full 3D simulations need to be done to capture the proper final state
- **Perhaps there are other cool new ways to look for the presence of a background oscillating electric field with plasmas**