

Higgs Rare Decays

On behalf of the ATLAS and CMS Collaborations

Andrzej Novak

March 20, 2026



**Phil
Harris
Lab@MIT**

60th Rencontres de Moriond (EWK), La Thuile (Italy)


Landscape of Higgs Searches

Over a decade since discovery

- Mass measured to per-mille precision
- ~90% of branching ratio observed

~24 Millions of Higgs bosons at ATLAS & CMS each

- Expecting ~170 Millions by end of HL-LHC
- Enabling rare decay exploration




Landscape of Higgs Searches

Over a decade since discovery

- Mass measured to per-mille precision
- ~90% of branching ratio observed

~24 Millions of Higgs bosons at ATLAS & CMS each

- Expecting ~170 Millions by end of HL-LHC
- Enabling rare decay exploration
- Rare decay channels now accessible:
 - $H \rightarrow c\bar{c}$ (+ H+c associated production)
 - $H \rightarrow Z\gamma$
 - $H \rightarrow \mu\mu$
 - γH



Outline

Analysis	ATLAS	CMS
$H \rightarrow cc$ (VBF)	HIGP-2024-21 (Nov 2025)	—
$H \rightarrow cc$ (ttH)	—	HIG-24-018 (Phys. Rev. Lett. 136 (2026) 011801)
$H \rightarrow WW + c$	—	HIG-24-009 (Aug 2025)
$H \rightarrow \gamma\gamma + c$	—	HIG-23-010 (JHEP 11 (2025) 060)
$H \rightarrow Z\gamma$	HIGP-2024-19 (Jul 2025)	HIG-25-012 (BRAND NEW!)
$H \rightarrow \mu\mu$	HIGP-2024-11 (Phys. Rev. Lett. 135 (2025) 231802)	—
γH	—	HIG-23-011 (Phys. Rev. D 112 (2025) 11200)



H → cc

2nd-generation quark Yukawa coupling

- Key SM test; with HH, next Higgs frontier


Experimentally challenging

- 20× smaller BR than H → bb
- Less distinct charm signature in detector





[ATLAS] $H \rightarrow cc$ in VBF Channel (Partial Run2+Run3 89fb^{-1})

- **First (at LHC) VBF $H \rightarrow c\bar{c}$ search**
 - Enabled by **inclusive VBF trigger**
 - No b-tag, $m_{jj} > 1\text{ TeV}$, $> 90/95\%$ eff
- **Key Strategy Elements:**
 - **GN2 transformer tagger** for flavour identification
 - WP-cut and count passing b/c jets
 - **Adversarial NN** decorrelated from di-jet Higgs-candidate mass
 - Six SRs; joint fit floating $\mu_{c\bar{c}}$ and $\mu_{b\bar{b}}$
 - **Data-driven multijet (99%)**
 - Shape from SR6, bias unc. 26–56% of stat.





[ATLAS] $H \rightarrow c\bar{c}$ in VBF Channel (Partial Run2+Run3 89fb^{-1})

- $H \rightarrow c\bar{c}$:
 - $\mu_{c\bar{c}} = 18 \pm 13 \rightarrow \mu_{c\bar{c}}^{95\%CL} < 41$ (28)
 - VBF+VH: $|\kappa_c| < 4.7$ (3.9 *exp*) – 8% over VH-alone
- $H \rightarrow b\bar{b}$:
 - $\mu = 0.97^{+0.57}_{-0.50}$
 - Combination w/ Run2 $3.2\sigma \rightarrow$ first VBF $H \rightarrow b\bar{b}$ evidence
- Statistically limited





[CMS] $H \rightarrow cc$ in $t\bar{t}H$ Channel (Run2 138fb^{-1})

- First (at LHC) $t\bar{t}H H \rightarrow c\bar{c}$ search (0L/1L/2L)
- Key challenges: many jets, complex combinatorics
- ParticleNet jet tagger (11 cats, $2\times$ rejection)
- Particle Transformer event classifier (10-class, no explicit reco).
- $t\bar{t}+HF$ background constrained from data CRs with floating normalizations.
- Validated with $t\bar{t}Z(Z \rightarrow c\bar{c}/bb)$



[CMS] $H \rightarrow cc$ in $t\bar{t}H$ Channel (Run2 138fb^{-1})


- $t\bar{t}H(H \rightarrow c\bar{c})$: $\mu_{c\bar{c}}^{95\%CL} < 7.8$ (8.7)
- Combined with VH: $|\kappa_c| < 3.5$ (2.7 *exp*) – best to date
- $t\bar{t}H(H \rightarrow b\bar{b})$: $\mu = 0.91^{+0.26}_{-0.22}$
- Sensitivity comparable to VH



H + c: Probing the Charm Yukawa Coupling

Complementary probe of κ_c

- $H \rightarrow c\bar{c}$ limited by ϵ^2 c-tagging penalty + QCD bkg
- $pp \rightarrow Hc$ alternative (Brivio et al., PRL 115)
 - Only 1 c-tag (charm at production vertex)
 - y_c interferes with $ggH \rightarrow$ **linear** κ_c sensitivity
 - $\Gamma \propto \kappa_c^2$ (weak); $\sigma \propto \kappa_c$ (tighter bounds)
- Two analyses presented:
 - $H \rightarrow \gamma\gamma + c$ – clean mass peak, high S/B
 - Earlier ATLAS result [HIGG-2021-06](#) – $5.3 \pm 3.2\text{pb}$ ($2.9 \pm 3.1\text{pb}$ expected)
 - $H \rightarrow WW + c$ – larger BR, leptonic W









[CMS] $H \rightarrow \gamma\gamma + c$ (Run2 138fb⁻¹)

- $H \rightarrow \gamma\gamma$ for clean $m_{\gamma\gamma}$ peak.
- Two BDTs (vs resonant Higgs bkg + vs continuum)
- 27 cats (9 x 3 years) \times $m_{\gamma\gamma}$ fit. c-tagging enriches cH over ggH.
- $\mu_{cH}^{95\%CL} < 243$ (355 exp); $|\kappa_c| < 38.1$ (72.5 exp)
- Complementary to $H \rightarrow c\bar{c}$; sensitivity $\propto \kappa_c^4 \rightarrow$ scales rapidly

CMS




138 fb⁻¹ (13 TeV)



$m_{\gamma\gamma}$ (GeV) 12


[CMS] $H \rightarrow WW + c$ (Run2 138fb⁻¹)

- $H \rightarrow WW$ larger BR (21.4% vs 0.227%) x harder bkg.
- $e\mu$ suppresses Z+jets (21.4% incl \rightarrow 1% leptonic)
- DeepJet CvsL/CvsB c-tagging for cH
- Two BDTs: D_{bkg} (vs tt) + $D_{\text{H-bkg}}$ (vs other Higgs)



[CMS] $H \rightarrow WW + c$ (Run2 138fb⁻¹)

- $\mu_{cH}^{95\%CL} < 1065$ (506 *exp*)
- Combined $cH(\gamma\gamma+WW)$: $|\kappa_c| < 47$ (**51 *exp***)
- Weaker than $H(\gamma\gamma)$ mostly due to $t\bar{t}$
- Validates cH in new decay mode
- Production-mode κ_c in early days
- Grows with lumi + c-tagging improvements



Higgs-Charm Coupling: Status of κ_c

Analysis	μ obs (exp)	$ \kappa_c $ obs (exp)	Lumi
CMS ttH ($H \rightarrow c\bar{c}$)	< 7.8 (8.7)	< 3.0 (3.3)	138 fb ⁻¹
CMS VH+ttH ($H \rightarrow c\bar{c}$)	< 9.3	< 3.5 (2.7)	138 fb ⁻¹
ATLAS VH ($H \rightarrow c\bar{c}$) upd.	< 11.5 (10.6)	< 4.2	140 fb ⁻¹
CMS VH ($H \rightarrow c\bar{c}$)	< 14.4	< 5.5 (3.4)	138 fb ⁻¹
ATLAS VBF+VH ($H \rightarrow c\bar{c}$)	< 41 (28)	< 4.7 (3.9)	89 fb ⁻¹
ATLAS VH ($H \rightarrow c\bar{c}$) '22	< 26 (31)	< 8.5 (12.4)	139 fb ⁻¹
CMS ggH ($H \rightarrow c\bar{c}$)	< 45	—	138 fb ⁻¹
CMS cH $\rightarrow \gamma\gamma$	< 243 (355)	< 38.1 (72.5)	138 fb ⁻¹
CMS cH $\rightarrow WW$	< 1065 (506)	—	138 fb ⁻¹
ATLAS cH $\rightarrow \gamma\gamma$	$\sigma < 10.6$ (8.8) pb	—	140 fb ⁻¹

- **Best direct $|\kappa_c| = 3.5$** (CMS ttH+VH combo); ATLAS updated VH reaches 4.2
- Direct $H \rightarrow c\bar{c}$ still 8–28 \times SM; cH production probes κ_c via production (complementary)
- Multiple channels converging; HL-LHC + improved c-taggers essential for SM sensitivity


Landscape of Higgs Searches

Over a decade since discovery

- Mass measured to per-mille precision
- ~90% of branching ratio observed



~24 Millions of Higgs bosons at ATLAS & CMS each

- Expecting ~170 Millions by end of HL-LHC
- Enabling rare decay exploration
- Rare decay channels now accessible:
 - $H \rightarrow c\bar{c}$ (+ H+c associated production)
 - $H \rightarrow Z\gamma$
 - $H \rightarrow \mu\mu$
 - γH




[ATLAS] $H \rightarrow Z\gamma$ (Run2 + Partial Run3 '22, '23, '24)

- $Z\gamma$ is loop-induced
- BSM-sensitive to new particles entering loop diagrams
 - Uniquely sensitive to composite Higgs models (1308.2676)
- BR = 0.154%
 - $Z \rightarrow ll$ (6.6%) limits rate.
 - $Z \rightarrow ee/\mu\mu$
- Key Strategy Elements:
 - 13 categories with XGBoost BDTs
 - Improvement over cut-based Run2 analysis
 - Z-mass kin fit improves $m_{Z\gamma}$ resolution
 - +17% electrons
 - +11% muons
 - p_T thresholds relaxed due to improved calibrations





[ATLAS] $H \rightarrow Z\gamma$ (Run2 + Partial Run3 '22, '23, '24)

- Run 3:
 - $\mu = 0.9_{-0.6}^{+0.7} \rightarrow 1.4\sigma$ (1.5 σ exp.)
- **Combined w/ Run2:**
 - $\mu = 1.3_{-0.5}^{+0.6} \rightarrow 2.5\sigma$ (1.9 σ exp.)
 - Best single-experiment (until this week)
 - 19% over ATLAS+CMS Run 2
 - 3.4 σ obs. due to excess in data ($\mu=2.2\pm0.7$)
- Completing EW decay suite
 - [$\gamma\gamma$ \checkmark , ZZ^* \checkmark , WW^* \checkmark , $Z\gamma \rightarrow$ next]



[CMS] $H \rightarrow Z\gamma$ (Run2 + Partial Run3 '22, '23)




- Brand new result approved last week
- Key Strategy Elements:
 - Mass fit across 13 categories - all production modes
 - BDTs used for ggF and VBF categories – 4 each
 - ttH and VH categories are newly included
 - Z kin fit improves mass resolution
 - 10% improvement due to improved triggers
- Run2 reanalyzed with updated methods



[CMS] $H \rightarrow Z\gamma$ (Run2 + Partial Run3 '22, '23)


Preliminary numbers:

- $\mu = 1.10^{+0.52}_{-0.61} \rightarrow 1.9\sigma$ (2.3σ exp.)
- $1.00^{+0.49}_{-0.46}$ expected
- Central value compatible with SM expectation
- Channel compatibility: $p = 0.75$





[ATLAS] $H \rightarrow \mu\mu$ (Run3 '22, '23, '24, 165fb^{-1})

- $H \rightarrow \mu\mu$: narrow resonance in $m_{\mu\mu}$ atop DY (S/B $\sim 10^{-3}$). BR=0.022%.
- 23 categories (ggF/VBF/VH/ttH), BDTs + NN. 5B NLO DY MC, vertex fit (+2% res.).
- Crystal Ball signal ($\sigma \approx 2.8\text{--}3.2$ GeV), bkg fits in $m_{\mu\mu}$ in 110–160 GeV.





[ATLAS] $H \rightarrow \mu\mu$ (Run2 + Run3 '22, '23, '24 - 205fb^{-1})

- Run 3: $\mu = 1.6 \pm 0.6, 2.8\sigma$
- **Combined: $\mu = 1.4 \pm 0.4, 3.4\sigma$ (2.5σ exp) – first ATLAS evidence**
- Direct 2nd-gen Yukawa; syst $\sim 5\%$ – stat-limited
- With CMS (3.0σ Run 2) $\rightarrow 5\sigma$ in reach with full Run 3




[CMS] γH + Light Quark Yukawas

- **Boosted Higgs recoiling against a high-energy photon (trigger)**
 - $H \rightarrow bb$ (boosted, ParticleNet AK8, $p_T > 300$ GeV, mass-decorrelated)
 - $H \rightarrow 4l$ f(MELA). γ -tagged + Untagged.
- **Effectively two analyses:**
 1. γH to probe anomalous couplings
 2. Light-quark Yukawas via $H \rightarrow 4l$.
 - Enhanced κ_q boosts $q\bar{q} \rightarrow H$
 - Simultaneous 4-param fit.
 - ggF loop must be refit simultaneously.







[CMS] γH + Light Quark Yukawas

- $\sigma_{\gamma H} < 16.4 \text{ fb}$
- 4 anomalous HV γ couplings (CP-even/odd)



- $\kappa_u = (0 \pm 1.5) \times 10^3$,
- $\kappa_d = (0 \pm 6.8) \times 10^2$,
- $\kappa_s = 0 \pm 31$
- $\kappa_c = 0.0_{-2.8}^{+2.3}$
- First simultaneous light-quark Yukawa constraints



Summary

Charm Yukawa – Tightest constraints yet

- ▶ CMS ttH+VH sets $|\kappa_c| < 3.5$ (2.7 exp) – best to date. ATLAS VBF H \rightarrow bb first evidence at 3.2σ

H \rightarrow $\mu\mu$ – Observation within reach

- ▶ ATLAS evidence at 3.4σ ($\mu=1.4\pm 0.4$);
- ▶ CMS 3.0σ , Together with ATLAS 5σ likely with full Run 3

H \rightarrow Z γ – EW decay suite nearing completion

- ▶ ATLAS combined 2.5σ ($\mu=1.3$)
- ▶ CMS combined 1.9σ ($\mu=1.1$)





Light quarks – First simultaneous multi-coupling constraints

- ▶ CMS: simultaneous $\kappa_u, \kappa_d, \kappa_s, \kappa_c$. Rules out 1st/2nd-gen = 3rd-gen at $>95\%$ CL

Thank You

Classifier - Jet Tagging

- FCNN - DeepCSV ~2016
- CNN->RNN->FCN - DeepDoubleX (1) ~2017
- GNN - ParticleNet (2) ~2019
- Transformer - ParT (3) ~2022



Classifier – Jet Tagging Calibration

- **No pure bb/cc signal region to calibrate on**
- Use $g \rightarrow bb/cc$ as a **proxy** (not entirely signal-like)
- Two ways of “forcing” a good proxy – compatible results
 - Use **soft-muon** presence to select Higgs-like cc jets
 - Train a **BDT** to select Higgs-like cc jet

