




# Multiboson and VBS measurements

S. Folgueras (Universidad de Oviedo)  
on behalf of the CMS and ATLAS Collaborations



Grant PID2023-147115NB-I00 funded by



MINISTERIO  
DE CIENCIA, INNOVACIÓN  
Y UNIVERSIDADES



AGENCIA  
ESTATAL DE  
INVESTIGACIÓN



Universidad de Oviedo



Funded by  
the European Union



erc  
European Research Council  
Established by the European Commission


# Why multiboson & VBS?

Multiboson LHC

Direct probe of  
EW self-  
interactions

Triple/quartic  
gauge couplings  
and EWSM

Rare  
interactions  
and new  
physics







JHEP 04 (2025) 115

Multiboson measurements offer a comprehensive test of the electroweak sector at the LHC

# Differential diboson



- Inclusive ZZ and ZZjj
- Differential distributions in agreement with NLO
- Sensitivity to neutral TGC and QGC



# $W\gamma$ production: precision probe



- Sufficient precision to discriminate between state-of-the-art theoretical predictions including higher-order NNLO QCD and (virtual) NLO EW.
- Inclusive  $W\gamma$  kinematic observables. RAZ probed.



## Polarization:

- Double differential measurement  $W$  decay angles.
- Probe the polarisation state of the  $W$  and provide direct sensitivity to the  $W$  boson spin-density matrix.
- Agreement with SM

# $W\gamma$ production: precision probe

- Sufficient precision to discriminate between state-of-the-art theoretical predictions including higher-order NNLO QCD and (virtual) NLO EW.



## CP-sensitive observables:

- sensitive to interference caused by CP-violating  $WW\gamma$  interactions

- $O_{NN}$ , constructed from the outputs of a multi-class NN trained to separate interference for a given operator.




$$O_{NN} = P_+ - P_-$$

- In agreement with SM
- Improve the expected limits on the CP-odd operators by x2 compared to the use of  $\Delta\phi_\ell^Y$  alone



# VBS topology





- Two vector bosons
- Two forward jets
- Large  $m_{jj}$ ,
- Large  $\Delta\eta_{jj}$
- VBS directly probes quartic gauge couplings and electroweak symmetry breaking.



VBS is now established experimentally, and Run 3 is the bridge to precision and EFT reach



# VBS observation in semileptonic modes

- Combined  $llqq / \ell\nu qq / \nu\nu qq$  (resolved + merged)
- RNN-based extraction
- Observed significance:  $7.4\sigma$  (expected  $6.1\sigma$ )
- ATLAS also sees  $3.3\sigma$  evidence for  $\geq 1$  longitudinal W in SSWW: the next VBS frontier [[Phys. Rev. Lett. 135 \(2025\) 111802](#)]



# VBS: CMS combination across 7 channels




- Simultaneous fit to fully-leptonic (SSWW, OSWW, WZ, ZZ) and semi-leptonic (WV, ZV) final states
- Run-2 dataset:  $138 \text{ fb}^{-1}$  at 13 TeV
- Measured  $\mu$  values (4-POI model) are all consistent with 1 within uncertainties



Provides a coherent baseline for EFT / aQGC interpretations and future Run-3 updates

# EW ZZ production: VBS fully established

- $\sigma_{EW} = 0.37_{-0.12}^{+0.14}(\text{stat}) \pm 0.06(\text{syst})$
- 3.1 sigma in this channel alone, and 5 sigma when combined with the four-lepton channel,
  - establishing electroweak ZZ production.






Limits on aQGC are derived in terms of dimension-8 EFT operators.

# First measurement in VBS @ 13.6 TeV






- First Run-3 leptonic VBS measurement: both EW  $W^\pm W^\pm$  and EW  $WZ$  observed with Run 3 data.
- **Analysis Strategy:** channels ( $W^\pm W^\pm$  and  $WZ$  leptonic modes), VBS topology cuts, BDT-based  $WZ$  discrimination, and the simultaneous ML fit structure
- **Key Results:**  $>5\sigma$  and  $\sim 7\sigma$  observation significances. Inclusive fiducial cross sections.  $W^\pm W^\pm$  differential.



# And already looking at 2025 data!



- More data is needed to improve sensitivity (to aQGC).
- Distributions in the 2022-25 data sample, both signal regions (with loose  $m_{jj}$  requirement) and control regions.







See Monika's talk this afternoon

CMS-PAS-SMP-25-013


# Triboson production: Run 2 + Run 3

- Triboson production directly probes quartic gauge couplings.
  - Extremely complex analyses, multiple final states to maximize sensitivity.
- CMS: evidence for triboson (WWZ) production at 13.6 TeV with  $4.5\sigma$  ( $5\sigma$ ) observed (expected) significance.
- ATLAS observed VVZ ( $6.4\sigma$ ) and evidence of WWZ



# Outlook

- Diboson measurements validate electroweak sector at precision level.
- VBS now observed in several channels and probes electroweak symmetry breaking
- First Run-3 VBS measurements at  $\sqrt{s} = 13.6$  TeV
- Triboson and radiative processes extend sensitivity to rare gauge interactions
- Run-3 and HL-LHC will transform multiboson physics into a precision probe of the electroweak sector.




# Back-up slides






**Diboson**



**VBS**




**Triboson**

# Longitudinal VBS: same-sign $WWjj$

- Same-sign dileptons + VBS jets (high  $m_{jj}$ , large  $\Delta\eta_{jj}$ )
- Two-stage DNN strategy:
  - (i) inclusive EW SSWW vs backgrounds;
  - (ii) polarisation separation
- Evidence for  $\geq 1$  longitudinal W:  $3.3\sigma$  (expected  $4.0\sigma$ )
- Upper limit for doubly-longitudinal component in fiducial region:  $0.45 \text{ fb}$  (95% CL)


Polarisation measurements:  
disentangle the longitudinal (L)  
and transverse (T) helicity  
fractions of the produced bosons.



# Diboson precision summary




JHEP 08 (2025) 142




Diboson measurements validate EW gauge sector and provide precision baseline for exploring rarer multiboson and VBS processes.


Multiboson LHC



JHEP 11 (2025) 006




JHEP 04 (2025) 115




# The “menu”

- WZ: [JHEP 04 \(2025\) 115](#) (Run 3)
- Zg: 13 TeV ( [arXiv:2601.14102](#)) and 13.6 TeV( [arXiv:2512.08582](#))
- ZZg: [CMS-PAS-SMP-24-014](#) (Run 2)
- VBF g: [arXiv:2512.00502](#)
- VBS ssWW and WZ:
- VBS ZV semileptonic: [arXiv:2510.00118](#)
- $\gamma\gamma \rightarrow WW$ : [arXiv:2601.21574](#)
- WWZ: [PRL 135 \(2025\) 091802](#) (Run 2 + Run 3)

- WW: [JHEP 08 \(2025\) 142](#)
- WZ: [JHEP 11 \(2025\) 006](#)
- ZZ: [arXiv:2511.15569](#) (Run 2)
- Wg:
- ZZg: [arXiv:2602.17165](#) (Run 2)
- VBS semileptonic: [arxiv.2503.17461](#)
- WWg: [Phys. Lett. B 873 \(2026\) 140050](#)
- ZZg: [arXiv:2602.17165](#)








# Why multiboson & VBS?



comprehensive  
test of the electroweak sector at the LHC







# Searching for new physics

- High-energy tails provide strong sensitivity to dimension-8 EFT operators.
- Complementary strength across the operator basis between ATLAS and CMS






# Differential diboson

- Inclusive ZZ and ZZjj (VBS topology)
- Differential distributions in agreement with NLO
- Sensitivity to neutral TGC and QGC



# Neutral TGC sensitivity: $Z\gamma$ @ 13 and 13.6 TeV

- $Z(\ell\ell)\gamma$  @ 13.6 TeV ( $34.8 \text{ fb}^{-1}$ )
  - $\sigma_{\text{fid}} = 1.896 - 0.066 \text{ pb}$





# ZZ $\gamma$

This results in an observed (expected) significance of the ZZ $\gamma$  of  $4.4\sigma$  ( $4.4\sigma$ ).

$$p_T^\gamma > 20 \text{ GeV}; |\eta_\gamma| < 2.37; \Delta R(\ell, \gamma) > 0.4$$

$$\sigma_{ZZ\gamma} = 0.144_{-0.051}^{+0.064} (\text{stat.})_{-0.005}^{+0.007} (\text{syst.}) \text{ fb}$$




Consistent with SM prediction, Run 3 to reach observation

First observation of  $4\ell\gamma$  ( $5.1\sigma$ ) and evidence of ZZ $\gamma$  ( $3.7\sigma$ )

$$p_T^\gamma > 20 \text{ GeV}; |\eta_\gamma| < 2.4; \Delta R(\ell, \gamma) > 0.5$$



$$\sigma_{ZZ\gamma} = 0.06 \pm 0.03 \text{ fb}$$



CMS-PAS-SMP-24-014

# Rare EW interactions


- Define signal in events with  $e\mu$  final state and no other tracks in the event.
- Detection of such exclusive events is still quite challenging due to the interaction of other protons in the beam (pileup).





first observation in CMS:  
 $\sigma(\gamma\gamma \rightarrow W^+W^-) = 659 \pm 80 \text{ fb}$

# VBF gamma

- First observation of EW  $\gamma jj$  @13 TeV
- Genuine VBF topology: large  $m_{jj}$ , large  $\Delta\eta_{jj}$ , central photon
- Direct probe of the  $WW\gamma$  vertex in pure EWK process
- Significance  $> 5\sigma$ ;  $\sigma = 202 \pm 36$  fb
- Limits on the Warsaw basis coef.:  $c_W = [-0.11, 0.16]$ ;  $c_{HWB} = [-1.6, 1.5]$







Consistent with SM



# VBS ZV semileptonic and fully leptonic


- VBS ZV in  $\ell\nu qq, \ell\ell qq$  @13 TeV ( $138 \text{ fb}^{-1}$ ) from CMS
- Resolved + merged topologies
- DNN-based signal extraction
- Statistically limited ( $1.3\sigma$ )



Rare SM processes with multiple gauge vertices; sensitive to EFT and high-energy behavior


Multiboson LHC

## ATLAS: VVZ observation (Run 2)



Observed significance:  $6.4\sigma$  (VVZ); evidence for WWZ




## CMS: WWZ + ZH (H→WW) simultaneous fit (Run 2+3)



Overall  $\mu \approx 1.03$  (Run 2+3); evidence at 13.6 TeV (Run 3)


# Radiative triboson: $WW\gamma$

- $WW\gamma$  Observation @13 TeV with  $5.9\sigma$
- Measured cross section:  $\sigma_{\text{fid}} = 6.2 \pm 1.0 \text{ fb}$
- aQGC dim-8 limits




*Phys. Lett. B 873 (2026) 140050*

## ATLAS: $WW\gamma$ observation (Run 2)



Significance:  $5.9\sigma$ ; fiducial  $\sigma(e\mu\nu\nu\gamma)$  measured

## CMS: $WZ\gamma$ observation (Run 2)



Observed significance:  $5.4\sigma$ ;  $\sigma(\text{fid}) = 5.48 \pm 1.11 \text{ fb}$