

Searches for BSM single Higgs bosons at the LHC (ATLAS + CMS)



Suzanne GASCON-SHOTKIN
IP2I Lyon (IN2P3-CNRS)/Université Claude Bernard Lyon 1
on behalf of the ATLAS and CMS Collaborations

Rencontres de Moriond, Electroweak Interactions &
Unified Theories 2026

Hotel Planibel, LA THUILE (IT), March 21 2026






Introduction



PLB 716 (2012)



- We found one scalar boson in 2012....
- Why do we still look for others?
- Many **BSM theories** allow additional (pseudo) scalar bosons!

This talk: Five recent (<1year) or **NEW** searches for BSM single Higgs bosons from ATLAS and CMS

What is a 'BSM single Higgs boson'? Either singly-produced, or produced in association with H_{125} :

• 'Dark' Higgs H_D (CMS) or s (ATLAS) model: A 2-mediator (Z' , H_D/s) DM model capable of reproducing the observed relic density

• $H_D \rightarrow bb$ (50-150 GeV) **CMS-PAS-SUS-23-013**

• $Z'^* \rightarrow Z' + s \rightarrow \chi\chi + bb$ (30-150 GeV) **ATLAS PRL 134 (2025) 121801**

• $Z'^* \rightarrow Z' + s \rightarrow \chi\chi + H_{125} H_{125}$ (250-400 GeV) **ATLAS JHEP 09 (2025) 067**

• Additional scalars A, H in IDM: $Z'^* \rightarrow AH \rightarrow Z' HH \rightarrow HH\ell\ell$ ($m_H = \{60-180\}$, $m_A - m_H = \{20-100\}$ GeV) **CMS-PAS-NPS-25-002**

• Additional spin-0/spin-1 resonances: $X \rightarrow ZZ$ or $Z' \rightarrow ZH \rightarrow bb\tau\tau$ ($m_{X/Z'} = \{200 \text{ GeV} - 6 \text{ TeV}\}$) **CMS-PAS-B2G-25-004**

Bonus: **New!** Updated summaries from CMS/ATLAS!!

Conclusion and perspectives


Acknowledgements

Please see other talks involving BSM Higgs boson searches:

- Searches for low-mass resonances in ATLAS and CMS: C. Collard 18/03
- Searches for new physics with photons in ATLAS and CMS: A. Agapitos 19/03
- Searches for new physics at high object masses A. Malara 19/03
- Searches for other exotica in ATLAS and CMS C. Koraka 19/03
- Multiple-Higgs searches at ATLAS and CMS in ATLAS and CMS. V. Dao 21/03




'Dark' Higgs $H_D \rightarrow bb$ (50-150 GeV)





- DM Z' produced with H_D mixing w/ H_{125} (θ_h)
- Signature: Large $E_{t,miss}$ + $H_{125} \rightarrow bb$



- Major backgrounds: V +jets, $t\bar{t}$, single t
- SR, 7 CR defined in $\{U(\text{non-hadronic } p_T), m_{SD}\}$ phase space



- Max joint LH over SR + CR to model background/extract $\mu = \sigma/\sigma_{th}$
- 95% CL limits on μ in $\{m_{Z'}, m_\chi\}$ phase space: Exclude Z' masses between 2.5-4.5 TeV (depending on m_{HD})




'Dark' Higgs s from $Z'^* \rightarrow Z' + s \rightarrow \chi\chi + bb$ (30-150 GeV)




- 1- μ and 2 ℓ CR for V+jets modeling \rightarrow Simultaneous ML fit to m_{bb} (SR) or event yields (CR)

- Signature: merged ($m_s < 50$ GeV) or resolved ($m_s > 70$ GeV) bb system, m_{bb} binned in $E_{T,miss}$
- Backgrounds from tt and V+jets
- Access lower m_s with $E_{T,miss}$ trigger and dedicated merged-bjet (large R) tagger





- Most significant uncertainties (% of total): Data statistics (75-80%), large-R bjet tagger calibration (20-47%)

'Dark' Higgs s from $Z'^* \rightarrow Z' + s \rightarrow \chi\chi + bb$ (30-150 GeV)




- Exclusions in $\{m_s, m_{Z'}\}$ plane:
Scenario 1: Z' - χ (DM) coupling fixed (comparable to CMS analysis)

- Scenario 2: Z' - χ coupling floats to match the relic density
- Largest deviations for $m_s \sim 50, 130$ GeV
- $m_{Z'} < 4.5$ TeV excluded for $m_s = 75$ GeV




- Scenario 3: $m_s = 70$ GeV (max sensitivity)
- Exclusion in $\{m_\chi, m_{Z'}\}$ plane: $m_{Z'} < 4.8$ TeV




- $m_{Z'} < 3.4$ TeV excluded for $m_s = 70$ GeV

'Dark' Higgs s from $Z'^* \rightarrow Z' + s \rightarrow \chi\chi + H_{125} H_{125}$ (250-400 GeV)





- Exclusions in $\{m_s, m_{Z'}\}$ plane:
Scenario 1: Z' - χ (DM) coupling fixed



- Signature: 3- or 4-bjets + E_{miss} , bb system merged ($m_{\text{bjet}} > 100$ GeV) or resolved
- Backgrounds from tt and Z+jets
- Kinematic NN > 0.95
- Simultaneous LH fits in 8 SR m_{HH} bins + 3CR ($n_b, m_{\text{T}^b_{\text{min}}} + 1\text{CR} (Z \rightarrow \mu\mu)$)


- Most significant uncertainties (% of total): Data statistics (95%), single t modeling/ σ (18%), MC statistics (17%)



- Exclude $m_s = \{250-400 \text{ GeV}\}$ for $500 < m_{Z'} < 2.3 \text{ TeV}$






$Z^* \rightarrow AH \rightarrow Z^* HH \rightarrow HH \ell\ell$ ($m_H = \{60-180\}$, $m_A - m_H = \{20-100\}$ GeV) IDM Run 2 + Run 3 (2022)




- **IDM (Inert Doublet Model):** 2HDM w/unbroken Z_2 symmetry \rightarrow 4 additional scalars: H^\pm + bosophilic A, H (**stable** DM candidate)
- Signature: Di-e or di- μ pair + E_{tmiss}
- PNN: Kinematical, transverse mass and jet-Z balance variables

- Major backgrounds after PNN > 0.9 : Diboson, $t\bar{t}$
- Simultaneous binned Max LH fit over SR + 7 CR to PNN output or dilepton $p_T \rightarrow$ model background/extract μ






$Z^* \rightarrow AH \rightarrow Z^* HH \rightarrow HH \ell \ell$ ($m_H = \{60-180\}$, $m_A - m_H = \{20-100\}$ GeV) IDM
 Run 2 + Run 3 (2022)



- Largely statistics-limited, but most important systematic uncertainties on μ : MC statistics + jet calibration


- 95% CL observed limits on m_H and m_A in $\{ (m_A - m_H), m_H \}$ phase space: Exclude $m_H < 108$ GeV for $m_A - m_H = 78$ GeV, and $m_A - m_H = \{35, 90$ GeV for $m_H = 60$ GeV




- **First limits on m_H and m_A in IDM Model** obtained by a dedicated search w/collision data
- Significant extension of constraints from previous direct/indirect measurements and DM searches: lower E_{tmiss} requirements





$X \rightarrow ZZ$ or $Z' \rightarrow ZH \rightarrow bb\tau\tau$ ($m_{X/Z'} = \{200 \text{ GeV} - 6 \text{ TeV}\}$)



- $bb\tau\tau$** final state, 12 categories: $\{\tau_\mu\tau_h, \tau_e\tau_h, \tau_h\tau_h\} \times \{\text{bb}/\tau\tau \text{ system boosted/resolved}\} \rightarrow 87.6\%$ of possible $\tau\tau$ decays



- Primary backgrounds: $t\bar{t}$, DY, QCD




- Dedicated reconstruction for boosted $\tau\tau$ pairs ($m_X > 1 \text{ TeV}$): identification of subjets within large-radius jets to re-seed τ_h reconstruction
- Elliptical selection in $\{m_{bb} - m_{\tau\tau}\}$ plane, $\epsilon_{\text{sig}} > 90\%$
- Diagonal selection to separate $Z_{bb}H_{\tau\tau}$ and $Z_{\tau\tau}H_{bb}$

- Signal extraction: Binned Max LH fit to output distribution of PNN with DNN architecture





$X \rightarrow ZZ$ or $Z' \rightarrow ZH \rightarrow bb\tau\tau$ ($m_{X/Z'} = \{200 \text{ GeV} - 6 \text{ TeV}\}$)



- 95% CL observed limits on
 - $\sigma(X \rightarrow ZZ)$ between 300pb-24fb
 - $\sigma(Z' \rightarrow ZH)$ between 0.4pb-12fb

• Dominant systematic uncertainties: DY and ttbar theoretical predictions, MC statistics, b-tagging efficiencies

• First analysis targeting $ZZ, ZH \rightarrow bb\tau\tau$ final state







New Run 2 Summaries: Spin 0 $X \rightarrow VV$

- ggF production



- VBF production



- 95% CL observed limits, **new** analyses added





New Summaries: Spin 0, MSSM reinterpretations



- hMSSM: $h=H_{125}$, $M_S \sim 1\text{TeV}$, Higgs sector describable by $\{\tan\beta, m_A\}$, and h phenomenology by couplings to V, t, b : Include latest H_{125} combination

- M_h^{125} EFT scenario: Heavy superparticles \rightarrow production/ decay of MSSM Higgs bosons little affected by them, extends M_h^{125} scenario to low $\tan\beta$



- 95% CL observed limits for m_A versus $\tan\beta$





New Summaries: Spin 0, MSSM reinterpretations

New!

- hMSSM: $h=H_{125}$, $M_S \sim 1\text{TeV}$, Higgs sector describable by $\{\tan\beta, m_A\}$, and h phenomenology by couplings to V, t, b

https://cms-summary-searches.docs.cern.ch/mssm_higgs_summary/

<https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/PUBNOTES/ATL-PHYS-PUB-2025-042/> includes new $H^+ \rightarrow \tau\nu$ result





- 95% CL observed limits for m_A versus $\tan\beta$



New Summaries: Spin-0, SM + singlet reinterpretation



- 95% CL observed limits for $\sin\alpha$ versus m_H in the SM+ singlet Higgs extension



- $\tan\beta=0.1$

- Include latest H_{125} combination, several other $\tan\beta$ values available

- $\tan\beta=4$






Conclusions and Perspectives



- No evidence for additional single BSM Higgs bosons found so far by CMS and ATLAS.
- Presented 4 results from Full LHC Run 2 data set + 1 including early Run 3 data (2022)
- Run 3 is almost over....but not the data analysis!
- Hope for $\sim 350\text{fb}^{-1} \rightarrow \sim 50\%$ (≥ 2025) largely unexplored for the moment!
- HL-LHC starts ~ 2030 , expect 3ab^{-1}
- Overall, still plenty of discovery potential at LHC!

https://twiki.cern.ch/twiki/bin/view/CMSPublic/LumiPublicResults#Multi_year_summary_proton_proton



<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/LuminosityPublicResultsRun3>

Acknowledgements


A. Agapitos, F. Beaudette, R. Camacho Toro, C. Charlot, C. Collard, A. Cortes-Gonzalez, T. Cuisset, A. De Iorio, V. Dao, M. Donega, N. Fomin, P. Francavilla, B. Hodgkinson, J. Hogan, C. Koraka, A-M Magnan, B. Maier, A. Malara, S. Mukherjee, A. Oliveira, P. Padley, T. Potter, T. Robens, H. Saka, S. Sekmen, L. Soffi, M. Verducci, D. Winterbottom, M. Xiao, H. Yang, ...and the Moriond EW organization!!




Backup




'Dark' Higgs s from $Z'^* \rightarrow Z' + s \rightarrow \chi\chi + bb$ (30-150 GeV)



- 1- and 2 μ CR for V+jets modeling \rightarrow Simultaneous ML fit to m_{bb} (SR) or event yields (CR)



- Signature: merged ($m_s < 50$ GeV) or resolved ($m_s > 70$ GeV) bb system, m_{bb} binned in $E_{t_{miss}}$
- Backgrounds from tt and V+jets
- Access lower m_s with $E_{t_{miss}}$ trigger and dedicated merged-bjet (large R) tagger



- Most significant uncertainties (% of total): Data statistics (75-80%), large-R bjet tagger calibration (20-47%)