

Probing Flavorful EFTs at the LHC

Matheus Martines

matheus.martines@ijclab.in2p3.fr

IJCLab (Orsay)

Based on:

- [D. Bečirević, **MM**, S. Rosauero-Alcaraz, O. Sumensari, 2603.25837]
- [O.J.P. Éboli, L.P.S. Leal, **MM**, O. Sumensari, 2509.08437]

IRN, October 20

The quest for physics beyond the Standard Model

The **Standard Model** is an **EFT** at low-energies of a more fundamental theory that remains **unknown**.

The quest for physics beyond the Standard Model


The **Standard Model** is an **EFT** at low-energies of a more fundamental theory that remains **unknown**.

Several questions that **cannot** be answered within the Standard Model:

Hierarchy problem



Flavor puzzle



and many more ...

The quest for physics beyond the Standard Model


The **Standard Model** is an **EFT** at low-energies of a more fundamental theory that remains **unknown**.

Several questions that **cannot** be answered within the Standard Model:

Hierarchy problem



Flavor puzzle



and many more ...

Addressing these fundamental problems **requires** physics **beyond the Standard Model**.

Direct searches

No **direct** evidence of New Physics (NP).


Presence of a **mass gap** between the electroweak scale and the scale of NP.

ATLAS Heavy Particle Searches* - 95% CL Upper Exclusion Limits

Status: Spring 2026

ATLAS Preliminary

∫L dt = (3.6 – 195) fb⁻¹ √s = 13, 13.6 TeV



*Only a selection of the available mass limits on new states or phenomena is shown.

[†]Small-radius (large-radius) jets are denoted by the letter j (J).

Direct searches

No **direct** evidence of New Physics (NP).

Presence of a **mass gap** between the electroweak scale and the scale of NP.




Indirect searches for New Physics.

ATLAS Heavy Particle Searches* - 95% CL Upper Exclusion Limits

Status: Spring 2026

ATLAS Preliminary

∫L dt = (3.6 – 195) fb⁻¹ √s = 13, 13.6 TeV



*Only a selection of the available mass limits on new states or phenomena is shown.

[†]Small-radius (large-radius) jets are denoted by the letter j (J).

1 TeV


The EFT approach

All possible interactions made of **SM fields** and respecting the **SM gauge symmetries**.

Standard Model EFT (SMEFT)

$$\mathcal{L}_{\text{SMEFT}} = \mathcal{L}_{\text{SM}} + \frac{1}{\Lambda} \mathcal{L}^{(5)} + \frac{1}{\Lambda^2} \mathcal{L}^{(6)} + \frac{1}{\Lambda^3} \mathcal{L}^{(7)} + \frac{1}{\Lambda^4} \mathcal{L}^{(8)} + \dots$$

$$\mathcal{L}^{(d)} = \sum_i c_i Q_i^{(d)}$$




- ▶ *New Physics* scale Λ controls the expansion;
- ▶ *Lepton* and *baryon* number conservation implies that *leading BSM* effects start at *dimension six*.

The EFT approach

All possible interactions made of **SM fields** and respecting the **SM gauge symmetries**.

Standard Model EFT (SMEFT)


$$\mathcal{L}_{\text{SMEFT}} = \mathcal{L}_{\text{SM}} + \frac{1}{\Lambda} \mathcal{L}^{(5)} + \frac{1}{\Lambda^2} \mathcal{L}^{(6)} + \frac{1}{\Lambda^3} \mathcal{L}^{(7)} + \frac{1}{\Lambda^4} \mathcal{L}^{(8)} + \dots$$
$$\mathcal{L}^{(d)} = \sum_i c_i Q_i^{(d)}$$






- ▶ *New Physics* scale Λ controls the expansion;
- ▶ *Lepton* and *baryon* number conservation implies that *leading BSM* effects start at **dimension six**.

*Same class of **EFT operators** can affect processes at **different scales**.*
e.g. high-energy LHC and low-energy flavor processes.

Synergy between low- and high-energy observables





Synergy between low- and high-energy observables



Can the **LHC** provide **competitive** bounds to those from **flavor**? **Yes!**


D. Becirevic *et al.* [2012.09872], O. Sumensari *et al.* [2312.14070] A. Greljo *et al.* [2306.09401, 2212.10497] ...

High- p_T Drell-Yan tails




J. Blas *et al.* [1307.5068], S. Dawson *et al.* [1811.12260], M. Farina *et al.* [1609.08157], L. Allwicher *et al.* [2207.10714, 2412.14162], T. Corbett *et al.* [2503.19962], G. Hiller *et al.* [2502.12250], S. Descostes-Genon *et al.* [2303.07521], A. Angelescu *et al.* [2002.05684], and many more....

High- p_T Drell-Yan tails




J. Blas *et al.* [1307.5068], S. Dawson *et al.* [1811.12260], M. Farina *et al.* [1609.08157], L. Allwicher *et al.* [2207.10714, 2412.14162], T. Corbett *et al.* [2503.19962], G. Hiller *et al.* [2502.12250], S. Descostes-Genon *et al.* [2303.07521], A. Angelescu *et al.* [2002.05684], and many more....

High- p_T Drell-Yan tails



J. Blas *et al.* [1307.5068], S. Dawson *et al.* [1811.12260], M. Farina *et al.* [1609.08157], L. Allwicher *et al.* [2207.10714, 2412.14162], T. Corbett *et al.* [2503.19962], G. Hiller *et al.* [2502.12250], S. Descostes-Genon *et al.* [2303.07521], A. Angelescu *et al.* [2002.05684], and many more....


High- p_T Drell-Yan tails




J. Blas *et al.* [1307.5068], S. Dawson *et al.* [1811.12260], M. Farina *et al.* [1609.08157], L. Allwicher *et al.* [2207.10714, 2412.14162], T. Corbett *et al.* [2503.19962], G. Hiller *et al.* [2502.12250], S. Descostes-Genon *et al.* [2303.07521], A. Angelescu *et al.* [2002.05684], and many more....

High- p_T Drell-Yan tails

Example: constraints on transitions mediated by the **charged current**.



$$\psi^4 : \mathcal{A}^{(6)} \sim E^2 / \Lambda^2$$



Energy-enhanced effects in the high- p_T tails.

J. Blas *et al.* [1307.5068], S. Dawson *et al.* [1811.12260], M. Farina *et al.* [1609.08157], L. Allwicher *et al.* [2207.10714, 2412.14162], T. Corbett *et al.* [2503.19962], G. Hiller *et al.* [2502.12250], S. Descotes-Genon *et al.* [2303.07521], A. Angelescu *et al.* [2002.05684], and many more....

Low-energy EFT for charged current processes


$$\begin{aligned}\mathcal{L}_{\text{LEFT}} = & -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ & \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}\end{aligned}$$

- Responsible for leptonic and semileptonic meson decays.

Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}$$

- Responsible for leptonic and **semileptonic** meson decays.




$$i\mathcal{A} = i\mathcal{C}_\Gamma \langle K(k) | \bar{s} \Gamma c | D(p) \rangle \langle \nu \bar{\mu} | \bar{\nu}_L \Gamma \mu | 0 \rangle$$

Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}$$

- Responsible for leptonic and **semileptonic** meson decays.




→ Evaluated using perturbative methods.

Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Ri} d_{Rj}) (\bar{\ell}_R \nu_L) \right]$$

- Responsible for leptonic and **semileptonic** meson decays

$$\langle K(k) | \bar{s} \gamma^\mu c | D(p) \rangle = \left[(p + k)^\mu - \frac{m_D^2 - m_K^2}{q^2} q^\mu \right] f_+(q^2) \\ + \frac{m_D^2 - m_K^2}{q^2} q^\mu f_0(q^2),$$



$$i\mathcal{A} = i\mathcal{C}_\Gamma \langle K(k) | \bar{s} \Gamma c | D(p) \rangle \langle \nu \bar{\mu} | \bar{\nu}_L \Gamma \mu | 0 \rangle$$

$$q^2 = (p - k)^2 = (p_\ell + p_\nu)^2$$


- Evaluated using perturbative methods.
- Non-perturbative methods are required, e.g. Lattice QCD.

Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Lj}) (\bar{\ell}_R \gamma^\mu \nu_L) \right]$$

$$\langle K(k) | \bar{s} \gamma^\mu c | D(p) \rangle = \left[(p + k)^\mu - \frac{m_D^2 - m_K^2}{q^2} q^\mu \right] f_+(q^2) \\ + \frac{m_D^2 - m_K^2}{q^2} q^\mu f_0(q^2),$$

► Respo



$$\langle K(k) | \bar{s} \Gamma c | D(p) \rangle \langle \nu \bar{\mu} | \bar{\nu}_L \Gamma \mu | 0 \rangle$$


$$q^2 = (p - k)^2 = (p_\ell + p_\nu)^2$$

g. Lattice QCD.

Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}$$

- Responsible for leptonic and **semileptonic** meson decays.



$$i\mathcal{A} = i\mathcal{C}_\Gamma \langle K(k) | \bar{s} \Gamma c | D(p) \rangle \langle \nu \bar{\mu} | \bar{\nu}_L \Gamma \mu | 0 \rangle$$

$$q^2 = (p - k)^2 = (p_\ell + p_\nu)^2$$

- ➔ Evaluated using perturbative methods.
- ➔ Non-perturbative methods are required, e.g. Lattice QCD.
- ➔ Encodes information about the underlining microscopic theory.


$$g_S = g_{S_R} + g_{S_L}$$

$$g_V = g_{V_R} + g_{V_L}$$

Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}$$

- Responsible for leptonic and **semileptonic** meson decays.




$$\frac{d^2\Gamma}{dq^2 d\cos\theta} = a(q^2) + b(q^2)\cos\theta + c(q^2)\cos^2\theta$$

Depend on the Wilson coefficients and the form-factors.

$$g_S = g_{S_R} + g_{S_L}$$


$$g_V = g_{V_R} + g_{V_L}$$



Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}$$

- Responsible for leptonic and **semileptonic** meson decays.




$$\frac{d^2\Gamma}{dq^2 d\cos\theta} = \underline{a(q^2)} + b(q^2) \cos\theta + \underline{c(q^2) \cos^2\theta}$$

$$\frac{d\Gamma}{dq^2} \propto a_V(q^2) |1 + g_V|^2 + a_S(q^2) |g_S|^2 + a_{VS}(q^2) \text{Re}[(1 + g_V)g_S^*] + \dots$$

Constrains **vector** and **scalar** couplings.

$$g_S = g_{S_R} + g_{S_L}$$


$$g_V = g_{V_R} + g_{V_L}$$



Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}$$

- Responsible for leptonic and **semileptonic** meson decays.




$$\frac{d^2\Gamma}{dq^2 d\cos\theta} = a(q^2) + \underline{b(q^2) \cos\theta} + c(q^2) \cos^2\theta$$

$$\frac{d\langle A_{\text{fb}} \rangle}{dq^2} \propto b_V(q^2) |1 + g_V|^2 + b_{VS}(q^2) \text{Re}[(1 + g_V)g_S^*] + \dots$$


$$g_S = g_{S_R} + g_{S_L}$$

$$g_V = g_{V_R} + g_{V_L}$$



Constrains the **real part** of the scalar coupling. [D. Bečirević, F. Jaffredo, A. Peñuelas, O. Sumensari, 2012.09872]

Low-energy EFT for charged current processes




Constrains **real part** of the scalar coupling.

Recently measured by BESIII Collaboration



Testing solutions at the LHC

A **mild tension** with the SM prediction is present in the BESIII differential decay width measurements of the $D \rightarrow K\mu\nu$ channels. [2601.21196, 2601.21185]



Testing solutions at the LHC

A **mild tension** with the SM prediction is present in the BESIII differential decay width measurements of the $D \rightarrow K\mu\nu$ channels. [2601.21196, 2601.21185]





Complex scalar coupling can alleviate the tension.

$$\mathcal{O}_S = (\bar{c}s)(\bar{\mu}_R\nu_L)$$

[D. Becirevic, **MM**, S. Rosauero-Alcaraz, O. Sumensari, 2603.25837]

Testing solutions at the LHC

A **mild tension** with the SM prediction is present in the BESIII differential decay width measurements of the $D \rightarrow K\mu\nu$ channels. [2601.21196, 2601.21185]



Complex scalar coupling can alleviate the tension.



$$\mathcal{O}_S = (\bar{c}s)(\bar{\mu}_R\nu_L)$$

The proposed solution is **not** consistent with **Drell-Yan limits**.

[D. Becirevic, **MM**, S. Rosauero-Alcaraz, O. Sumensari, 2603.25837]

Testing solutions at the LHC

A **mild tension** with the SM prediction is present in the BESIII differential decay width measurements of the $D \rightarrow K\mu\nu$ channels. [2601.21196, 2601.21185]



Complex scalar coupling can alleviate the tension.

$$\mathcal{O}_S = (\bar{c}s)(\bar{\mu}_R\nu_L)$$



The proposed solution is **not** consistent with **Drell-Yan limits**.

*High-energy observables provide a **consistency check** on solutions to **low-energy flavor tensions**.*

[D. Becirevic, **MM**, S. Rosauero-Alcaraz, O. Sumensari, 2603.25837]

Testing solutions at the LHC

A **mild tension** with the SM prediction is present in the BESIII differential decay width measurements of the $D \rightarrow K\mu\nu$ channels. [2601.21196, 2601.21185]



Complex scalar coupling can alleviate the tension.

$$\mathcal{O}_S = (\bar{c}s)(\bar{\mu}_R\nu_L)$$

The proposed solution is **not** consistent with **Drell-Yan limits**.

*High-energy observables provide a **consistency check** on solutions to **low-energy flavor tensions**.*

Not all couplings affecting low-energy **semileptonic transitions** mediated by **charged currents** can be constrained by **Drell-Yan** data.

Low-energy EFT for charged current processes

$$\begin{aligned} \mathcal{L}_{\text{LEFT}} = & -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ & \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.} \end{aligned}$$

Matching with SMEFT @ d=6:

$$g_{V_L}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^2 H^2 D}^{ij} - \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$

$$g_{V_R}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^2 H^2 D}^{ij},$$

$$g_{S_L}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$

$$g_{S_R}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$

$$g_T^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell}.$$

Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}$$

Matching with SMEFT @ d=6:

$$g_{V_L}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^2 H^2 D}^{ij} - \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$

$$g_{V_R}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^2 H^2 D}^{ij},$$


$$g_{S_L}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$

$$g_{S_R}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$

$$g_T^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell}.$$

Well constrained by $pp \rightarrow \ell \nu$ data

$$\psi^4 : \mathcal{A}^{(6)} \sim E^2 / \Lambda^2$$



J. Fuentes-Martín et al. [2003.12421], M. Farina et al. [1609.08157],
A. Greljo et al. [2212.10497], A. Angelescu et al. [2002.05684], ...

Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}$$

Matching with SMEFT @ d=6:

$$g_{V_L}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^2 H^2 D}^{ij} - \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$

$$g_{V_R}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^2 H^2 D}^{ij},$$

$$g_{S_L}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$


$$g_{S_R}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$

$$g_T^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell}.$$

Poorly constrained by $pp \rightarrow \ell \nu$ data.

$$\psi^2 H^2 D : \mathcal{M}^{(6)}(u\bar{d} \rightarrow \ell^+ \nu) \sim v^2 / \Lambda^2$$

Only generated by a $\psi^2 H^2 D$ operator @ d=6.



Low-energy EFT for charged current processes

$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}$$

Matching with SMEFT @ d=6:

$$g_{V_L}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^2 H^2 D}^{ij} - \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$

$$g_{V_R}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^2 H^2 D}^{ij},$$

$$g_{S_L}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$


$$g_{S_R}^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell},$$

$$g_T^{ij(\ell)} \propto \frac{v^2}{\Lambda^2} \mathcal{C}_{\psi^4}^{ij\ell\ell}.$$

Poorly constrained by $pp \rightarrow \ell \nu$ data.

$$\psi^2 H^2 D : \mathcal{M}^{(6)}(u\bar{d} \rightarrow \ell^+ \nu) \sim v^2/\Lambda^2$$


Only generated by a $\psi^2 H^2 D$ operator @ d=6.



The operators in $\psi^2 H^2 D$ class can be **energy-enhanced** in **Higgs** production channels.

Example: Right-Handed Charged Currents at the LHC

$$\mathcal{O}_{Hud} = (\tilde{H}^\dagger i D_\mu H)(\bar{u}_R \gamma^\mu d_R) + \text{h.c.}$$



$$\psi^2 H^2 D : \mathcal{M}^{(6)}(u\bar{d} \rightarrow \ell^+ \nu) \sim v^2 / \Lambda^2$$




Poorly constrained by DY.


Example: Right-Handed Charged Currents at the LHC

$$\mathcal{O}_{Hud} = (\tilde{H}^\dagger i D_\mu H)(\bar{u}_R \gamma^\mu d_R) + \text{h.c.}$$

Energy-enhancement effects



$$\mathcal{M}(u_R^i \bar{d}_R^j \rightarrow W_0 h) = -\frac{i}{\sqrt{2}} \frac{E^2 \sin \theta}{\Lambda^2} \mathcal{C}_{Hud}^*{}_{ij}$$



$$\propto \frac{[\mathcal{C}_{Hud}]_{ij}}{\Lambda^2} P_R$$

$$\psi^2 H^2 D : \mathcal{M}^{(6)}(u\bar{d} \rightarrow \ell^+ \nu) \sim v^2/\Lambda^2$$




Poorly constrained by DY.

Example: Right-Handed Charged Currents at the LHC

$$\mathcal{O}_{Hud} = (\tilde{H}^\dagger i D_\mu H)(\bar{u}_R \gamma^\mu d_R) + \text{h.c.}$$


Energy-enhancement effects




$$\propto \frac{[\mathcal{C}_{Hud}]_{ij}}{\Lambda^2} P_R$$

$$\psi^2 H^2 D : \mathcal{M}^{(6)}(u\bar{d} \rightarrow \ell^+ \nu) \sim v^2 / \Lambda^2$$

Poorly constrained by DY.



$$\mathcal{M}(u_R^i \bar{d}_R^j \rightarrow W_0 h) = -\frac{i}{\sqrt{2}} \frac{E^2 \sin \theta}{\Lambda^2} \mathcal{C}_{Hud}^*{}_{ij}$$




Modifications to the SM couplings **spoil** the cancellations.

$$i\mathcal{M}_{\text{SM}} \simeq i\sqrt{2} V_{ij}^* \sin \theta \left[c_W^2 - \left(T^{(u)} - s_W^2 Q^{(u)} \right) + \left(T^{(d)} - s_W^2 Q^{(d)} \right) \right] \frac{E^2}{v^2} = 0$$

Example: Right-Handed Charged Currents at the LHC

$$\mathcal{O}_{Hud} = (\tilde{H}^\dagger i D_\mu H)(\bar{u}_R \gamma^\mu d_R) + \text{h.c.}$$


Energy-enhancement effects




$$\propto \frac{[\mathcal{C}_{Hud}]_{ij}}{\Lambda^2} P_R$$

$$\psi^2 H^2 D : \mathcal{M}^{(6)}(u\bar{d} \rightarrow \ell^+ \nu) \sim v^2 / \Lambda^2$$

Poorly constrained by DY.



$$\mathcal{M}(u_R^i \bar{d}_R^j \rightarrow W_0 h) = -\frac{i}{\sqrt{2}} \frac{E^2 \sin \theta}{\Lambda^2} \mathcal{C}_{Hud}^*{}_{ij}$$




Modifications to the SM couplings **spoil** the cancellations.

$$\mathcal{M}(u_R^i \bar{d}_R^j \rightarrow W_0 Z_0) = -\frac{i}{\sqrt{2}} \frac{E^2 \sin \theta}{\Lambda^2} \mathcal{C}_{Hud}^*{}_{ij}$$

Example: Right-Handed Charged Currents at the LHC

$$\mathcal{O}_{Hud} = (\tilde{H}^\dagger i D_\mu H)(\bar{u}_R \gamma^\mu d_R) + \text{h.c.}$$


Energy-enhancement effects




$$\propto \frac{[\mathcal{C}_{Hud}]_{ij}}{\Lambda^2} P_R$$

$$\psi^2 H^2 D : \mathcal{M}^{(6)}(u\bar{d} \rightarrow \ell^+ \nu) \sim v^2 / \Lambda^2$$

Poorly constrained by DY.



$$\mathcal{M}(u_R^i \bar{d}_R^j \rightarrow W_0 h) = -\frac{i}{\sqrt{2}} \frac{E^2 \sin \theta}{\Lambda^2} \mathcal{C}_{Hud}^*_{ij}$$



Modifications to the SM couplings **spoil** the cancellations.

$$\mathcal{M}(u_R^i \bar{d}_R^j \rightarrow W_0 Z_0) = -\frac{i}{\sqrt{2}} \frac{E^2 \sin \theta}{\Lambda^2} \mathcal{C}_{Hud}^*_{ij}$$

These amplitudes are related by the **Goldstone Boson Equivalence Theorem.**

Goldstone Boson Equivalence Theorem

In the limit $E \gg m_W, m_Z$,

$$\mathcal{M}(q_i \bar{q}_j \rightarrow V_0 V_0') \propto \mathcal{M}(q_i \bar{q}_j \rightarrow \phi \phi')$$

$V_0 \rightarrow$ longitudinal polarization

$\phi \rightarrow$ Goldstone boson

B. W. Lee *et al.*, *Phys. Rev. D* 16, 1519, G. J. Gounaris *et al.*, *Phys. Rev. D* 34, 3257,
M. Chanowitz, *et al.*, *Phys. Rev. D* 36, 1490, *Nucl.Phys.B* 261 (1985) 379-431, A.
Wulzer [1309.6055]

See also R. Franceschini *et al.* [1712.01310]

Goldstone Boson Equivalence Theorem

In the limit $E \gg m_W, m_Z$,

$$\mathcal{M}(q_i \bar{q}_j \rightarrow V_0 V_0') \propto \mathcal{M}(q_i \bar{q}_j \rightarrow \phi \phi')$$

$V_0 \rightarrow$ longitudinal polarization

$\phi \rightarrow$ Goldstone boson

B. W. Lee *et al.*, *Phys. Rev. D* 16, 1519, G. J. Gounaris *et al.*, *Phys. Rev. D* 34, 3257,
M. Chanowitz, *et al.*, *Phys. Rev. D* 36, 1490, *Nucl.Phys.B* 261 (1985) 379-431, A.
Wulzer [1309.6055]

$$H = \begin{pmatrix} -iw^+ \\ \frac{1}{\sqrt{2}}(v + h + iz) \end{pmatrix}$$

$\rightarrow$

$$\mathcal{O}_{Hud} = (\tilde{H}^\dagger i D_\mu H) (\bar{u}_R \gamma^\mu d_R) + \text{h.c.}$$

See also R. Franceschini *et al.* [1712.01310]

Goldstone Boson Equivalence Theorem

In the limit $E \gg m_W, m_Z$,

$$\mathcal{M}(q_i \bar{q}_j \rightarrow V_0 V_0') \propto \mathcal{M}(q_i \bar{q}_j \rightarrow \phi \phi')$$

$V_0 \rightarrow$ longitudinal polarization

$\phi \rightarrow$ Goldstone boson

B. W. Lee *et al.*, *Phys. Rev. D* 16, 1519, G. J. Gounaris *et al.*, *Phys. Rev. D* 34, 3257,
M. Chanowitz, *et al.*, *Phys. Rev. D* 36, 1490, *Nucl.Phys.B* 261 (1985) 379-431, A.
Wulzer [1309.6055]

$$H = \begin{pmatrix} -iw^+ \\ \frac{1}{\sqrt{2}}(v + h + iz) \end{pmatrix}$$


$\rightarrow$

$$\mathcal{O}_{Hud} = (\tilde{H}^\dagger i D_\mu H) (\bar{u}_R \gamma^\mu d_R) + \text{h.c.}$$

$\downarrow$

$$\mathcal{O}_{Hud} = \sqrt{2} (h + iz) (\partial_\mu w^+) (\bar{u}_i \gamma^\mu P_R d_j) + \dots$$

$\nearrow$



See also R. Franceschini *et al.* [1712.01310]

Goldstone Boson Equivalence Theorem

In the limit $E \gg m_W, m_Z$,

$$\mathcal{M}(q_i \bar{q}_j \rightarrow V_0 V_0') \propto \mathcal{M}(q_i \bar{q}_j \rightarrow \phi \phi')$$

$V_0 \rightarrow$ longitudinal polarization

$\phi \rightarrow$ Goldstone boson

B. W. Lee *et al.*, *Phys. Rev. D* 16, 1519, G. J. Gounaris *et al.*, *Phys. Rev. D* 34, 3257,
M. Chanowitz, *et al.*, *Phys. Rev. D* 36, 1490, *Nucl.Phys.B* 261 (1985) 379-431, A.
Wulzer [1309.6055]

$$H = \begin{pmatrix} -iw^+ \\ \frac{1}{\sqrt{2}}(v + h + iz) \end{pmatrix}$$


$\rightarrow$

$$\mathcal{O}_{Hud} = (\tilde{H}^\dagger i D_\mu H) (\bar{u}_R \gamma^\mu d_R) + \text{h.c.}$$

$\downarrow$

$$\mathcal{O}_{Hud} = \sqrt{2} (h + iz) (\partial_\mu w^+) (\bar{u}_i \gamma^\mu P_R d_j) + \dots$$

$\nearrow$



$\rightarrow$ At energies $E^2 \gg m_W^2$, these amplitudes are related by $SU(2)_L$ gauge invariance.

See also R. Franceschini *et al.* [1712.01310]

Goldstone Boson Equivalence Theorem

In the limit $E \gg m_W, m_Z$,

$$\mathcal{M}(q_i \bar{q}_j \rightarrow V_0 V_0') \propto \mathcal{M}(q_i \bar{q}_j \rightarrow \phi \phi')$$

$V_0 \rightarrow$ longitudinal polarization

$\phi \rightarrow$ Goldstone boson

B. W. Lee *et al.*, *Phys. Rev. D* 16, 1519, G. J. Gounaris *et al.*, *Phys. Rev. D* 34, 3257,
M. Chanowitz, *et al.*, *Phys. Rev. D* 36, 1490, *Nucl.Phys.B* 261 (1985) 379-431, A.
Wulzer [1309.6055]

$$H = \begin{pmatrix} -iw^+ \\ \frac{1}{\sqrt{2}}(v + h + iz) \end{pmatrix}$$


$\rightarrow$

$$\mathcal{O}_{Hud} = (\tilde{H}^\dagger i D_\mu H) (\bar{u}_R \gamma^\mu d_R) + \text{h.c.}$$

$\downarrow$

$$\mathcal{O}_{Hud} = \sqrt{2} (h + iz) (\partial_\mu w^+) (\bar{u}_i \gamma^\mu P_R d_j) + \dots$$

$\nearrow$



- $\rightarrow$ At energies $E^2 \gg m_W^2$, these amplitudes are related by $SU(2)_L$ **gauge invariance**.
- $\rightarrow$ Characteristic of all dim-6 operators modifying **neutral** & **charged** currents.

See also R. Franceschini *et al.* [1712.01310]

Dimension-six Higgs current operators

Only Zh & WW

Only Wh & WZ

Vh and WV

$\mathcal{O}_{\psi^2 H^2 D}$	Operator
$\mathcal{O}_{Hq}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_i \gamma^\mu q_j)$
$\mathcal{O}_{Hq}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H) (\bar{q}_i \gamma^\mu \tau^I q_j)$
$\mathcal{O}_{Hu}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{u}_i \gamma^\mu u_j)$
$\mathcal{O}_{Hd}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{d}_i \gamma^\mu d_j)$
$\mathcal{O}_{Hud}$	$(\tilde{H}^\dagger i D_\mu H) (\bar{u}_i \gamma^\mu d_j) + \text{h.c.}$


Channel	Distribution	Collaboration	N_{obs}	$\mathcal{L}$
$pp \rightarrow WW$	$\frac{d\sigma}{dp_T^{\ell^{\text{lead}}}}$	ATLAS	14	36.1 fb ⁻¹ [42]
	$\frac{dN_{\text{ev}}}{dm_{e\mu}}$	CMS	11	35.9 fb ⁻¹ [43]
$pp \rightarrow WZ$	$\frac{d\sigma}{dm_T^{WZ}}$	ATLAS	12	140 fb ⁻¹ [44]
	$\frac{1}{\sigma} \frac{d\sigma}{dm_{WZ}}$	CMS	5	137 fb ⁻¹ [45]
$pp \rightarrow Zh$	$\frac{d\sigma}{dp_T^Z}$	ATLAS	5	140 fb ⁻¹ [30]
		CMS	3	138 fb ⁻¹ [31]
$pp \rightarrow Wh$	$\frac{d\sigma}{dp_T^W}$	ATLAS	5	140 fb ⁻¹ [30]
		CMS	3	138 fb ⁻¹ [31]



Kinematic variables related to the **center-of-mass energy**.


Dimension-six Higgs current operators

Only Zh & WW	Only Wh & WZ	Vh and WV
$\mathcal{O}_{\psi^2 H^2 D}$	Operator	
$\mathcal{O}_{Hq}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_i \gamma^\mu q_j)$	
$\mathcal{O}_{Hq}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H) (\bar{q}_i \gamma^\mu \tau^I q_j)$	
$\mathcal{O}_{Hu}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{u}_i \gamma^\mu u_j)$	
$\mathcal{O}_{Hd}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{d}_i \gamma^\mu d_j)$	
$\mathcal{O}_{Hud}$	$(\tilde{H}^\dagger i D_\mu H) (\bar{u}_i \gamma^\mu d_j) + \text{h.c.}$	





Higgs differential distributions in the format of **Simplified Template Cross-Sections (STXS)** are available. D. De Florian *et al.* [1610.07822] 14

Single parameter limits




$$\sigma = \sigma_{\text{SM}} + \frac{1}{\Lambda^2} \sigma_{\text{Int}} + \frac{1}{\Lambda^4} \sigma_{\text{NP}^2}$$





**Considering a single Wilson coefficient at the time.*

Single parameter limits





Heavier initial states lead to **weaker** constraints due to **PDF** suppression.

Limits from Vh are a factor of 2 **stronger** than WV when considering Run 2 data.



**Considering a single Wilson coefficient at the time.*

Differential distributions





In diboson production channels, **transverse** polarizations **dominate** compared to **longitudinal** modes.

R. Franceschini *et al.* [1712.01310]

Differential distributions

Examples where **Higgs** and **DY** data are important.



In diboson production channels, **transverse** polarizations **dominate** compared to **longitudinal** modes.





R. Franceschini *et al.* [1712.01310]

LHC & Flavor I: $c \rightarrow s\mu\nu$

Assuming all coefficients to be **real**.

DY & VH data allows to constrain **all** parameter space.

All scenarios are **consistent** with the **SM** within 2σ .



[D. Becirevic, **MM**, S. Rosauo-Alcaraz, O. Sumensari, 2603.25837]




LHC & Flavor I: $c \rightarrow s\mu\nu$

Assuming all coefficients to be **real**.

DY & VH data allows to constrain **all** parameter space.

All scenarios are **consistent** with the **SM** within 2σ .

HL-LHC will further improve the limits on charged-current interactions.







[D. Becirevic, **MM**, S. Rosauro-Alcaraz, O. Sumensari, 2603.25837]

LHC & Flavor II: First-row CKM unitarity

$$\Delta_{\text{CKM}}^{\text{global}} \equiv |V_{ud}|_{\text{global}}^2 + |V_{us}|_{\text{global}}^2 + |V_{ub}|^2 - 1 = -0.001\,51(53) \quad \sim 3\sigma \text{ tension with CKM unitarity.}$$


A. Crivellin *et al.* [2212.06862], [2008.01113], B. Belfatto *et al.* [1906.02714], Y. Grossman *et al.* [1911.07821], M. Kirk [2008.0113], A. K. Alok *et al.* [2108.05614], and many more...




LHC & Flavor II: First-row CKM unitarity

$$\Delta_{\text{CKM}}^{\text{global}} \equiv |V_{ud}|_{\text{global}}^2 + |V_{us}|_{\text{global}}^2 + |V_{ub}|^2 - 1 = -0.001\,51(53) \quad \sim 3\sigma \text{ tension with CKM unitarity.}$$


A. Crivellin *et al.* [2212.06862], [2008.01113], B. Belfatto *et al.* [1906.02714], Y. Grossman *et al.* [1911.07821], M. Kirk [2008.0113], A. K. Alok *et al.* [2108.05614], and many more...



Assuming the discrepancy arises from NP, **RH CC** could offer a viable explanation. V. Cirigliano *et al.* [2208.11707]




LHC & Flavor II: First-row CKM unitarity



LHC and HL-LHC limits are dominated by Wh production.

HL-LHC has the potential to provide constraints that are **competitive** with those from **flavor observables**.



$$\mathcal{M}(u_R^i \bar{d}_R^j \rightarrow W_0 h) = -\frac{i}{\sqrt{2}} \frac{E^2 \sin \theta}{\Lambda^2} C_{Hud}^*{}_{ij}$$

[O. J. P. Éboli, L. P. S. Leal, **MM**, O. Sumensari, 2509.08437]

Summary

LHC data offer **complementary constraints** to low-energy processes, benefiting from **energy-enhanced** effects in the high- p_T tails.

Diboson (**VV**) and Higgs-associated (**VH**) production processes can **probe** Wilson coefficients that are **poorly constrained** by Drell–Yan data.

The **combination** of **DY** and **VH** production allows to constrain all couplings affecting **charged-current** interactions.

Soon to be implemented in **HighPT Mathematica** package. [github.com/HighPT/HighPT]

*The **HL-LHC** could help clarify **flavor anomalies**, such as the Cabibbo angle anomaly, and provide meaningful constraints on several transitions.*

Summary

LHC data offer **complementary constraints** to low-energy processes, benefiting from **energy-enhanced** effects in the high- p_T tails.

Diboson (**VV**) and Higgs-associated (**VH**) production processes can **probe** Wilson coefficients that are **poorly constrained** by Drell–Yan data.

The **combination** of **DY** and **VH** production allows to constrain all couplings affecting **charged-current** interactions.

Soon to be implemented in **HighPT Mathematica** package. [github.com/HighPT/HighPT]

*The **HL-LHC** could help clarify **flavor anomalies**, such as the Cabibbo angle anomaly, and provide meaningful constraints on several transitions.*

Thank you!

Backup

Interpreting the results on exclusive $c \rightarrow s\mu\nu$ modes

Illustration II: BESIII measurements on $c \rightarrow s\mu\nu$ modes

BESIII measurements on total branching ratio and forward-backward asymmetry are **consistent** with the SM prediction. [2601.21196, 2601.21185]

Mild tension present in the q^2 distribution.









Illustration II: BESIII measurements on $c \rightarrow s\mu\nu$ modes

BESIII measurements on total branching ratio and forward-backward asymmetry are **consistent** with the SM prediction. [2601.21196, 2601.21185]

Mild tension present in the q^2 distribution.





Tension is coming from **large q^2 bins in $\Delta\Gamma$ distribution.**

Illustration II: BESIII measurements on $c \rightarrow s\mu\nu$ modes

BESIII measurements on total branching ratio and forward-backward asymmetry are **consistent** with the SM prediction. [2601.21196, 2601.21185]

Mild tension present in the q^2 distribution.


Which **EFT operators** could alleviate this tension?



Tension is coming from **large q^2 bins in $\Delta\Gamma$ distribution.**

Complex scalar coupling

Solution proposed by BESIII [2601.21185]




[D. Becirevic,, **MM**, S. Rosauero-Alcaraz, O. Sumensari, 2603.25837]


Complex scalar coupling can alleviate the tension.

The tension is present regardless of the **form-factor parametrization**.

Tension is coming from **large q^2 bins**.



Complex scalar coupling




Solution proposed by BESIII Collaboration [2601.21185]

Complex scalar coupling can alleviate the tension.
The tension is present regardless of the **form-factor parametrization**.

[D. Becirevic, **MM**, S. Rosauero-Alcaraz, O. Sumensari, 2603.25837]

Complex scalar coupling




Solution proposed by BESIII Collaboration [2601.21185]

Complex scalar coupling can alleviate the tension.
The tension is present regardless of the **form-factor parametrization**.

[D. Becirevic, **MM**, S. Rosauro-Alcaraz, O. Sumensari, 2603.25837]

Is the proposed solution consistent with high- p_T bounds?

Complex scalar coupling




A_{FB} is **linearly** sensitive to the real part of the **scalar** coupling.

D. Becirevic et al. [2012.09872], S. Fajfer et al. [1502.07488]

→ A_{FB} and $\Delta\Gamma$ are complementary to each other.

[D. Becirevic, **MM**, S. Rosauero-Alcaraz, O. Sumensari, 2603.25837]

Complex scalar coupling



A_{FB} is **linearly** sensitive to the real part of the **scalar** coupling.


D. Becirevic et al. [2012.09872], S. Fajfer et al. [1502.07488]

➔ A_{FB} and $\Delta\Gamma$ are complementary to each other.

➔ **High- p_T** bounds provides stringent constraints, **excluding** the 1σ preferred region.

[D. Becirevic, **MM**, S. Rosauero-Alcaraz, O. Sumensari, 2603.25837]

Complex scalar coupling



A_{FB} is **linearly** sensitive to the real part of the **scalar** coupling.


D. Becirevic et al. [2012.09872], S. Fajfer et al. [1502.07488]

- A_{FB} and $\Delta\Gamma$ are complementary to each other.
- **High- p_T** bounds provides stringent constraints, **excluding** the 1σ preferred region.

Which other scenarios are possible?

[D. Becirevic, **MM**, S. Rosauero-Alcaraz, O. Sumensari, 2603.25837]

Real couplings



See also C. Bolognani, M. Reboud, D. Van Dyk, K. Keri Vos [2407.06145]

Low-energy EFT for charged current processes


$$\mathcal{L}_{\text{LEFT}} = -2\sqrt{2}G_F V_{ij} \left[(1 + g_{V_L}^{ij(\ell)}) (\bar{u}_{Li} \gamma_\mu d_{Lj}) (\bar{\ell}_L \gamma^\mu \nu_L) + g_{V_R}^{ij(\ell)} (\bar{u}_{Ri} \gamma_\mu d_{Rj}) (\bar{\ell}_L \gamma^\mu \nu_L) \right. \\ \left. + g_{S_L}^{ij(\ell)} (\bar{u}_{Ri} d_{Lj}) (\bar{\ell}_R \nu_L) + g_{S_R}^{ij(\ell)} (\bar{u}_{Li} d_{Rj}) (\bar{\ell}_R \nu_L) + g_T^{ij(\ell)} (\bar{u}_{Ri} \sigma_{\mu\nu} d_{Lj}) (\bar{\ell}_R \sigma^{\mu\nu} \nu_L) \right] + \text{h.c.}$$

- Responsible for **leptonic** and semileptonic meson decays.

$$g_A \equiv g_{V_R} - g_{V_L}$$

$$g_S \equiv g_{S_R} - g_{S_L}$$

decay constant




$$\mathcal{B}(D_s \rightarrow \mu\nu) = \tau_{D_s} \frac{G_F^2 |V_{cs}|^2 f_{D_s}^2 m_{D_s} m_\mu^2}{8\pi} \left| 1 - g_A + g_P \frac{m_{D_s}^2}{m_\mu(m_c + m_s)} \right|^2$$

Helicity suppressed

Constraints on **axial** and **pseudoscalar** couplings.

Probing Flavorful EFTs at the LHC

Confronting LHC with EWPO







Considering a **single non vanishing** coefficient at a time.



Confronting LHC with EWPO

Vh production **breaks** the blind-directions present in the **EWPO** involving the **first generation of quarks**.

Forward-backward asymmetry limits taken from V. Bresó-Pla et al.[2103.12074].




Limits on VLQs



$$\mathcal{L}_{\text{VLQ}} \supset -\xi_{fi}^{(u)} \bar{Q}_f \tilde{H} u_i - \xi_{fi}^{(d)} \bar{Q}_f H d_i + \text{h.c.}$$

Diboson production



SM prediction for W_0W_0 production

$$\mathcal{M}(u_i\bar{u}_j \rightarrow W_0W_0) \stackrel{\text{SM}}{=} i\hat{S} \frac{e^2 \sin \theta}{2m_W^2} \left[\textcolor{green}{Q_q \delta_{ij}} + \textcolor{brown}{\frac{1}{s_W^2} (T_q^3 - s_W^2 Q_q) \delta_{ij}} - \textcolor{blue}{\frac{T_q^3}{s_W^2} (V^\dagger \cdot V)_{ij}} \right] + \mathcal{O}(\hat{s}^0),$$

Center of mass energy


t -channel

s -channel Z

s -channel γ

C. Grojean, M. Montull, M. Riembau [1810.05149]
 R. Franceschini, G. Panico, A. Pomarol, F. Riva, A. Wulzer [1712.01310]
 T. Corbett, O. J. P. Éboli, M. C. Gonzalez-Garcia [1705.09294]

Diboson production



SM prediction for $W_0 W_0$ production

$$\mathcal{M}(u_i \bar{u}_j \rightarrow W_0 W_0) \stackrel{\text{SM}}{=} i\hat{S} \frac{e^2 \sin \theta}{2m_W^2} \left[Q_q \delta_{ij} + \frac{1}{s_W^2} (T_q^3 - s_W^2 Q_q) \delta_{ij} - \frac{T_q^3}{s_W^2} (V^\dagger \cdot V)_{ij} \right] + \mathcal{O}(\hat{s}^0),$$

t -channel
s-channel Z
s-channel γ

Center of mass energy

CKM is unitary


Modifications to the Z and W couplings can **spoil** the SM cancellation.

C. Grojean, M. Montull, M. Riembau [1810.05149]

R. Franceschini, G. Panico, A. Pomarol, F. Riva, A. Wulzer [1712.01310]

T. Corbett, O. J. P. Éboli, M. C. Gonzalez-Garcia [1705.09294]

Diboson production



SM prediction for $W_0 W_0$ production

$$\mathcal{M}(u_i \bar{u}_j \rightarrow W_0 W_0) \stackrel{\text{SM}}{=} i\hat{S} \frac{e^2 \sin \theta}{2m_W^2} \left[Q_q \delta_{ij} + \frac{1}{s_W^2} (T_q^3 - s_W^2 Q_q) \delta_{ij} - \frac{T_q^3}{s_W^2} (V^\dagger \cdot V)_{ij} \right] + \mathcal{O}(\hat{s}^0),$$

t -channel
 s -channel Z
 s -channel γ

Center of mass energy

CKM is unitary

Modifications to the Z and W couplings can **spoil** the SM cancellation.

At high-energies, $\mathcal{M}(qq' \rightarrow W_0 V_0) \sim \mathcal{M}(qq' \rightarrow h V_0') \longrightarrow$ **Goldstone Boson Equivalence Theorem**

B. W. Lee *et al.*, *Phys. Rev. D* 16, 1519, G. J. Gounaris *et al.*, *Phys. Rev. D* 34, 3257,
M. Chanowitz, *et al.*, *Phys. Rev. D* 36, 1490, *Nucl.Phys.B* 261 (1985) 379-431, A.
Wulzer [1309.6055]

High-energy amplitudes

At dimension $d = 6$, for neutral processes

$$\mathcal{M}(u_L^i \bar{u}_L^j \rightarrow W_0^- W_0^+) = -c_W \mathcal{M}(d_L^i \bar{d}_L^j \rightarrow Z_0 h) = -i \frac{\hat{S} \sin \theta}{\Lambda^2} \mathcal{C}_{Hq}^{(1+3)}{}_{ij},$$


$$\mathcal{M}(u_R^i \bar{u}_R^j \rightarrow W_0^- W_0^+) = -c_W \mathcal{M}(d_R^i \bar{d}_R^j \rightarrow Z_0 h) = -i \frac{\hat{S} \sin \theta}{\Lambda^2} \mathcal{C}_{Hd}{}_{ij},$$

$$\mathcal{M}(d_L^i \bar{d}_L^j \rightarrow W_0^- W_0^+) = -c_W \mathcal{M}(u_L^i \bar{u}_L^j \rightarrow Z_0 h) = -i \frac{\hat{S} \sin \theta}{\Lambda^2} \mathcal{C}_{Hq}^{(1-3)}{}_{ij},$$

$$\mathcal{M}(d_R^i \bar{d}_R^j \rightarrow W_0^- W_0^+) = -c_W \mathcal{M}(u_R^i \bar{u}_R^j \rightarrow Z_0 h) = -i \frac{\hat{S} \sin \theta}{\Lambda^2} \mathcal{C}_{Hu}{}_{ij}.$$

WW and Zh probe the **same combination** of couplings.

Weak basis



$$\mathcal{C}_{Hq}^{(1\pm 3)} = \mathcal{C}_{Hq}^{(1)} \pm \mathcal{C}_{Hq}^{(3)}$$

$\mathcal{O}_{\psi^2 H^2 D}$	Operator
$\mathcal{O}_{Hq}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_i \gamma^\mu q_j)$
$\mathcal{O}_{Hq}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H) (\bar{q}_i \gamma^\mu \tau^I q_j)$
$\mathcal{O}_{Hu}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{u}_i \gamma^\mu u_j)$
$\mathcal{O}_{Hd}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{d}_i \gamma^\mu d_j)$
$\mathcal{O}_{Hud}$	$(\tilde{H}^\dagger i D_\mu H) (\bar{u}_i \gamma^\mu d_j) + \text{h.c.}$

High-energy amplitudes


At dimension $d = 6$, for charged processes

$$-c_W \mathcal{M}(u_L^i \bar{d}_L^j \rightarrow W_0 Z_0) = \mathcal{M}(u_L^i \bar{d}_L^j \rightarrow W_0 h) = -i\sqrt{2} \frac{\hat{S} \sin \theta}{\Lambda^2} \mathcal{C}_{Hq}^{(3)}{}_{ij},$$

$$c_W \mathcal{M}(u_R^i \bar{d}_R^j \rightarrow W_0 Z_0) = \mathcal{M}(u_R^i \bar{d}_R^j \rightarrow W_0 h) = -\frac{i}{\sqrt{2}} \frac{\hat{S} \sin \theta}{\Lambda^2} \mathcal{C}_{Hud}^{(3)}{}_{ij}.$$

WZ and Wh probe the **same combination** of couplings.

Weak basis



$\mathcal{O}_{\psi^2 H^2 D}$	Operator
$\mathcal{O}_{Hq}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_i \gamma^\mu q_j)$
$\mathcal{O}_{Hq}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H) (\bar{q}_i \gamma^\mu \tau^I q_j)$
$\mathcal{O}_{Hu}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{u}_i \gamma^\mu u_j)$
$\mathcal{O}_{Hd}$	$(H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{d}_i \gamma^\mu d_j)$
$\mathcal{O}_{Hud}$	$(\tilde{H}^\dagger i D_\mu H) (\bar{u}_i \gamma^\mu d_j) + \text{h.c.}$

On the validity of Drell-Yan tails

EFT validity of Drell-Yan tails

LHC EFT WG note:

Truncation, validity, uncertainties

See also

T. Corbett [2405.04570]

I. Brivio et al. [2201.04974]

S. Dawson et al. [2205.01561]

T. Kim, A. Martin [2203.11976]

R. Contino et al. [1604.06444]

T. Cohen et al. [2111.09895]

T. Corbett et al. [2503.19962]

and more.

Main questions

1. How accurate is the EFT description of DY?
2. Whenever the EFT expansion is valid, which terms of the EFT series should be considered?

Our work

A **direct comparison** between the **EFT** and concrete **UV models**, considering different truncations of the EFT series.

- ↪ Level of agreement between UV model and EFT in different kinematic ranges.
- ↪ Comparison of limits from the UV model and EFT.
- ↪ Benefit: not relying on general methods.

Factors that can play an important role

The **initial quark flavors** (i. e., PDFs)

light vs. heavy quarks.

The **topology** of the **NP contribution**

s- or t-channel mediator.

The **experimental sensitivity**

Depends on the available kinematic information and signal region.

Drell-Yan cross-section — PDF importance

See F. Wilsch presentation [\[link\]](#).

Amplitudes for $q_i \bar{q}_j \rightarrow \ell_\alpha \bar{\ell}_\beta$ consistent with $SU(3)_c \times U(1)_{em}$

$$\mathcal{A}(\bar{q}_i q_j \rightarrow \ell_\alpha^- \ell_\beta^+) = \frac{1}{v^2} \sum_{\Gamma \otimes \Gamma'} \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} (\bar{\ell}_\alpha \Gamma \ell_\beta) (\bar{q}_i \Gamma' q_j) \rightarrow \Gamma \otimes \Gamma' \in \{P_X \otimes P_Y, \gamma_\mu P_X \otimes \gamma^\mu P_Y, \sigma_{\mu\nu} P_X \otimes \sigma^{\mu\nu} P_X\}$$

$X, Y \in \{L, R\}$

$$\mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} \equiv \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij}(\hat{s}, \hat{t})$$

Includes **contact-interactions** and/or **pole contributions**.

Drell-Yan cross-section — PDF importance

See F. Wilsch presentation [\[link\]](#).

Amplitudes for $q_i \bar{q}_j \rightarrow \ell_\alpha^- \bar{\ell}_\beta^+$ consistent with $SU(3)_c \times U(1)_{em}$

$$\mathcal{A}(\bar{q}_i q_j \rightarrow \ell_\alpha^- \ell_\beta^+) = \frac{1}{v^2} \sum_{\Gamma \otimes \Gamma'} \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} (\bar{\ell}_\alpha \Gamma \ell_\beta) (\bar{q}_i \Gamma' q_j) \rightarrow \Gamma \otimes \Gamma' \in \{P_X \otimes P_Y, \gamma_\mu P_X \otimes \gamma^\mu P_Y, \sigma_{\mu\nu} P_X \otimes \sigma^{\mu\nu} P_X\}$$

$$X, Y \in \{L, R\}$$

$$\mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} \equiv \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij}(\hat{s}, \hat{t})$$

Includes **contact-interactions** and/or **pole contributions**.

Parton Luminosities/PDFs

Hadronic LHC cross-section:

$$\sigma(pp \rightarrow \ell_\alpha^- \ell_\beta^+) = \frac{1}{48\pi v^4} \sum_{\Gamma \otimes \Gamma'} \sum_{i,j} \int \frac{d\hat{s}}{s} \int_{-\hat{s}}^0 d\hat{t} \mathcal{L}_{\bar{q}_i q_j} \mathcal{M}_{\Gamma, \Gamma'} \left| \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} \right|^2$$

$$\mathcal{L}_{\bar{q}_i q_j}(\hat{s}) \equiv \int_{\hat{s}/s}^1 \frac{dx}{x} \left[f_{\bar{q}_i}(x, \mu_F) f_{q_j}\left(\frac{\hat{s}}{sx}, \mu_F\right) + (\bar{q}_i \leftrightarrow q_j) \right]$$

Drell-Yan cross-section — PDF importance

See F. Wilsch presentation [\[link\]](#).

Amplitudes for $q_i \bar{q}_j \rightarrow \ell_\alpha^- \bar{\ell}_\beta^+$ consistent with $SU(3)_c \times U(1)_{em}$

$$\mathcal{A}(\bar{q}_i q_j \rightarrow \ell_\alpha^- \ell_\beta^+) = \frac{1}{v^2} \sum_{\Gamma \otimes \Gamma'} \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} (\bar{\ell}_\alpha \Gamma \ell_\beta) (\bar{q}_i \Gamma' q_j) \rightarrow \Gamma \otimes \Gamma' \in \{P_X \otimes P_Y, \gamma_\mu P_X \otimes \gamma^\mu P_Y, \sigma_{\mu\nu} P_X \otimes \sigma^{\mu\nu} P_X\}$$

$$X, Y \in \{L, R\}$$

$$\mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} \equiv \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij}(\hat{s}, \hat{t})$$


Includes **contact-interactions** and/or **pole contributions**.

Hadronic LHC cross-section:

Parton Luminosities/PDFs

$$\sigma(pp \rightarrow \ell_\alpha^- \ell_\beta^+) = \frac{1}{48\pi v^4} \sum_{\Gamma \otimes \Gamma'} \sum_{i,j} \int \frac{d\hat{s}}{s} \int_{-\hat{s}}^0 d\hat{t} \mathcal{L}_{\bar{q}_i q_j} \mathcal{M}_{\Gamma, \Gamma'} \left| \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} \right|^2$$

$$\mathcal{L}_{\bar{q}_i q_j}(\hat{s}) \equiv \int_{\hat{s}/s}^1 \frac{dx}{x} \left[f_{\bar{q}_i}(x, \mu_F) f_{q_j}\left(\frac{\hat{s}}{sx}, \mu_F\right) + (\bar{q}_i \leftrightarrow q_j) \right]$$



Drell-Yan cross-section — PDF importance

See F. Wilsch presentation [\[link\]](#).

Amplitudes for $q_i \bar{q}_j \rightarrow \ell_\alpha^- \bar{\ell}_\beta^+$ consistent with $SU(3)_c \times U(1)_{em}$

$$\mathcal{A}(\bar{q}_i q_j \rightarrow \ell_\alpha^- \ell_\beta^+) = \frac{1}{v^2} \sum_{\Gamma \otimes \Gamma'} \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} (\bar{\ell}_\alpha \Gamma \ell_\beta) (\bar{q}_i \Gamma' q_j) \rightarrow \Gamma \otimes \Gamma' \in \{P_X \otimes P_Y, \gamma_\mu P_X \otimes \gamma^\mu P_Y, \sigma_{\mu\nu} P_X \otimes \sigma^{\mu\nu} P_X\}$$

$$X, Y \in \{L, R\}$$

$$\mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} \equiv \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij}(\hat{s}, \hat{t})$$

Includes **contact-interactions** and/or **pole contributions**.


Hadronic LHC cross-section:

Parton Luminosities/PDFs

$$\sigma(pp \rightarrow \ell_\alpha^- \ell_\beta^+) = \frac{1}{48\pi v^4} \sum_{\Gamma \otimes \Gamma'} \sum_{i,j} \int \frac{d\hat{s}}{s} \int_{-\hat{s}}^0 d\hat{t} \mathcal{L}_{\bar{q}_i q_j} \mathcal{M}_{\Gamma, \Gamma'} \left| \mathcal{F}_{\Gamma, \Gamma'}^{\alpha\beta ij} \right|^2$$

$$\mathcal{L}_{\bar{q}_i q_j}(\hat{s}) \equiv \int_{\hat{s}/s}^1 \frac{dx}{x} \left[f_{\bar{q}_i}(x, \mu_F) f_{q_j}\left(\frac{\hat{s}}{sx}, \mu_F\right) + (\bar{q}_i \leftrightarrow q_j) \right]$$

The **flavor of the initial quarks** can impact the convergence of the EFT.



EFT convergence — resonant mediator

Example 1:

$$Z' \sim (1, 1, 0)$$


(s-channel)

Integrating out the heavy Z' at tree-level

$$\mathcal{L}_{Z'} = -\frac{1}{4}Z'_{\mu\nu}Z'^{\mu\nu} + \frac{m_{Z'}^2}{2}Z'_\mu Z'^\mu + J^\mu Z'_\mu \longrightarrow \mathcal{L}_{\text{eff}}^{Z'} \supset -\frac{J_\mu J^\mu}{2m_{Z'}^2} - \frac{1}{2m_{Z'}^4}(D_\mu J_\nu)(D^\mu J^\nu) + \dots$$

q and $l \rightarrow$ quark and lepton doublets.

Example:




EFT convergence — resonant mediator


EFT parton-level cross-section computed in different orders in Λ^{-1} and normalized to the full model.

Initial Quark Flavor $\rightarrow$


1st gen. quarks



2nd gen. quarks



3rd gen. quarks



$$\begin{aligned} \hat{\sigma} &\propto \left| \mathcal{A}_{\text{SM}} + \frac{\mathcal{A}_6}{\Lambda^2} + \frac{\mathcal{A}_{6 \times 6} + \mathcal{A}_8}{\Lambda^4} + \mathcal{O}_\mathcal{A}(\Lambda^{-6}) \right|^2 \\ &= \underbrace{|\mathcal{A}_{\text{SM}}|^2 + \frac{2\text{Re}(\mathcal{A}_{\text{SM}}^* \mathcal{A}_6)}{\Lambda^2} + \frac{|\mathcal{A}_6|^2}{\Lambda^4}}_{\mathcal{O}_\mathcal{A}(\Lambda^{-2})} + \underbrace{\frac{2\text{Re}(\mathcal{A}_{\text{SM}}^* \mathcal{A}_{6 \times 6} + \mathcal{A}_{\text{SM}}^* \mathcal{A}_8)}{\Lambda^4} + \frac{2\text{Re}(\mathcal{A}_6^* \mathcal{A}_{6 \times 6} + \mathcal{A}_6^* \mathcal{A}_8)}{\Lambda^6} + \frac{|\mathcal{A}_{6 \times 6}|^2 + |\mathcal{A}_8|^2}{\Lambda^8} + \dots}_{\mathcal{O}_\mathcal{A}(\Lambda^{-4})}, \end{aligned}$$

$$\frac{1}{1 - x_V^{-1}} = - \sum_{n=1}^{\infty} x_V^n$$

$$x_V \equiv \hat{s}/m_{Z'}^2$$

*neglecting the width.







Initial Quark Flavor →

Coupling
size ↑

1st gen. quarks

2nd gen. quarks

3rd gen. quarks



EFT convergence — non-resonant mediator

Example 2:

$U_1 \sim (\mathbf{3}, \mathbf{1}, 2/3)$

(t -channel)

Integrating out the heavy U_1 at **tree-level**.



$$\mathcal{L}_{U_1} \supset -\frac{1}{2}U_{1\mu\nu}^\dagger U_1^{\mu\nu} + m_U^2 U_1^{\mu\dagger} U_{1\mu} \\ + (J_\mu^\dagger U_1^\mu + \text{h.c.})$$

$$J_\mu^\dagger = x_L^{i\alpha} \bar{q}_i \gamma_\mu l_\alpha$$

q and $l \rightarrow$ quark and lepton doublets.

$$\mathcal{L}_{\text{eff}}^{U_1} \supset -\frac{J_\mu^\dagger J^\mu}{m_{U_1}^2} - \frac{1}{m_{U_1}^4} (D_\mu J_\nu)^\dagger (D^\mu J^\nu) \\ + \frac{1}{m_{U_1}^4} (D_\mu J_\nu)^\dagger (D^\nu J^\mu) \\ + \dots$$

Example:




EFT convergence — non-resonant mediator


EFT parton-level cross-section computed in different orders in Λ^{-1} and normalized to the full model.

Initial Quark Flavor $\rightarrow$


1st gen. quarks



2nd gen. quarks



3rd gen. quarks




$$\begin{aligned} \hat{\sigma} &\propto \left| \mathcal{A}_{\text{SM}} + \frac{\mathcal{A}_6}{\Lambda^2} + \frac{\mathcal{A}_{6 \times 6} + \mathcal{A}_8}{\Lambda^4} + \mathcal{O}_\mathcal{A}(\Lambda^{-6}) \right|^2 \\ &= \underbrace{|\mathcal{A}_{\text{SM}}|^2 + \frac{2\text{Re}(\mathcal{A}_{\text{SM}}^* \mathcal{A}_6)}{\Lambda^2} + \frac{|\mathcal{A}_6|^2}{\Lambda^4}}_{\mathcal{O}_\mathcal{A}(\Lambda^{-2})} + \underbrace{\frac{2\text{Re}(\mathcal{A}_{\text{SM}}^* \mathcal{A}_{6 \times 6} + \mathcal{A}_{\text{SM}}^* \mathcal{A}_8)}{\Lambda^4} + \frac{2\text{Re}(\mathcal{A}_6^* \mathcal{A}_{6 \times 6} + \mathcal{A}_6^* \mathcal{A}_8)}{\Lambda^6} + \frac{|\mathcal{A}_{6 \times 6}|^2 + |\mathcal{A}_8|^2}{\Lambda^8} + \dots}_{\mathcal{O}_\mathcal{A}(\Lambda^{-4})} \end{aligned}$$

$$\begin{aligned} \frac{1}{1 - x_V^{-1}} &= - \sum_{n=1}^{\infty} (-1)^n x_V^n \\ x_V &\equiv - \frac{t}{m_{U_1}^2} > 0 \end{aligned}$$


Initial Quark Flavor →

Coupling
size ↑





1st gen. quarks



2nd gen. quarks




3rd gen. quarks




Initial Quark Flavor →

Coupling
size


1st gen. quarks






2nd gen. quarks



3rd gen. quarks



In all cases, the EFT appears to converge better towards the model when **truncating the series on amplitude level**.






Initial Quark Flavor →

1st gen. quarks




2nd gen. quarks

3rd gen. quarks

Coupling
size



In all cases, the EFT appears to converge better towards the model when **truncating the series on amplitude level**.



Will the same conclusion hold when we compare the predictions to **data**?

Example: low vs. high- p_T constraints

Motivation:

$R_D^{\text{exp}} = 0.342 \pm 0.026,$ $R_{D^*}^{\text{exp}} = 0.287 \pm 0.012$



In combination, 3σ above the average SM predictions.



Measured at LHCb and B-factories

Sw. Banerjee et al. [2411.18639]

$$R_{D^{(*)}} = \frac{\mathcal{B}(B \rightarrow D^{(*)} \tau \bar{\nu})}{\mathcal{B}(B \rightarrow D^{(*)} l \bar{\nu})} \Bigg|_{l \in \{e, \mu\}}$$



Example: low vs. high- p_T constraints

Motivation:

$$R_D^{\text{exp}} = 0.342 \pm 0.026, \quad R_{D^*}^{\text{exp}} = 0.287 \pm 0.012$$


In combination, 3σ above the average SM predictions.



Measured at LHCb and B-factories


Sw. Banerjee et al. [2411.18639]

$$R_{D^{(*)}} = \left. \frac{\mathcal{B}(B \rightarrow D^{(*)} \tau \bar{\nu})}{\mathcal{B}(B \rightarrow D^{(*)} l \bar{\nu})} \right|_{l \in \{e, \mu\}}$$



Vector leptoquarks can provide a good solution for the discrepancies.

D. Buttazzo et al. [1706.07808], A. Angelescu et al. [1808.08179], C. Cornella et al. [1903.11517], and more.



Example: low vs. high- p_T constraints

Motivation:

$$R_D^{\text{exp}} = 0.342 \pm 0.026, \quad R_{D^*}^{\text{exp}} = 0.287 \pm 0.012$$


In combination, 3σ above the average SM predictions.



Measured at LHCb and B-factories


Sw. Banerjee et al. [2411.18639]

$$R_{D^{(*)}} = \left. \frac{\mathcal{B}(B \rightarrow D^{(*)} \tau \bar{\nu})}{\mathcal{B}(B \rightarrow D^{(*)} l \bar{\nu})} \right|_{l \in \{e, \mu\}}$$



Vector leptoquarks can provide a good solution for the discrepancies.



D. Buttazzo et al. [1706.07808], A. Angelescu et al. [1808.08179], C. Cornella et al. [1903.11517], and more.



High- p_T DY tails are **complementary** to low-energy observables, and allows the study of the **EFT validity**.

Experimental search for the analysis

Recast of $pp \rightarrow \tau\tau$ ATLAS search [G. Aad et al. [2002.12223]].




Allwicher, Faroughy, Wilsch, Jaffredo, Sumensari
[2207.10741], [2207.10756]

Observables convoluted with

- Parton Shower and Hadronization — **Pythia8**.
- Detector simulation — **Delphes**.

Not possible to fully reconstruct the kinematics of the events.



Example: low vs. high- p_T constraints



Flavor observables

$$\frac{R_{D^{(*)}}}{R_{D^{(*)}}^{\text{SM}}} \simeq 1 - \frac{v^2}{\Lambda^2} \text{Re} \left(\frac{1}{V_{cb}^*} [\mathcal{C}_{lq}^3]_{3323} + [\mathcal{C}_{lq}^3]_{3333} \right) = 1 + \frac{v^2}{2m_{U_1}^2} \text{Re} \left(\frac{x_L^{s\tau} x_L^{b\tau*}}{V_{cb}^*} + |x_L^{b\tau}|^2 \right)$$

Example: low vs. high- p_T constraints




Largest effect in EWPO comes from $Z\nu\bar{\nu}$ vertex corrections

$$\delta g_L^\nu \propto [\mathcal{C}_{Hl}^{(1-3)}]_{33}(m_Z) \simeq \frac{6|y_t|^2}{16\pi^2} \log \frac{m_Z}{\Lambda} [\mathcal{C}_{lq}^{(1+3)}]_{3333} = \frac{6|y_t|^2}{16\pi^2} \log \left[-\frac{m_Z}{m_{U_1}} \frac{|x_L^{b\tau}|^2}{2} \right]$$

Inputs taken from V. Bresó-Pla, A. Falkowski and M. González-Alonso [2103.12074].

Example: low vs. high- p_T constraints





- $\mathcal{O}_\sigma(\Lambda^{-2})$ ■ Full model
- $\mathcal{O}_A(\Lambda^{-2})$
- .-.-.- $\mathcal{O}_A(\Lambda^{-4})$

High- p_T DY tails - $pp \rightarrow \tau\tau$

- ➔ The EFT series has alternating signs, which helps with its convergence.
- ➔ $\mathcal{O}_A(\Lambda^{-2})$: EFT series exhibits $\sim 10\%$ difference from the model for $m_{U_1} = 1.5$ TeV.
- ➔ $\mathcal{O}_A(\Lambda^{-4})$: EFT is an excellent approximation for the model.

Rare decays

$b \rightarrow s\tau\tau$ transitions



[L. Allwicher, D. A. Faroughy, MM, O. Sumensari, F. Wilsh,
In preparation]