

COSMIC-RAY DRIVEN MAGNETIC FIELD DYNAMO IN GALAXIES

Michał Hanasz,
Centre for Astronomy
Nicolaus Copernicus University, Toruń

COLLABORATION:

Toruń:

- Michał Hanasz
- Dominik Wóltański
- Kacper Kowalik

Munich:

- Harald Lesch
- Hanna Kotarba

Kraków:

- Katarzyna Otmianowska-Mazur
- Katarzyna Kulpa-Dybeł
- Hubert Siejkowski
- Grzegorz Kowal
- Marian Soida
- Barbara Kulesza

Bochum:

- Dominik Bomans


MAGNETIC FIELDS IN SPIRAL GALAXIES - RADIO OBSERVATIONS

M51




A. Fletcher et al. 2008

NGC891



M. Krause et al. 2008

Parker instability in the ISM (Parker 1966, 1967)




(from Longair 1994,
*High Energy
 Astrophysics*)


- Cosmic ray gas: an important ingredient - continuously supplied by SN remnants (diffusive shock acceleration), lead to strong buoyancy effects.
 - Kinetic energy of SN II explosion $\sim 10^{51}$ erg \Rightarrow 10 % of E_{SN} \rightarrow acceleration of cosmic rays - charged particles (protons, electrons) accelerated in shocks to relativistic energies

(Hanasz & Lesch 2000, ApJ, 543, 235)

- ⇒ helical magnetic loops form on initially azimuthal magnetic field due to buoyancy of cosmic rays and the Coriolis force



- ⇒ small scale loops reconnect to form larger loops
- ⇒ generation of the large-scale radial m.f.
- ⇒ differential rotation: generation of the azimuthal m.f.



MHD EQUATIONS

$$\frac{\partial \mathbf{V}}{\partial t} + (\mathbf{V} \cdot \nabla) \mathbf{V} = -\frac{1}{\rho} \nabla(p + p_{CR}) + \mathbf{g} + \frac{1}{\rho} \nabla \left(\frac{B^2}{8\pi} \right) + \frac{\mathbf{B} \cdot \nabla \mathbf{B}}{4\pi\rho}$$

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{V}) = 0$$

$$p = c_s^2 \rho \quad (\text{isoth.approx})$$

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{V} \times \mathbf{B}) + \eta \nabla^2 \mathbf{B}$$

CR TRANSPORT EQUATION

Diffusion - advection approximation

(eg. Schlickeiser & Lerche 1985, A&A, 151, 151)

$$\frac{\partial e_{\text{cr}}}{\partial t} + \nabla(e_{\text{cr}} \mathbf{V}) = -p_{\text{cr}} \nabla \mathbf{V} + \nabla(\hat{K} \nabla e_{\text{cr}}) \quad (1)$$

+ CR sources (SN remnants)

$$\rho_{\text{cr}} = (\gamma_{\text{cr}} - 1) e_{\text{cr}} \quad (2)$$

Anisotropic diffusion of CRs

(Giaccalone & Jokipii 1998 , Jokipii 1999, Ryu et al. 2003)

$$K_{ij} = K_\perp \delta_{ij} + (K_\parallel - K_\perp) n_i n_j, \quad n_i = B_i / B, \quad (3)$$

$$K_{\parallel} = 3 \cdot 10^{28} \text{ cm}^{-2}\text{s}^{-1}, \quad K_{\perp} = (1 - 10)\% (K_{\parallel})$$

Original idea: Parker (1992)

Shearing box model:

Hanasz, Kowal, Otmianowska-Mazur & Lesch, 2004, ApJL, 605, 33

Hanasz, Otmianowska-Mazur, Kowal & Lesch, 2009, A&A, 498, 335

- the cosmic ray component: anisotropic diffusion-advection transport (Hanasz and Lesch 2003 - numerical algorithm).
- localized sources of cosmic rays: supernova remnants, exploding randomly in the disk volume, SN shocks & thermal effects neglected
- resistivity of the ISM (see Hanasz, Otmianowska-Mazur and Lesch 2002, and Hanasz and Lesch 2003, Kowal, Hanasz & Otmianowska-Mazur 2003) ⇒ magnetic reconnection.
- differential rotation (+ Coriolis and tidal forces in local simulations)
- realistic vertical disk gravity following the model of ISM in the Milky Way by Ferriere (1998)

GALACTIC DISK MODEL


- Galactic gravitational potential: halo+bulge+disk
(Allen & Santillan 1991)
- Interstellar gas: Global model of ISM for the Milky Way
(Ferriere 1998)
- **No magnetic field at $t = 0$**
- SN rate \propto star formation rate \propto to gas column density: maximum of SN activity at $R_G = 4.5\text{kpc}$, Gaussian distribution of SNe in z-coordinate ($H = 200\text{pc}$)
- 10% of SN energy output is converted to CR energy.
- **weak ($10^{-4}\mu\text{G}$) dipolar, small scale ($r \sim 50\text{pc}$) randomly oriented magnetic field is supplied locally with every SN explosion for $t \leq 1\text{Gyr}$**
- resistive dissipation of small-scale magnetic fields.

MAGNETOHYDRODYNAMICAL SIMULATIONS OF GALACTIC DISKS

Multifluid, parallel (MPI) magnetohydrodynamical code PIERNIK

(Hanasz et al. 2008): <http://piernik.astri.uni.torun.pl>,


GALERIA, TASK Gdańsk, resol. 500x500x200 up to 1000x1000x200 grid cells, $\simeq 100k\text{-}250k$ CPU h per experiment, 400-1600 CPU cores.



Gas density + vectors of gas velocity (left), a and cosmic ray energy density + vectors of magnetic field at $t = 4\text{Gyr}$.


Hanasz, Woltański, Kowalik 2009, ApJ Letter 706L, 155

MAGNETOHYDRODYNAMICAL SIMULATIONS OF GALACTIC DISKS




Colours: – azimuthal (toroidal) magnetic field component blue: $B_\varphi < 0$,
red: $B_\varphi > 0$

Exploding magnetized stars spread weak irregular magnetic fields in the interstellar medium





Colours: – azimuthal (toroidal) magnetic field component blue: $B_\varphi < 0$,
red: $B_\varphi > 0$

Cosmic rays resulting from Supernova Explosions, and disk rotation cause amplification and ordering of magnetic field in the interstellar medium




Hanasz & Lesch (1998)

GROWTH OF MAGNETIC FLUX AND ENERGY IN THE GLOBAL DISK




Amplification timescale of the large-scale magnetic field component:

$$T_{\langle B \rangle} = 270 \text{ Myr} \simeq T_{\text{rot}}$$




SYNTHETIC RADIO MAPS OF POLARIZED SYNCHROTRON EMISSION



X-type structure in edge-on view


Synthetic radio-maps of the simulated galaxy reproduce the main features observed in real galaxies



Spiral structure of magnetic field in the disk plain


MAGNETIC FIELDS IN SPIRAL GALAXIES - RADIO OBSERVATIONS

M51



A. Fletcher et al. 2008


NGC891



M. Krause et al. 2008

BARRED GALAXY – ANALYTICAL POTENTIAL

Axisymmetric gravitational potential + elliptical bar





Colours: – azimuthal magnetic field component blue: $B_\varphi < 0$, red: $B_\varphi > 0$
– antisymmetry of the azimuthal m.f with respect to galactic midplane.

Kulpa-Dybel et al. 2011, ApJL, 733,L18

BARRED GALAXY – ANALYTICAL POTENTIAL



Synthetic radio maps



time = 2.00 Gyr





Beck & Hoernes 1996, Beck 2011

Kulpa-Dybel et al. 2011, ApJL,
733,L18





STARS + GAS



N-body simulations: VINE code (Wetzstein et al 2008)




Evolution of gas density and toroidal magnetic field in the gravitational field of N-body system



Evolution of gas density and toroidal magnetic field in the gravitational field of N-body system



MAGNETIC FIELD AMPLIFICATION IN THE LIVE DISK



Amplification of magnetic energy (red) and azimuthal magnetic flux (green)

- COSMIC RAY DYNAMICS LEADS TO A VERY EFFICIENT MAGNETIC FIELD AMPLIFICATION IN GALACTIC DISKS
- Amplification timescale $\sim t_{\text{rot}}$ \Rightarrow growth of the large-scale magnetic field by several orders of magnitude, fast enough to expect $\sim 1 \mu\text{G}$ magnetic field in galaxies at $z \sim 1 \div 2$
- Dipolar small-scale magnetic fields supplied by exploding stars build up a large scale magnetic field \Rightarrow no need for seed fields of cosmological origin.
- Efficient regularization of the random magnetic field component.
- Growth of magnetic field, driven by SNe, far outside the star forming ring.
- Synthetic radio-maps resemble magnetic field structures in real galaxies.
- Magnetic field in models based on N-body simulations tend to converge to real galaxies.

IN PROGRESS:

- Multicomponent, energy dependent transport of CR nuclei, radioactive decay & nuclear reactions with ISM nuclei \Rightarrow towards GALPROP functionality in dynamically evolving galactic magnetic fields.
- Synchrotron cooling of CR electrons \Rightarrow selfconsistent global galactic models of synchrotron radioemission