

Euro-GRD SuSy

12 November 2007 - ULB Brussels

Minimal Dark Matter

Marco Cirelli

(CNRS, SPHT-CEA/Saclay)

in collaboration with:
N.Fornengo (Torino)
A.Strumia (INFN Pisa)
M.Tamburini (Pisa)

Nuclear Physics B 753 (2006)
and
Nuclear Physics B 787 (2007)
and
work in progress

Statement

DM exists

Statement

DM exists, it requires New Physics beyond the SM

Statement

DM exists, it requires New Physics beyond the SM,
nice examples of which are SuSy, xDims, LH...

Statement

DM exists, it requires New Physics beyond the SM,
nice examples of which are SuSy, xDims, LH...,
that may (or may not) solve crucial problems related
to the EW scale (EW symmetry breaking, hierarchy...)

Statement

DM exists, it requires New Physics beyond the SM,
nice examples of which are SuSy, xDims, LH...,
that may (or may not) solve crucial problems related
to the EW scale (EW symmetry breaking, hierarchy...)
and, on the way, provide DM as a byproduct (LSP, LKP, LTOP...)

Statement

DM exists, it requires New Physics beyond the SM,
nice examples of which are SuSy, xDims, LH...,
that may (or may not) solve crucial problems related
to the EW scale (EW symmetry breaking, hierarchy...)
and, on the way, provide DM as a byproduct (LSP, LKP, LTOP...)
which is a WIMP with $M \sim \text{TeV}$

Statement

DM exists, it requires New Physics beyond the SM,
nice examples of which are SuSy, xDims, LH...,
that may (or may not) solve crucial problems related
to the EW scale (EW symmetry breaking, hierarchy...)
and, on the way, provide DM as a byproduct (LSP, LKP, LTOP...)
which is a WIMP with $M \sim \text{TeV}$ and is stable, provided
that there is a discrete symmetry (R-parity, KK-parity, T-parity...)

Statement

DM exists, it requires New Physics beyond the SM, nice examples of which are SuSy, xDims, LH..., that may (or may not) solve crucial problems related to the EW scale (EW symmetry breaking, hierarchy...) and, on the way, provide DM as a byproduct (LSP, LKP, LTOP...), which is a WIMP with $M \sim \text{TeV}$ and is stable, provided that there is a discrete symmetry (R-parity, KK-parity, T-parity...), and since these are complex theories there are many parameters.

Statement

DM exists, it requires **New Physics** beyond the SM, nice examples of which are SuSy, xDims, LH..., that may (or **may not**) solve crucial problems related to the EW scale (EW symmetry breaking, hierarchy...) and, on the way, provide DM as a byproduct (LSP, LKP, LTOP...) which is a **WIMP** with $M \sim \text{TeV}$ and is stable, provided that there is a **discrete symmetry** (R-parity, KK-parity, T-parity...), and since these are complex theories there are **many parameters**.

Minimalistic approach

On top of the SM, add **only** one extra multiplet $\mathcal{X} = \begin{pmatrix} \chi_1 \\ \chi_2 \\ \vdots \end{pmatrix}$

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \bar{\chi}(i\not{D} + M)\chi \quad \text{if } \mathcal{X} \text{ is a fermion}$$

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + |D_\mu \mathcal{X}|^2 - M^2 |\mathcal{X}|^2 \quad \text{if } \mathcal{X} \text{ is a scalar}$$

and systematically search for the ideal DM candidate...


Minimalistic approach

On top of the SM, add **only** one extra multiplet $\mathcal{X} = \begin{pmatrix} \chi_1 \\ \chi_2 \\ \vdots \end{pmatrix}$

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \bar{\mathcal{X}}(i\not{D} + M)\mathcal{X} \quad \text{if } \mathcal{X} \text{ is a fermion}$$

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + |D_\mu \mathcal{X}|^2 - M^2 |\mathcal{X}|^2 \quad \text{if } \mathcal{X} \text{ is a scalar}$$

gauge interactions



the only parameter,
and will be fixed by Ω_{DM} .

(other terms in the

(one loop mass splitting)

and systematically search for the ideal DM candidate...

The ideal DM candidate is

weakly int., massive, neutral, stable

The ideal DM candidate is

weakly int., massive, neutral, stable

$SU(2)_L$	$U(1)_Y$	spin
<u>2</u>		
<u>3</u>		
<u>4</u>		
<u>5</u>		
<u>7</u>		

$$\mathcal{X} = \begin{pmatrix} \chi_1 \\ \chi_2 \\ \vdots \\ \chi_n \end{pmatrix}$$

these are all possible choices:

$n \leq 5$ for fermions

$n \leq 7$ for scalars

to avoid explosion in the running coupling

$$\alpha_2^{-1}(E') = \alpha_2^{-1}(M) - \frac{b_2(n)}{2\pi} \ln \frac{E'}{M}$$

← (6 is similar to 4)

The ideal DM candidate is

weakly int., massive, neutral, stable

$SU(2)_L$	$U(1)_Y$	spin
$\underline{2}$	$1/2$	
$\underline{3}$	0	
	1	
$\underline{4}$	$1/2$	
	$3/2$	
$\underline{5}$	0	
	1	
	2	
$\underline{7}$	0	

Each multiplet contains a neutral component with a proper assignment of the hypercharge, according to

$$Q = T_3 + Y \equiv 0$$

e.g. for $n = 2$: $T_3 = \begin{pmatrix} +\frac{1}{2} \\ -\frac{1}{2} \end{pmatrix} \Rightarrow |Y| = \frac{1}{2}$

e.g. for $n = 3$: $T_3 = \begin{pmatrix} +1 \\ 0 \\ -1 \end{pmatrix} \Rightarrow |Y| = 0 \text{ or } 1$

etc.

The ideal DM candidate is

weakly int., massive, neutral, stable

$SU(2)_L$	$U(1)_Y$	spin
$\underline{2}$	$1/2$	S
		F
$\underline{3}$	0	S
		F
	1	S
		F
$\underline{4}$	$1/2$	S
		F
	$3/2$	S
		F
$\underline{5}$	0	S
		F
	1	S
		F
	2	S
		F
$\underline{7}$	0	S

Each multiplet contains a neutral component with a proper assignment of the hypercharge, according to

$$Q = T_3 + Y \equiv 0$$

e.g. for $n = 2$: $T_3 = \begin{pmatrix} +\frac{1}{2} \\ -\frac{1}{2} \end{pmatrix} \Rightarrow |Y| = \frac{1}{2}$

e.g. for $n = 3$: $T_3 = \begin{pmatrix} +1 \\ 0 \\ -1 \end{pmatrix} \Rightarrow |Y| = 0 \text{ or } 1$

etc.

The ideal DM candidate is

weakly int., massive, neutral, stable

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)
<u>2</u>	1/2	S	0.43
		F	1.2
<u>3</u>	0	S	2.0
		F	2.6
	1	S	1.4
		F	1.8
<u>4</u>	1/2	S	2.4
		F	2.5
	3/2	S	2.4
		F	2.5
<u>5</u>	0	S	5.0
		F	4.5
	1	S	3.5
		F	3.2
	2	S	3.5
		F	3.2
<u>7</u>	0	S	8.5

The **mass** M is determined by the relic abundance:


$$\Omega_{\text{DM}} = \frac{6 \cdot 10^{-27} \text{ cm}^3 \text{ s}^{-1}}{\langle \sigma_{\text{ann}} v \rangle} \cong 0.24$$

for χ scalar

$$\langle \sigma_{Av} \rangle \simeq \frac{g_2^4 (3 - 4n^2 + n^4) + 16 Y^4 g_Y^4 + 8g_2^2 g_Y^2 Y^2 (n^2 - 1)}{64\pi M^2 g_\chi}$$

for χ fermion

$$\langle \sigma_{Av} \rangle \simeq \frac{g_2^4 (2n^4 + 17n^2 - 19) + 4Y^2 g_Y^4 (41 + 8Y^2) + 16g_2^2 g_Y^2 Y^2 (n^2 - 1)}{128\pi M^2 g_\chi}$$



(- include co-annihilations)
 (- computed for $M \gg M_{Z,W}$)

The ideal DM candidate is

weakly int., massive, neutral, stable

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)
$\underline{2}$	$1/2$	S	1.0
		F	
$\underline{3}$	0	S	2.5
		F	2.7
	1	S	
		F	
$\underline{4}$	$1/2$	S	
		F	
	$3/2$	S	
		F	
$\underline{5}$	0	S	9.4
		F	10
	1	S	
		F	
	2	S	
		F	
$\underline{7}$	0	S	25

Non-perturbative corrections


(and other smaller corrections)

(more later)

induce modifications:

$$\langle \sigma_{\text{ann}} v \rangle \rightsquigarrow R \cdot \langle \sigma_{\text{ann}} v \rangle + \langle \sigma_{\text{ann}} v \rangle_{p\text{-wave}}$$

with $R \sim \mathcal{O}(\text{few}) \rightarrow \mathcal{O}(10^2)$




The ideal DM candidate is

weakly int., massive, neutral, stable

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)	ΔM (MeV)
<u>2</u>	1/2	S		348
		F	1.0	342
<u>3</u>	0	S	2.5	166
		F	2.7	166
	1	S		540
		F		526
<u>4</u>	1/2	S		353
		F		347
	3/2	S		729
		F		712
<u>5</u>	0	S	9.4	166
		F	10	166
	1	S		537
		F		534
	2	S		906
		F		900
<u>7</u>	0	S	25	166

EW loops induce
a **mass splitting** ΔM
inside the n-uplet:

tree level



$$M_Q - M_{Q'} = \frac{\alpha_2 M}{4\pi} \left\{ (Q^2 - Q'^2) s_W^2 f\left(\frac{M_Z}{M}\right) + (Q - Q')(Q + Q' - 2Y) \left[f\left(\frac{M_W}{M}\right) - f\left(\frac{M_Z}{M}\right) \right] \right\}$$

with $f(r) \xrightarrow{r \rightarrow 0} -2\pi r$


The neutral component
is the lightest



The ideal DM candidate is
weakly int., massive, neutral, stable

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)	ΔM (MeV)	decay ch.
<u>2</u>	1/2	S		348	EL
		F	1.0	342	EH
<u>3</u>	0	S	2.5	166	HH^*
		F	2.7	166	LH
	1	S		540	HH, LH
		F		526	LH
<u>4</u>	1/2	S		353	HHH^*
		F		347	(LHH^*)
	3/2	S		729	HHH
		F		712	(LHH)
<u>5</u>	0	S	9.4	166	(HHH^*H^*)
		F	10	166	—
	1	S		537	$(HH^*H^*H^*)$
		F		534	—
	2	S		906	$(H^*H^*H^*H^*)$
		F		900	—
<u>7</u>	0	S	25	166	—

List all **allowed SM couplings**:



e.g. $\chi_{\underline{4}}^{1/2 -1/2} L_{\underline{2}}^{1/2 -1/2} H_{\underline{2}}^{1/2} H_{\underline{2}}^{-1/2}$

dim=5 operator, induces

$$\tau \sim \Lambda^2 \text{TeV}^{-3} \ll t_{\text{universe}}$$


for $\Lambda \sim M_{\text{Pl}}$

The ideal DM candidate is

weakly int., massive, neutral, stable

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)	ΔM (MeV)	decay ch.
<u>2</u>	1/2	S		348	EL
		F	1.0	342	EH
<u>3</u>	0	S	2.5	166	HH^*
		F	2.7	166	LH
	1	S		540	HH, LH
		F		526	LH
<u>4</u>	1/2	S		353	HHH^*
		F		347	(LHH^*)
	3/2	S		729	HHH
		F		712	(LHH)
<u>5</u>	0	S	9.4	166	(HHH^*H^*)
		F	10	166	—
	1	S		537	$(HH^*H^*H^*)$
		F		534	—
	2	S		906	$(H^*H^*H^*H^*)$
		F		900	—
<u>7</u>	0	S	25	166	—

List all **allowed SM couplings**:



e.g. $\chi_{\underline{4}}^{1/2 -1/2} L_{\underline{2}}^{1/2 -1/2} H_{\underline{2}} H_{\underline{2}}^*$

dim=5 operator, induces $\tau \sim \Lambda^2 \text{TeV}^{-3} \ll t_{\text{universe}}$ for $\Lambda \sim M_{\text{Pl}}$

No allowed decay!
Automatically stable!

The ideal DM candidate is

weakly int., massive, neutral, stable

and
not excluded
by direct searches!

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)	ΔM (MeV)	decay ch.
<u>2</u>	1/2	S		348	EL
		F	1.0	342	EH
<u>3</u>	0	S	2.5	166	HH^*
		F	2.7	166	LH
	1	S		540	HH, LH
		F		526	LH
<u>4</u>	1/2	S		353	HHH^*
		F		347	(LHH^*)
	3/2	S		729	HHH
		F		712	(LHH)
<u>5</u>	0	S	9.4	166	(HHH^*H^*)
		F	10	166	—
	1	S		537	$(HH^*H^*H^*)$
		F		534	—
	2	S		906	$(H^*H^*H^*H^*)$
		F		900	—
<u>7</u>	0	S	25	166	—


The ideal DM candidate is

weakly int., massive, neutral, stable

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)	ΔM (MeV)	decay ch.
$\underline{2}$	1/2	S		348	EL
		F	1.0	342	EH
$\underline{3}$	0	S	2.5	166	HH^*
		F	2.7	166	LH
	1	S		540	HH, LH
		F		526	LH
$\underline{4}$	1/2	S		353	HHH^*
		F		347	(LHH^*)
	3/2	S		729	HHH
		F		712	(LHH)
$\underline{5}$	0	S	9.4	166	(HHH^*H^*)
		F	10	166	—
	1	S		537	$(HH^*H^*H^*)$
		F		534	—
	2	S		906	$(H^*H^*H^*H^*)$
		F		900	—
$\underline{7}$	0	S	25	166	—

and
not excluded
by direct searches!

Candidates with $Y \neq 0$
interact as



$$\sigma \simeq G_F^2 M_N^2 Y^2$$

\gg present bounds
e.g. **Xenon**

need $Y = 0$

The ideal DM candidate is

weakly int., massive, neutral, stable

and
not excluded
by direct searches!

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)	ΔM (MeV)	decay ch.
<u>2</u>	1/2	S		348	EL
		F	1.0	342	EH
<u>3</u>	0	S	2.5	166	HH^*
		F	2.7	166	LH
	1	S		540	HH, LH
		F		526	LH
<u>4</u>	1/2	S		353	HHH^*
		F		347	(LHH^*)
	3/2	S		729	HHH
		F		712	(LHH)
<u>5</u>	0	S	9.4	166	(HHH^*H^*)
		F	10	166	—
	1	S		537	$(HH^*H^*H^*)$
		F		534	—
	2	S		906	$(H^*H^*H^*H^*)$
		F		900	—
<u>7</u>	0	S	25	166	—

The ideal DM candidate is

weakly int., massive, neutral, stable
 and **not excluded**

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)	ΔM (MeV)	decay ch.
<u>2</u>	1/2	S		348	EL
		F	1.0	342	EH
<u>3</u>	0	S	2.5	166	HH^*
		F	2.7	166	LH
	1	S		540	HH, LH
		F		526	LH
<u>4</u>	1/2	S		353	HHH^*
		F		347	(LHH^*)
	3/2	S		729	HHH
		F		712	(LHH)
<u>5</u>	0	S	9.4	166	(HHH^*H^*)
		F	10	166	—
	1	S		537	$(HH^*H^*H^*)$
		F		534	—
	2	S		906	$(H^*H^*H^*H^*)$
		F		900	—
<u>7</u>	0	S	25	166	—

The ideal DM candidate is

weakly int., massive, neutral, stable
 and **not excluded**

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)	ΔM (MeV)	decay ch.
<u>2</u>	1/2	S		348	EL
		F	1.0	342	EH
<u>3</u>	0	S	2.5	166	HH^*
		F	2.7	166	LH
	1	S		540	HH, LH
		F		526	LH
<u>4</u>	1/2	S		353	HHH^*
		F		347	(LHH^*)
	3/2	S		729	HHH
		F		712	(LHH)
<u>5</u>	0	S	9.4	166	(HHH^*H^*)
		F	10	166	—
	1	S		537	$(HH^*H^*H^*)$
		F		534	—
	2	S		906	$(H^*H^*H^*H^*)$
		F		900	—
<u>7</u>	0	S	25	166	—

The ideal DM candidate is

weakly int., massive, neutral, stable

and
not excluded

$SU(2)_L$	$U(1)_Y$	spin	M (TeV)	ΔM (MeV)	decay ch.
<u>2</u>	1/2	S		348	EL
		F	1.0	342	EH
<u>3</u>	0	S	2.5	166	HH^*
		F	2.7	166	LH
	1	S		540	HH, LH
		F		526	LH
<u>4</u>	1/2	S		353	HHH^*
		F		347	(LHH^*)
	3/2	S		729	HHH
		F		712	(LHH)
<u>5</u>	0	S	9.4	166	(HHH^*H^*)
		F	10	166	—
	1	S		537	$(HH^*H^*H^*)$
		F		534	—
	2	S		906	$(H^*H^*H^*H^*)$
		F		900	—
<u>7</u>	0	S	25	166	—

← We have a winner!

← and a 2^o place

(other terms in the scalar potential)

Recap:

A fermionic $SU(2)_L$ quintuplet with $Y = 0$ provides a DM candidate with $M = 10$ TeV, which is fully successful:

- neutral

- ***automatically*** stable 

like proton stability in SM!

and

not _{yet} discovered by DM searches.

A scalar $SU(2)_L$ septuplet with $Y = 0$ also does.

(Other candidates can be cured via non-minimalities.)

Detection and Phenomenology

DM detection

direct detection

production at colliders

indirect

γ from annihil in galactic halo or center
(line + continuum)

e^+ from annihil in galactic halo or center

\bar{p} from annihil in galactic halo or center




\bar{D} from annihil in galactic halo or center

$\nu, \bar{\nu}$ from annihil in massive bodies

tracing in Cosmic Rays?

1. Direct Detection

one-loop processes



$$\mathcal{L}_{\text{eff}}^W = (n^2 - (1 - 2Y)^2) \frac{\pi \alpha_2^2}{16 M_W} \sum_q \left[\left(\frac{1}{M_W^2} + \frac{1}{m_h^2} \right) [\bar{\chi} \chi] m_q [\bar{q} q] - \frac{2}{3M} [\bar{\chi} \gamma_\mu \gamma_5 \chi] [\bar{q} \gamma_\mu \gamma_5 q] \right]$$

larger for higher n

Spin-Independent


$$\propto \frac{m_q}{M_W^3}$$

Spin-Dependent

$$\propto \frac{1}{M M_W}$$



$$\langle N | \sum_q m_q \bar{q} q | N \rangle \equiv f m_N \quad \left(f \simeq \frac{1}{3} \right)$$

1. Direct Detection



(NB: no free parameters \Rightarrow one predicted point per candidate)

4. Tracing in Cosmic Rays?



at U high Energy:
 - high production
 - χ^\pm lives long

A **clear track!** DM is no more dark!

But: - production?

requires non-standard acceleration mechanism

- flux?

few events/km² yr above 10¹⁷ eV

- particle ID?

it's fat and fast, but looks like a light slow muon

$$\frac{dE}{dx} \propto \frac{1}{M} E$$

MDM can cross the Earth
 with chain regeneration (like ν_τ).

Small ΔM makes χ^\pm long-living.

Conclusions

The DM problem requires **physics beyond the SM**.

Introducing the **minimal** amount of it, we find some fully successful DM candidates: massive, neutral, *automatically* stable.

The “best” is the
fermionic $SU(2)_L$ quintuplet with $Y = 0$.
($M = 10$ TeV)

Its phenomenology is **precisely computable**:

- can be found in next gen **direct detection** exp's,
- too heavy to be produced at LHC,
- could give signals in indirect detection exp's,
- can be searched for in UHE CR.

Back-up slides

Comparison with SplitSuSy-like models

A-H, Dimopoulos and/or Giudice, Romanino 2004

Pierce 2004; Arkani-Hamed, Dimopoulos, Kachru 2005

Mahbubani, Senatore 2005

SplitSuSy-like

- Higgsino (a fermion doublet)
- + something else (a singlet)
- stabilization by R-parity
- want unification also
- unification scale is low,
need to embed in 5D
to avoid proton decay

Mahbubani, Senatore 2005

MDM

- arbitrary multiplet, scalar or fermion
- nothing else (with $Y=0$)
- automatically stable
- forget unification, it's SM
- nothing

Common feature: the focus is on DM, not on SM hierarchy problem.

The Evidence for DM



1) galaxy rotation curves

$$v_c(r) = \sqrt{\frac{2G_N M(r)}{r}}$$

$$v_c(r) \sim \text{const} \Rightarrow \rho_M(r) \sim \frac{1}{r^2}$$




$$\Omega_M \gtrsim 0.1$$



The Evidence for DM

1) galaxy rotation curves



$$\Omega_M \gtrsim 0.1$$

2) clusters of galaxies

- “rotation curves”
- gravitation lensing
- X-ray gas temperature



$$\Omega_M \sim 0.2 \div 0.4$$




“bullet cluster” - NASA
astro-ph/0608247

[further developments]


The Evidence for DM

1) galaxy rotation curves



$$\Omega_M \gtrsim 0.1$$

2) clusters of galaxies



$$\Omega_M \sim 0.2 \div 0.4$$





3) CMB+LSS(+SNIa:)

WMAP-3yr Boomerang
ACbar DASI
CBI VSA

SDSS, 2dFRGS
LyA Forest Croft
LyA Forest SDSS




$$\Omega_M \approx 0.26 \pm 0.05$$




The Evidence for DM

1) galaxy rotation curves



$$\Omega_M \gtrsim 0.1$$

2) clusters of galaxies



$$\Omega_M \sim 0.2 \div 0.4$$


3) CMB+LSS(+SNIa:)

WMAP-3yr Boomerang
 ACbar DASI
 CBI VSA

SDSS, 2dFRGS
 LyA Forest Croft
 LyA Forest SDSS




$$\Omega_M \approx 0.26 \pm 0.05$$



The Evidence for DM


1) galaxy rotation curves



$$\Omega_M \gtrsim 0.1$$

details


2) clusters of galaxies



$$\Omega_M \sim 0.2 \div 0.4$$

details

3) CMB+LSS(+SNIa:)



$$\Omega_M \approx 0.26 \pm 0.05$$

details


DM is there.

What is DM?

The Evidence for DM

How would the power spectra be **without DM**?
(and no other extra ingredient)

LSS



The thrilling story of the bullet cluster

Farrar, Rosen (2006) astro-ph/0610298

“The bullet goes too fast!”

With a surprising twist, the bullet cluster that just killed MOND repents and reverts into an advocate of a 5th force in the DM sector, that pulled in the merger.



The thrilling story of the bullet cluster

Farrar, Rosen (2006) astro-ph/0610298

“The bullet goes too fast!”

With a surprising twist, the bullet cluster that just killed MOND repents and reverts into an advocate of a 5th force in the DM sector, that pulled in the merger.



Springel, Farrar (2007) astro-ph/0703232

“Not too fast for the law.”

In a breath-taking finale,
Newton and hydro
dynamical laws regain
control: the bullet is a
uncommon guy (7%), but
he is not too fast for them.

The thrilling story of the bullet cluster

Farrar, Rosen (2006) astro-ph/0610298

“The bullet goes too fast!”

With a surprising twist, the bullet cluster that just killed MOND repents and reverts into an advocate of a 5th force in the DM sector, that pulled in the merger.




Springel, Farrar (2007) astro-ph/0703232

“Not too fast for the law.”

In a breath-taking finale, Newton and hydro dynamical laws regain control: the bullet is a uncommon guy (7%), but he is not too fast for them.

The Max Planck Studios in Hollywood seize the opportunity and make a 2.3-billion-years long blockbuster movie.



Non-Minimal terms in the scalar case

Quadratic and quartic terms in \mathcal{X} and H :

$$\lambda_H (\mathcal{X}^* T_{\mathcal{X}}^a \mathcal{X}) (H^* T_H^a H) + \lambda'_H |\mathcal{X}|^2 |H|^2 + \frac{\lambda_{\mathcal{X}}}{2} (\mathcal{X}^* T_{\mathcal{X}}^a \mathcal{X})^2 + \frac{\lambda'_{\mathcal{X}}}{2} |\mathcal{X}|^4$$

[1] [2] [3] [4]

- do not induce decays (even number of \mathcal{X} , and $\langle \mathcal{X} \rangle = 0$)

- [3] and [4] do not give mass terms

- after EWSB, [2] gives a common mass $\sqrt{\lambda'_H} v \approx \mathcal{O}(\lesssim 100 \text{ GeV})$
to all \mathcal{X}_i components;

negligible for $M = \mathcal{O}(\text{TeV})$

- after EWSB, [1] gives mass splitting $\Delta M_{\text{tree}} = \frac{\lambda_H v^2 |\Delta T_{\mathcal{X}}^3|}{4M} = \lambda_H \cdot 7.6 \text{ GeV} \frac{\text{TeV}}{M}$
between \mathcal{X}_i components;

assume $\lambda_H \lesssim 0.01$ so that $\Delta M_{\text{tree}} \ll \Delta M$

- [1] (and [2]) gives annihilations $\bar{\mathcal{X}} \mathcal{X} \rightarrow \bar{H} H$

assume $|\lambda'_H| \ll g_Y^2, g_2^2$ so that these are subdominant

(Anyway, scalar MDM is less interesting.)

[\[back to Lagrangian\]](#)

[\[back to table\]](#)

Neutralino “properties”

neutralino mass matrix in MSSM ($\tilde{B} - \tilde{W}^3 - \tilde{H}_1^0 - \tilde{H}_2^0$ basis)

$$M_\chi = \begin{pmatrix} M_1 & 0 & -m_Z c_\beta s_W & m_Z s_\beta s_W \\ 0 & M_2 & m_Z c_\beta c_W & -m_Z s_\beta c_W \\ -m_Z c_\beta s_W & m_Z c_\beta c_W & 0 & -\mu \\ m_Z s_\beta s_W & -m_Z s_\beta c_W & -\mu & 0 \end{pmatrix}$$

superpotential

$$\mathcal{W} = -\mu \mathcal{H}_1 \mathcal{H}_2 + \mathcal{H}_1 h_e^{ij} \mathcal{L}_{Li} \mathcal{E}_{Rj} + \mathcal{H}_1 h_d^{ij} \mathcal{Q}_{Li} \mathcal{D}_{Rj} - \mathcal{H}_2 h_u^{ij} \mathcal{Q}_{Li} \mathcal{U}_{Rj}$$

soft SUSYB terms

$$\mathcal{L}_{\text{soft}} = -\frac{1}{2} \left(M_1 \bar{\tilde{B}} \tilde{B} + M_2 \bar{\tilde{W}}^a \tilde{W}^a + M_3 \bar{\tilde{G}}^a \tilde{G}^a \right) + \dots$$





$$\tan \beta = \frac{\langle v_1 \rangle}{\langle v_2 \rangle}$$

Direct detected *already*?

DAMA annual modulation:

however:

- raw data??
- bkgd (Rn emission)
- higher bins not expon suppressed




- DATA listed top to bottom on plot
- DAMA 2000 58k kg-days NaI Ann.Mod. 3sigma,w/o DAMA 1996 limit
- ZEPLIN I Preliminary 2002 result
- Edelweiss, 32 kg-days Ge 2000+2002+2003 limit
- CDMS (Soudan) 2004 Blind 53 raw kg-days Ge
- XENON10 (10 kg) projected sensitivity
- Bottino et al. Neutralino Configurations ($\Omega_{\text{WIMP}} < \Omega_{\text{CDMmin}}$)
- Bottino et al. Neutralino Configurations ($\Omega_{\text{WIMP}} \geq \Omega_{\text{CDMmin}}$)
- CDMSII (Projected) Development ZBG
- XENON100 (100 kg) projected sensitivity
- Chattopadhyay et. al Theory results - post WMAP
- Lahanas and Nanopoulos 2003
- Baer et. al 2003
- Kim/Nihei/Roszkowski/de Austri 2002 JHEP
- Ellis et. al Theory region post-LEP benchmark points
- Masiero, Profumo and Ullio: general Split SUSY
- Baltz and Gondolo 2003

[back to DM detection]

Hints from photons?

EGRET excess



however:


- source not centered
- variability...

+ CANGAROO (2004)

+ HESS (2004)

Ullio et al., ApJ 21 (2004), astro-ph/0308075

WMAP "haze"




Finkbeiner, ApJ 614 (2004)

(Synchrotron rad from e^+e^- from DM annihilations)

The Galactic emission found by Finkbeiner (2004) in the *WMAP* data in excess of the expected foreground Galactic ISM signal may be a signature of such dark matter annihilation.

Hints from positrons?

HEAT excess (1994+95 & 2000)



Baltz, Edsjo, Gondolo, Freese PRD65 (2002)


however:

- random trajectories in magnetic field
- flux requires too much DM...

Neutrinos from DM


Sun

Earth



DM

up-going muons:




SUPERKAMICKANDE INSTITUTE FOR COSMIC RAY RESEARCH UNIVERSITY OF TOYO


[back to DM detection]

“Neutrino Telescopes”

UnderGround



UnderWater



UnderIce



Size: “small”
 Energy thres: GeV
 Energy resol: GeV
 Angle resol: degree

large
 tens GeV
 10 GeV
 few degrees

large/huge
 100 GeV
 tens GeV
 tens degrees

[back to DM detection]

2. Production at colliders

$$\hat{\sigma}_{u\bar{d}} = \frac{g_{\mathcal{X}} g_2^4 (n^2 - 1)}{13824 \pi \hat{s}} \beta \cdot \begin{cases} \beta^2 \\ 3 - \beta^2 \end{cases}$$

if \mathcal{X} is a fermion
if \mathcal{X} is a scalar

(similarly $\hat{\sigma}_{u\bar{u}}, \hat{\sigma}_{d\bar{d}}, \hat{\sigma}_{d\bar{u}}$) $\beta = \sqrt{1 - 4M^2/\hat{s}}$

Large production for small M .

$2 \times$ LHC to produce heavy candidates.

A clean signature:


$$\begin{aligned} \mathcal{X}^{\pm} \rightarrow \mathcal{X}^0 \pi^{\pm} & : \Gamma_{\pi} = (n^2 - 1) \frac{G_F^2 V_{ud}^2 \Delta M^3 f_{\pi}^2}{4\pi} \sqrt{1 - \frac{m_{\pi}^2}{\Delta M^2}}, & \text{BR}_{\pi} = 97.7\% \\ \mathcal{X}^{\pm} \rightarrow \mathcal{X}^0 e^{\pm} (\bar{\nu}_e) & : \Gamma_e = (n^2 - 1) \frac{G_F^2 \Delta M^5}{60\pi^3} & \text{BR}_e = 2.05\% \\ \mathcal{X}^{\pm} \rightarrow \mathcal{X}^0 \mu^{\pm} (\bar{\nu}_{\mu}) & : \Gamma_{\mu} = 0.12 \Gamma_e & \text{BR}_{\mu} = 0.25\% \end{aligned}$$

$$\tau \simeq 44\text{cm}/(n^2 - 1)$$

Events at LHC	
$\int \mathcal{L} dt = 100/\text{fb}$	
$(0.7 \div 2) \cdot 10^3$	
120 \div 260	
0.2 \div 1.0	
0.4 \div 2.2	
11 \div 33	
26 \div 80	
0.1 \div 0.7	
3.6 \div 18	
0.1 \div 0.6	
2.7 \div 14	
$\ll 1$	●
$\ll 1$	
$\ll 1$	◆


Interlude: the “DMtron”

Can one have **CC** DM interactions?
(tree level!)



Need to provide $\Delta M = M_{\chi^+} - M_{\chi} = 166 \text{ MeV}$

Accelerate nuclei and
use DM as diffuse target.



$$\hat{\sigma}(a \chi \rightarrow a' \chi^{\pm}) = \sigma_0 \frac{n^2 - 1}{4} \left[1 - \frac{\ln(1 + 4E^2/M_W^2)}{4E^2/M_W^2} \right]$$

$$\sigma_0 = \frac{G_F^2 M_W^2}{\pi} = 1.1 \cdot 10^{-34} \text{ cm}^2$$

$$\frac{dN}{dt} = \epsilon N_p \sigma \frac{\rho_{\text{DM}}}{M} = \epsilon \frac{10}{\text{year}} \frac{N_p}{10^{20}} \frac{\rho_{\text{DM}}}{0.3 \text{ GeV/cm}^3} \frac{\text{TeV}}{M} \frac{\sigma}{3\sigma_0}$$

“efficiency”
number of bullets
number of targets

not
unreasonable?
tagging χ^+ ...

[skip to conclusions]


3. Indirect Detection

i.e. $\nu, \bar{p}, e^+, \gamma, \bar{D}$ from MDM annihilations in halo or body.

Signal in ν : promising at neutrino telescopes


Enhanced cross section in vector bosons due to resummed diagrams when Non-Relativistic $\bar{\chi}\chi$ are a “bound state”:

$$\alpha_2 M_W \sim \Delta M \approx E_B \sim \alpha_2^2 M$$



Hisano et al., 2004,
Hisano et al., 2005

e.g.



resonances match M for $n = \underline{3}$

Signal in \bar{p}, e^+, γ : promising if enhanced

3. Indirect Detection

For instance, predicted signal in γ rays:

