

# COMPASS results on gluon polarisation




Luís Silva

LIP – Lisbon

[l Silva@lip.pt](mailto:l Silva@lip.pt)


22 Jul 2011



On behalf of the COMPASS Collaboration

## Outline:

- Brief Motivation
- High  $p_T$  and Open Charm analyses
- $\Delta G/G$  results
- Summary and Conclusion



# THE COMPASS EXPERIMENT

Beam:  $2 \cdot 10^8 \mu^+$ / spill  
Luminosity:  $\sim 5 \cdot 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$   
Beam polarisation: 80%  
Beam momentum: 160 GeV/c

Data taken: 2002 - 2011, ...




~250 physicists  
25 institutes  
11 countries

# The COMPASS Spectrometer

Common Muon and Proton Apparatus for Structure and Spectroscopy

- Trackers
- Magnets
- RICH
- Electromagnetic Calorimeters
- Hadronic Calorimeters
- Absorbers
- Target

Two staged spectrometer:  
LAS and SAS  
Polarised beam and target



NIM A577 (2007) 455

Acceptance:


70 mrad (2002-04)

180 mrad (2006)

About 350 detector planes

Track reconstruction  $p > 0.5 \text{ GeV}/c$

# The Nucleon Spin



$$S_N = \frac{1}{2} = \frac{1}{2} \Delta\Sigma + \Delta G + L$$

Quarks

Gluons

Partons  
Orbital Angular  
Momenta

Future !  
GPDs

Well known !

In 1988 EMC measured

$\Delta\Sigma = 0.12 \pm 0.17$  (Phys.Lett.B206,364)

A recent result, including COMPASS, gives:

$\Delta\Sigma = 0.30 \pm 0.01$ (stat.)  $\pm 0.02$ (evol.) Phys.Lett.B647,8


Poorly known

Exploratory and discovery stage.  
Some experiments and data  
might give hints.

COMPASS, HERMES, CLAS,  
STAR, PHOENIX

# Direct measurement of $\Delta G/G$

$$\gamma^* g \rightarrow q \bar{q}$$



To select this process there are two methods :

- **High transverse momentum hadrons ( $Q^2 < 1$  and  $Q^2 > 1$  (GeVc) $^2$ )**
  - ☺ High statistics.
  - ☹ Physical background: strongly model dependent, requires a very good agreement between MC and Data.
- **Open-charm meson (D mesons)**
  - ☺ Provides the purest sample of PGF events, almost free from background contamination. Small dependence on MC.
  - ☹ Low statistics.


Photon-gluon fusion process (PGF)

# Direct measurement of $\Delta G/G$




2002-2004

$$\gamma^* g \rightarrow q \bar{q}$$



Photon-gluon fusion process (PGF)



$$A_{PGF} = \frac{N_{PGF}^{\rightarrow} - N_{PGF}^{\leftarrow}}{N_{PGF}^{\rightarrow} + N_{PGF}^{\leftarrow}}$$




Experiments with polarised beam and target are sensitive to gluon helicity

# High $p_T$ Analysis

$$A_{LL}^{2h}(x) = \frac{\Delta G}{G}(x_g) a_{LL}^{PGF} \frac{\sigma^{PGF}}{\sigma^{Tot}} + A_1^{LO}(x_C) a_{LL}^C \frac{\sigma^C}{\sigma^{Tot}} + A_1^{LO}(x_{Bj}) D \frac{\sigma^{LO}}{\sigma^{Tot}}$$

$A_1^{LO}$  : estimated by an inclusive sample

Final formula for the gluon polarisation



→ 
$$\frac{\Delta G}{G}(x_g^{av}) = \frac{1}{\beta} \left[ A_{LL}^{2h}(x_{Bj}) + A_{corr} \right]$$

$$\beta = a_{LL}^{PGF} R_{PGF} - a_{LL}^{PGF, incl} R_{PGF}^{incl} \frac{R_{LO}}{R_{LO}^{incl}} - a_{LL}^{PGF, incl} \frac{R_C R_{PGF}^{incl}}{R_{LO}^{incl}} \frac{a_{LL}^C}{D}$$

$$A_{corr} = - \left( A_1(x_{Bj}) D \frac{R_{LO}}{R_{LO}^{incl}} - A_1(x_C) \beta_1 + A_1(x_C') \beta_2 \right)$$

- $A_{LL}^{2h}$  : measured from the two hadron sample.
- $a_{LL}^i$  and  $R_i$  : estimated from MC and parametrised using a Neural Network.




# MC Simulation and Neural Network




- Full chain of MC has been used:  
Generator (LEPTO) + Apparatus  
Simulation (GEANT) +  
Reconstruction Program.
- PDF: MSTW2008LO.
- High  $p_T$  sample:
  - MC with parton shower ON.
  - A new tuning was performed to improve the hadron description.

Data-MC comparison:

$Q^2$ ,  $p_T$  and Hadron Multiplicities.



# High $p_T$ Results, $Q^2 > 1 \text{ (GeV/c)}^2$





# High $p_T$ Results, $Q^2 > 1 \text{ (GeV/c)}^2$



The whole statistics was divided, for the first time, resulting in 3 independent measurements, for  $Q^2 > 1 \text{ GeV/c}$ .

# High $p_T$ Analysis, $Q^2 < 1 \text{ (GeV/c)}^2$

~90 % of our statistics in this sample



2002-2004 Preliminary:

$$\Delta G/G = 0.016 \pm 0.058(\text{stat}) \pm 0.055(\text{syst})$$

2002-2003 Published:

$$\Delta G/G = 0.024 \pm 0.089(\text{stat}) \pm 0.057(\text{syst}) \quad \textit{Phys. Lett. B} 633, 25$$

# Open Charm

- The relation between the number of reconstructed  $D^0$  (for each target cell configuration) and  $\Delta G/G$  is given by:

$$N_t = \mathbf{a} \phi \mathbf{n} (\mathbf{S} + \mathbf{B}) \left( 1 + f P_T P_\mu \left[ a_{LL} \frac{\mathbf{S}}{\mathbf{S} + \mathbf{B}} \frac{\Delta G}{G} + D \frac{\mathbf{B}}{\mathbf{S} + \mathbf{B}} A^{bg} \right] \right), \quad t = (\mathbf{u}, \mathbf{d}, \mathbf{u}', \mathbf{d}')$$

acceptance, muon flux, number of target nucleons

Open Charm event probability

- Each equation is weighted with a signal weight  $w_s = f P_m a_{LL} S / (S+B)$  and also with a background weight  $w_B = f P_m D B / (S+B)$ :

**8 equations with 7 unknowns:**  $\Delta G/G$ ,  $A^{bg}$  + 5 independent  $\alpha = (a\phi n)$  factors

The system is solved by a  $\chi^2$  minimisation

# Neural Network parametrisation

- Two real data samples (with the same cuts applied) are compared by a Neural Network (using some kinematic variables as a learning vector):

- Signal model**

$$gcc = K^+ \pi^- \pi_s^- + K^- \pi^+ \pi_s^+$$

( $D^0$  spectrum: signal + background)






- Background model**

$$wcc = K^+ \pi^+ \pi_s^- + K^- \pi^- \pi_s^+$$

(no  $D^0$  is allowed)


- If the background model is good enough: The Neural Network is able to distinguish the signal from the combinatorial background on a event by event basis (inside *gcc*)

$D^0_{K\pi}$  tagged spectrum in bins of  $\Sigma = S/(S+B)_{NN}$




$$\delta \left( \frac{\Delta G}{G} \right) \propto \frac{1}{FOM}$$

# $\Delta G/G$ Results (LO)





# $\Delta G/G$ Results (LO)




$$\frac{\Delta G}{G} = -0.08 \pm 0.21_{\text{(stat)}} \pm 0.08_{\text{(syst)}} \quad \langle \mathbf{x}_g \rangle = 0.11^{+0.11}_{-0.05}, \quad \langle \mu^2 \rangle = 13 \text{ (GeV/c)}^2$$

# NLO corrections for Open Charm analysis

NLO corrections to the analysing power  $a_{LL}$





# $\Delta G/G$ new Result (NLO)



$$\frac{\Delta G}{G} = -0.20 \pm 0.21 \pm 0.08 \text{ (syst)} \quad \langle x_g \rangle = 0.28^{+0.19}_{-0.10}, \quad \langle \mu^2 \rangle = 13 (\text{GeV}/c)^2$$

**Preliminary:** theoretical uncertainties still under study ( $a_{LL}$ )

# Summary and Conclusion



## Summary:






- The direct measurement methods used in COMPASS experiment were explained.
- Several gluon polarisations results presented are in full agreement among themselves.

## Conclusion:

- All measurements of  $\Delta G/G$  are compatible with zero, around  $X_g \sim 0.1$
- The  $\Delta G$  seems to be small contribution.
- The missing contribution could be in  $L_{\text{partons}}$ .
- COMPASS-II program foresees to measure  $L_{\text{partons}}$  via GPDs.

# Spares

# $D^0$ invariant mass spectra: 2002-2007 data



## Number of $D^0$ :

- Total = 90600
- ${}^6\text{LiD}$  = 65600
- $\text{NH}_3$  = 25000

# Monte Carlo Simulation



This analysis uses **information** from the MC, thus a **strong effort** and **care** to ensure that the MC simulation describes as **good** as possible the **data** was undertaken.

Two **MC samples** were used in the analysis: **high  $p_T$**  and **inclusive** samples.

- Full chain of MC has been used:

Generator (LEPTO) + Apparatus Simulation (GEANT) +  
Reconstruction Program.

- PDF: MSTW2008LO.
- High  $p_T$  sample:

- MC with **parton shower ON** has been used in the analysis.
  - A **new tuning** was performed to **improve** the hadron description.

# MC Tuning




- The purpose of the **MC tuning** is to correct the shapes of the **hadron variables** (momenta) and **fragmentation** (multiplicity).
- In **LEPTO** this can be **achieved** by changing **JETSET** parameters:

PARJ(21)	PARJ(23)	PARJ(24)	PARJ(41)	PARJ(42)
Transverse momentum of the hadron fragmentation			Fragmentation function	

- These **parameters** can be **divided** into **two sets** regarding the **component** of the **trajectory** of the particles: **Transverse** and **longitudinal** variable components.
- The **sets** can be **tuned independently**.  
⇒ The tuning improves substantially the Data-MC agreement.

# Monte Carlo Simulation



$a = \text{PARJ}(41)$


$b = \text{PARJ}(42)$

$$f(z) \propto \frac{1}{2} (1-z)^a \exp\left(-\frac{b m_T^2}{z}\right)$$

COMPASS new tuning  
LEPTO default tuning


PARJ(21)	PARJ(23)	PARJ(24)	PARJ(41)	PARJ(42)
0.34	0.04	2.8	0.025	0.075
0.36	0.01	2.0	0.3	0.58
Transverse momentum of the hadron fragmentation			Fragmentation function	

# Data – Monte Carlo comparison









# Data – Monte Carlo comparison

high- $p_T$  sample: hadron variables ( $p_{T1}$ ,  $p_{T2}$  and  $\sum p_T^2$ )



# Data – Monte Carlo comparison


high- $p_T$  sample: hadron variables ( $p_1$ ,  $p_2$  and multiplicity)



# Weighted method

A Neural Network is used to assign to each event a probability to be originated from one of the three processes (LO, PGF or Compton).

- A **MC** sample is used to train the Neural Network (NN).
- A parametrisation is constructed for all variables involved in the weight.
- A **Data** sample is weighted on an event-by-event basis.



Optimal usage of the data sample statistics

# Weighted method

- A weight is applied on event-by-event basis:

$W = fDP_b \beta$  , where  $\beta$  is a factor depending on  $a_{LL}^i$  and  $R^i$

- Therefore for every event we have to know:

$R_{PGF}, R_C, R_{LO}, R_{PGF}^{incl}, R_C^{incl}, R_{LO}^{incl},$

$a_{LL}^{PGF}, a_{LL}^C, a_{LL}^{PGF,incl}, a_{LL}^{C,incl},$

$x_C, x_G,$




$f, D, P_b$

$f, D, P_b$  are directly obtained from data.

The all the others variables have to be estimated/parametrised.

# Example: Stability plots for NN

We parametrise the  $R^i$  fractions as probabilities.




# Results

$$\frac{\Delta G}{G} = 0.125 \pm 0.060 \pm 0.063$$

$$x_G = 0.09^{+0.08}_{-0.04}$$

$$\langle \mu^2 \rangle = 3.4 \text{ (GeV/c)}^2$$



# Systematic Uncertainties



Sources of Systematic Uncertainties	$\delta(\Delta G/G)$	
	High pT	Open Charm
MC Simulation	0.05	
Formula Simplification	0.04	
False Asymmetries	0.02	0.08
$A_1$ Parametrisation	0.02	
NN Parametrisation	0.01	
$P_B, P_T, f$	0.004	0.01
$a_{LL}$		0.01
$s/(s+b)$		0.01
Total	0.06	0.08





# S/(S+B): Obtaining final probabilities for a $D^0$ candidate

- Events with small  $S/(S+B)_{NN}$

- Mostly combinatorial background is selected


$S/(S+B)$  is obtained from a fit inside this bins (correcting with the NN parameterisation)

$D^0_{K\pi}$  tagged spectrum in bins of  $\Sigma = S/(S+B)_{NN}$



- Events with large  $S/(S+B)_{NN}$

- Mostly Open Charm are selected





$$\delta \left( \frac{\Delta G}{G} \right) \propto \frac{1}{FOM}$$

# Neural Network qualification of events

- Two real data samples (with the same cuts applied) are compared by a Neural Network (using some kinematic variables as a learning vector):
  - Signal model →  $gcc = K^+ \pi^- \pi_s^- + K^- \pi^+ \pi_s^+$  ( $D^0$  spectrum: signal + background)
  - Background model →  $wcc = K^+ \pi^+ \pi_s^- + K^- \pi^- \pi_s^+$  (no  $D^0$  is allowed)
- If the background model is good enough: The Neural Network is able to distinguish the signal from the combinatorial background on a event by event basis (inside  $gcc$ )

Example of a good learning variable



# Analysing power (muon-gluon asymmetry $a_{LL}$ )


- $a_{LL}$  is dependent on the full knowledge of the partonic kinematics:

$$a_{LL} = \frac{\Delta \sigma^{\text{PGF}}}{\sigma_{\text{PGF}}} (y, Q^2, x_g, z_C, \phi)$$



Can't be experimentally obtained: only one charmed meson is reconstructed

- $a_{LL}$  is obtained from Monte-Carlo (in LO), to serve as input for a Neural Network parameterisation on some reconstructed kinematical variables:  $y$ ,  $x_{Bj}$ ,  $Q^2$ ,  $z_D$  and  $p_T$



Parameterised  $a_{LL}$ , shows a strong correlation with the generated one (using AROMA)

# Comparison of $a_{LL}$ (LO) with $a_{LL}$ (NLO)

- The AROMA generator is used to simulate the phase space for the NLO (PS on) / LO (PS off) calculations of  $a_{LL}$ . The resulting  $D^0$  mesons are reconstructed in the COMPASS spectrometer like real events. The respective  $a_{LL}$  distributions are:

