

NLO QCD and electroweak corrections to Higgs strahlung off W/Z bosons at Tevatron and the LHC with HAWK

Thursday, 21 July 2011 17:35 (15 minutes)

Higgs strahlung off W/Z bosons at Tevatron and the LHC, $p\bar{p}/pp \rightarrow WH/ZH$, is an important process class for discovering a light Higgs boson. In the talk, first results on electroweak (and QCD) corrections for these reactions are shown that support the full differential information and the decays of the W/Z bosons, i.e. the actually considered processes are $p\bar{p}/pp \rightarrow l \nu H / l l H$. The precise knowledge of differential distributions is particularly important in view of the fact that reconstructing Higgs strahlung at the LHC has to proceed via the investigation of “fat b jets” of highly boosted Higgs bosons and that electroweak corrections usually show strong variations in transverse-momentum distributions. Both results and details of the calculation, which is embedded in the HAWK Monte Carlo generator first designed for Higgs production via vector-boson fusion, are presented.

Primary author: Prof. DITTMAYER, Stefan Dittmaier (University of Freiburg)

Co-authors: Dr MUECK, Alexander (RWTH Aachen); Prof. DENNER, Ansgar (University of Wuerzburg); Dr KALLWEIT, Stefan (PSI Villigen)

Presenter: Prof. DITTMAYER, Stefan Dittmaier (University of Freiburg)

Session Classification: Higgs and New Physics