

5th LEA-COLLIGA Meeting



Rapport sur les contributions

ID de Contribution: 0

Type: **Non spécifié**

In Flight Fast Timing measurements at VAMOS.

mardi 15 novembre 2011 10:50 (30 minutes)

In septembre 2011 we performed the first In Flight Fast Timing (IFFT) measurements at GANIL on ^{197}Au , ^{152}Sm and ^{172}Yb nuclei presenting 1.9ns, 1.4ns, 100ps and 57ps isomers.

States of interest in the target nuclei were populated via Coulomb excitation induced by a ^{136}Xe beam accelerated at 5 A.MeV.

Desexcitation gamma-rays were detected using an array consisting on 16 LaBr3 fast scintillators. The recoiling nuclei were identified in the VAMOS large acceptance spectrometer that also served for trajectory reconstruction, mandatory for gamma ray detection efficiency correction.

In this talk, I propose to present preliminary results showing that IFFT method is a powerful tool for measuring lifetimes in the range of some tens of ps up to about 10 ns hardly reachable, in flight, using other experimental methods.

Author: Dr GAUDEFROY, Laurent (CEA-France)

Orateur: Dr GAUDEFROY, Laurent (CEA-France)

ID de Contribution: 1

Type: **Non spécifié**

Multinucleon transfer reactions in the $^{40}\text{Ar}+^{208}\text{Pb}$ and $^{36}\text{S}+^{208}\text{Pb}$ systems

mardi 15 novembre 2011 15:00 (30 minutes)

Two multinucleon transfer experiments performed at the LNL will be presented.

- The $^{40}\text{Ar}+^{208}\text{Pb}$ reaction was used to populate 40-43Ar isotopes via the 0n to 3n channels. Residues were identified in the Prisma spectrometer and the coincident gamma rays in the Clara gamma array.
- More recently, the $^{36}\text{S}+^{208}\text{Pb}$ reaction (July 2011) allowed the study of lifetimes of intruder states in N~20 Si, P and S isotopes. Recoils were tagged in the Prisma spectrometer at the grazing angle and gamma-rays were detected in the AGATA demonstrator associated with the Köln plunger device for lifetimes measurement.

Author: Dr COURTIN, Sandrine (IPHC and UDS)

Orateur: Dr COURTIN, Sandrine (IPHC and UDS)

ID de Contribution: 2

Type: **Non spécifié**

Study and comparison of the decay modes of the systems formed in the reactions $^{78}\text{Kr}+^{40}\text{Ca}$ and $^{86}\text{Kr}+^{48}\text{Ca}$ at 10 AMeV

mardi 15 novembre 2011 17:20 (30 minutes)

The first results of the ISODEC experiment, performed at the INFN-Laboratori Nazionali del Sud (LNS) by using the CHIMERA detector, will be presented. The principal aim of this experiment is to study the competition between the various disintegration modes of $^{118,134}\text{Ba}$ produced in the reactions $^{78}\text{Kr}+^{40}\text{Ca}$ and $^{86}\text{Kr}+^{48}\text{Ca}$ at 10 AMeV, exploring the isospin dependence of the decay modes of medium mass compound nuclei. The experiment complements data already obtained at 5.5 MeV/A for $^{78,82}\text{Kr}+^{40}\text{Ca}$ reactions, previously realized with beams delivered by GANIL facility and by using the INDRA detector.

Staggering effects are evident in the Z distributions, as well as different isotopic composition and enrichment for the reaction products in the two systems.

Absolute cross sections calculations of the reaction products are in progress, to provide important indication on the isospin influence on the reaction mechanism and fragments production.

Comparisons with theoretical models are in progress to estimate the influence of structural effects during the separation phase in asymmetric fission.

Author: POLITI, Giuseppe (Dipartimento di Fisica e Astronomia - Università di Catania - ITALY)

Orateur: POLITI, Giuseppe (Dipartimento di Fisica e Astronomia - Università di Catania - ITALY)

ID de Contribution: 3

Type: **Non spécifié**

BETA-DECAY SPECTROSCOPIC STUDIES OF THE NEUTRON-RICH $^{211,212,213}\text{Tl}$ and ^{219}Bi ISOTOPES

mardi 15 novembre 2011 16:00 (20 minutes)

The study of the beta-decay process in heavy neutron-rich systems is of main importance to probe the nuclear models used in r-process calculations. Experimental evidence is particularly interesting in nuclei approaching the waiting point $A \sim 195$, since the r-process nuclei are still inaccessible in laboratory and the beta-decay models used to extrapolate their properties show strong discrepancies in their predictions [1,2].

Here we present the first results of an experiment focused on the investigation of the neutron-rich Tl isotopes, carried out within the “Stopped beam Campaign” of the RISING collaboration at GSI. The nuclei of interest were produced in fragmentation reactions of a relativistic Uranium beam impinging on a thick Be target. The residues were subsequently identified in the magnetic spectrometer Fragment Separator (FRS) and were finally implanted in the RISING Active Stopper [3]. This device consisted of nine Double Sided Silicon Strip Detectors (DSSSD) that recorded the position and time of implantations and beta-electrons. The characteristic gamma-ray transitions of the daughter Pb nuclei were registered using the RISING gamma-ray spectrometer [4], placed in close isotropic geometry around the Active Stopper.

The event-by-event position and time correlations between implantations and gamma-labeled radioactive electrons allowed us to measure the beta-decay half-lives of $^{211,212,213}\text{Tl}$, as well as the low-energy structure of their daughter nuclei $^{211,212,213}\text{Pb}$. The comparison of the new lifetimes with the calculations of the nuclear models proposed to describe the r-process provides a significant experimental constraint to their validity near the third r-process abundance peak, confirming previous half-life measurements near the shell closure $N=126$ [5-7].

- [1] P. Möller et al., Phys. Rev. C 67 (2003) 055802
- [2] I.N. Borzov, Phys. Rev C 67 (2003) 025802
- [3] R. Kumar et al., Nucl. Instr. Meth. A 598 (2009) 754
- [4] S. Pietri et al., Nucl. Instr. Meth B 261 (2007) 1079
- [5] T. kurtukian et al., Nucl. Instr. Meth A 589 (2008) 472
- [6] N. Alkhomashi et al., Phys. Rev. C 80 (2009) 064308
- [7] A.I. Morales, PhD thesis. Universidad de Santiago de Compostela, 2011

Author: MORALES LOPEZ, Anabel (INFN-Milano)

Co-auteurs: Dr GOTTARDO, Andrea (INFN-Legnaro); Dr BENZONI, Giovanna (INFN-Milano); Dr VALIENTE-DOBON, Jose Javier (INFN-Legnaro); NICOLINI, Roberto (INFN-Milano)

Orateur: MORALES LOPEZ, Anabel (INFN-Milano)

ID de Contribution: 4

Type: **Non spécifié**

Beyond-mean-field theories with zero-range effective interactions: A way to handle the ultraviolet divergence.

Zero-range effective interactions are commonly used in nuclear physics and in other domains to describe many-body systems within the mean field model. If they are used in a beyond mean-field framework, contributions to the total energy that display an ultraviolet divergence are found. We propose a general strategy to regularize this divergence and we illustrate it in the case of the second-order corrections to the equation of state (EOS) of uniform symmetric matter. By setting a momentum cut-off Λ , we show that for every (physically meaningful) value of Λ it is possible to determine a new interaction such that the EOS with the second-order corrections reproduces the empirical EOS, with a fit of the same quality as that obtained at the mean-field level.

Author: GRASSO, Marcella (IPNO)

Orateur: GRASSO, Marcella (IPNO)

ID de Contribution: 6

Type: **Non spécifié**

Search for X-rays characteristic of element with $Z = 120$

mardi 15 novembre 2011 14:30 (30 minutes)

Super-heavy compound nuclei ($Z = 120$ and 124) with long fission times ($t \geq 10$ - 18 s) have been recently evidenced through experiments [1] applying the crystal blocking technique. This method, because it

requires high quality monocrystalline targets, cannot be generalized to define and locate possible islands of stability in the super-heavy region. An alternative approach is to use the fluorescence technique which looks for emission of X-rays characteristic of the compound nucleus formed. The first reaction studied in this way has been $^{238}\text{U}+^{64}\text{Ni}$ at 6.6 MeV/nucleon leading to $Z=120$ compound nuclei. We will discuss this method and its application for the very first time in the super-heavy elements

region, and present our latest results and conclusions.

[1] M. Morjean et al., PRL 101 (2008) 072701

Author: Dr FRÉGEAU, Marc Olivier (GANIL)

Co-auteurs: CHBIHI, Abdou (GANIL); CORSI, Anna (INFN Milano); DROUART, Antoine (CEA/IRFU/SPHN Saclay); CIORTEA, Constantin (IFIN Bucarest); DUMITRIU, D. (IFIN Bucarest); FLUERASU, D. (IFIN Bucarest); FABRIS, Daniela (INFN Padova); JACQUET, Dominique (IPN Orsay); GRAMEGNA, Fabiana (INFN Legnaro); FRANKLAND, John (GANIL); NALPAS, Laurent (CEA/IRFU/SPHN Saclay); TASSAN-GOT, Laurent (IPN Orsay); GUGIU, M (IFIN Bucarest); PARLOG, Marian (LPC Caen et IFIN Bucarest); RIVET, Marie-France (IPN Orsay); MORJEAN, Maurice (GANIL); BARLINI, Sandro (INFN Firenze); MARCHI, Tommaso (INFN Legnaro); KRAVTCHOUK, V. L. (INFN Legnaro); LEDOUX, Xavier (CEA/DAM/DIF F-91297 Arpajon); BONNET, Éric (GANIL)

Orateur: Dr FRÉGEAU, Marc Olivier (GANIL)

ID de Contribution: 7

Type: **Non spécifié**

News from the MAGNEX-EDEN facility at the LNS

mardi 15 novembre 2011 15:30 (30 minutes)

An intense experimental activity has been accomplished during 2011 at the INFN-LNS by the MAGNEX spectrometer team. An overview of the experiments, including some of the main results will be presented. A special emphasis will be given to the activities strongly connecting our group with the French collaborators in the field of two-neutron transfer reactions and their implications on the understanding of neutron-neutron correlations in atomic nuclei.

In addition in the same period the EDEN array of 36 Ne²¹³ liquid scintillators has been moved from IN2P3-IPN Orsay and successfully installed at the INFN-LNS around the MAGNEX scattering chamber. This has opened unprecedented opportunities to face old and new puzzles of nuclear structure and nuclear reactions. A brief report of the status of installation and the commissioning will be given.

Author: CAPPUZZELLO, Francesco (Univ. Catania and INFN-LNS)

Orateur: CAPPUZZELLO, Francesco (Univ. Catania and INFN-LNS)

ID de Contribution: 9

Type: **Non spécifié**

Interdisciplinary activities connected to astrophysical issues at the Orsay Tandem Facility.

lundi 14 novembre 2011 18:30 (30 minutes)

I will present an overview of interdisciplinary activities connected to astrophysical issues at the Orsay Tandem Facility. It will concern the simulation of the Cosmic Rays on the Plank bolometer, the Cosmic Ray processing of interstellar analogue dusts and, the molecule-atom collision program with the AGAT multi-detector for astrochemical purpose.

Author: Dr CHABOT, Marin (Institut de physique Nucléaire d'Orsay)

Orateur: Dr CHABOT, Marin (Institut de physique Nucléaire d'Orsay)

ID de Contribution: 10

Type: **Non spécifié**

SPES Project: a Neutron Rich ISOL Facility for re-accelerated RIBs

lundi 14 novembre 2011 14:30 (30 minutes)

SPES (Selective Production of Exotic Species) is an INFN project to develop a Radioactive Ion Beam (RIB) facility as an intermediate step toward EURISOL.

The SPES Project is under realization at the INFN Legnaro National Laboratories site.

The SPES Project main goal is to provide a production and accelerator system of exotic beams to perform forefront research in nuclear physics by studying nuclei far from stability. The SPES Project is concentrating on the production of neutron-rich radioactive nuclei with mass in the range 80-160.

The final energy of the radioactive beams on target will range from few MeV/u up to 11 MeV/u for $A=130$.

The SPES acceleration system will be presented, together with the facility realization status.

Author: PIAZZA, Leandro (INFN - LNL)

Orateur: PIAZZA, Leandro (INFN - LNL)

ID de Contribution: 11

Type: **Non spécifié**

Beyond-mean-field theories with zero-range effective interactions: A way to handle the ultraviolet divergence.

lundi 14 novembre 2011 15:45 (30 minutes)

Zero-range effective interactions are commonly used in nuclear physics and in other domains to describe many-body systems within the mean field model. If they are used in a beyond mean-field framework, contributions to the total energy that display an ultraviolet divergence are found. We propose a general strategy to regularize this divergence and we illustrate it in the case of the second-order corrections to the equation of state (EOS) of uniform symmetric matter. By setting a momentum cut-off Λ , we show that for every (physically meaningful) value of Λ it is possible to determine a new interaction such that the EOS with the second-order corrections reproduces the empirical EOS, with a fit of the same quality as that obtained at the mean-field level.

Author: Dr GRASSO, Marcella (IPNO)**Orateur:** Dr GRASSO, Marcella (IPNO)

ID de Contribution: 12

Type: **Non spécifié**

In-beam and isomer spectroscopy in the third island of inversion at EXOGAM+VAMOS

mardi 15 novembre 2011 09:30 (30 minutes)

Excited states in a wide range of neutron rich nuclei around ^{68}Ni has been studied in multi nucleon transfer reactions in inverse kinematics using a ^{238}U beam at the Coulomb barrier and the large acceptance spectrometer VAMOS and the segmented germanium clover detectors EXOGAM at GANIL. In a first experiment, the lifetimes of the first excited states in even-even Fe and even-odd Co isotopes towards $N=40$ have been measured for the first time using the recoil-distance Doppler shift method. A sudden increase of collectivity from ^{62}Fe to ^{64}Fe has been observed and probe the weakness of the $N=40$ subshell gap. This observation is confirmed by the onset of a collective character in the neutron rich zinc and germanium isotopes. Moreover, collective $7/2^-$ states observed in the structure of copper isotopes above nickel can be interpreted as a one proton in the $p_{3/2}$ orbital coupled to the first 2^+ state in nickel isotones. Similarly, a one proton hole in the $f_{7/2}$ orbital coupled to the first 2^+ state in nickel isotones would induce some collectivity in the cobalt isotopes as suggested by D. Pauwels et al. To demonstrate this, the lifetime of the first excited $9/2^-$ and $3/2^-$ states in $^{63-65}\text{Co}$ have been measured and the $B(E2)$ transition probability extracted. The data extracted from these experiments have been compared with large scale shell model calculations performed with ANTOINE code. Evidence of coexisting neutron, proton intruder states and “normal” state will be presented. A second experiment was recently done at the GANIL facility aiming to perform combined prompt and delayed spectroscopy at the VAMOS spectrometer in solenoid configuration. Preliminary results on new isomers and prompt decays in odd-mass in the third island of inversion will be presented.

Author: Dr CLEMENT, Emmanuel (CNRS GANIL)

Orateur: Dr CLEMENT, Emmanuel (CNRS GANIL)

ID de Contribution: 13

Type: **Non spécifié**

Upgrading the RIPEN Apparatus with Digital Electronics

mercredi 16 novembre 2011 10:00 (20 minutes)

The RIPEN apparatus is a neutron detector array composed of BC501 liquid scintillators specifically suited for neutron detection and time of flight measurement. It was installed at Legnaro National Laboratory in early '90s, while the last measurement campaign was performed in 2007. At present the apparatus is undergoing a process of complete substitution of readout/acquisition electronics. The capabilities of digital electronics have been tested using CAEN V1720 VME digitizers (12 bit, 250MS/s). Analogue RC/CR emulation filters have been developed to perform neutron gamma discrimination: zero crossing technique as well as gate integrated method have been implemented. In June 2011 a subset of 8 detectors was successfully used to perform an in-beam experiment to measure neutron production cross sections. This required the use of 2 VME synchronized acquisition boards and the development of a specific on-line analysis software. We will present a short description of the RIPEN apparatus at LNL and the digital electronic setup. We will also show preliminary results from the above mentioned in-beam test.

Author: MARCHI, Tommaso (INFN Laboratori Nazionali di Legnaro - Università di Padova)

Orateur: MARCHI, Tommaso (INFN Laboratori Nazionali di Legnaro - Università di Padova)

ID de Contribution: 14

Type: **Non spécifié**

Recent news from the S3 spectrometer at SPIRAL2

mercredi 16 novembre 2011 09:00 (30 minutes)

In this presentation, recent new on the design and construction of the S3 spectrometer will be presented (spectromter design, target station, detection system).

Authors: Dr CLEMENT, Emmanuel (CNRS GANIL); Dr SAVAJOLS, Hervé (CNRS GANIL)

Orateur: Dr CLEMENT, Emmanuel (CNRS GANIL)

Classification de Session: Detection arrays and RIBS

ID de Contribution: 15

Type: **Non spécifié**

Neutron-rich ^{71}Cu by transfer reactions at Ganil

mardi 15 novembre 2011 11:20 (30 minutes)

The series of copper isotopes ($Z=29$) is of prime interest for nuclear structure. The first $5/2$ level in the odd isotopes, carrying most of the $f_{5/2}$ proton strength, sharply drops in energy beyond $N=40$ and becomes the ground state in ^{75}Cu . The position and the strength distribution of the $f_{7/2}$ spin-orbit partner is essential to understand the underlying mechanism of this effect, bent to have major implications for structure variations towards ^{78}Ni . With this purpose in mind the $^{72}\text{Zn}(d,^3\text{He})^{71}\text{Cu}$ transfer reaction was performed at Ganil with the Must-2 particle array, giving precisely access to proton-hole states in copper. The data are currently under analysis.

Author: FRANCHOO, serge (ipn)**Orateur:** FRANCHOO, serge (ipn)

ID de Contribution: 16

Type: **Non spécifié**

Microscopic theory of the gamma decay of nuclear giant resonances

mardi 15 novembre 2011 10:00 (30 minutes)

A microscopic formalism that allows the calculation of the gamma decay of nuclear excited states has been developed. In particular, it has been applied to the direct gamma decay of the Isoscalar Giant Quadrupole Resonance in ^{208}Pb to the ground state and to the low-lying octupole state, as well. The phonons are calculated within fully self-consistent RPA, while the calculation of the gamma decay width is performed at the lowest contributing order of the perturbation theory within the framework of the Nuclear Field Theory (NFT), including consistently the whole effective Skyrme interaction in the particle-vibration coupling vertex. The decay width to the lowest 3- state turns out to be only a few percent of the decay width to the ground state, as indicated by the experiment.

Author: M. BRENNNA, Marco (Dipartimento di Fisica, Università degli Studi di Milano and INFN, Sezione Milano)

Co-auteurs: Prof. COLO', Gianluca (Dipartimento di Fisica, Università degli Studi di Milano and INFN, Sezione Milano); Prof. BORTIGNON, Pier Francesco (Dipartimento di Fisica, Università degli Studi di Milano and INFN, Sezione Milano)

Orateur: M. BRENNNA, Marco (Dipartimento di Fisica, Università degli Studi di Milano and INFN, Sezione Milano)

ID de Contribution: 17

Type: **Non spécifié**

Progress on the development and construction of the 50 kW Neutron Converter for Spiral 2 project

mercredi 16 novembre 2011 09:00 (20 minutes)

LNL-INFN is in charge of the research, development and construction of the 50 kW neutron converter for the Spiral 2 facility for producing radioactive ion beams. In the Spiral 2 project, the deuterons primary beam 40 MeV and 5 mA interacts with a graphite converter for producing a neutron flux of 1×10^{12} neutrons/s/cm². The high neutron flux is required for inducing fissions on the UCx fissile target that leads to reach high fission rates and high intensity of the radioactive ion beams. The design of the converter is based on a graphite rotating wheel, that allows the dissipation of the total power of the primary beam, avoiding complex cooling systems. The design had been conceived in such a way to house also the 200 kW converter. A brief summary of the recent developments and state of art on the construction will be presented.

Author: Mme BERMUDEZ, Judilka (LNL-INFN)

Co-auteurs: Dr UDUP, Emil (HHNI-Bucharest); Dr ACOSTA, Gabriela (LNL-INFN); Dr TECCHIO, Luigi (LNL-INFN)

Orateur: Mme BERMUDEZ, Judilka (LNL-INFN)

ID de Contribution: 18

Type: **Non spécifié**

About Predictive Power of Nuclear Theories and Strengthening the Link with Experiment

lundi 14 novembre 2011 16:45 (30 minutes)

It is (too) often thought (and even said) that the theory has predictive power, when the comparison between the data and the theory graph looks good. [Something looking good for someone is perhaps quite unsatisfactory for someone else, and with such a definition we have an infinity of different predictive powers in the circulation! - in-acceptable in the XXIst century!]

This presentation is oriented for the experimental audiences; The idea originates from a sub-field of Applied Mathematics known under the name of “Inverse Problem” but this latter term has in fact not much in common with the “inverse problem of” quantum mechanics where one finds the potential for the Schrodinger equation, out of the energy spectra and scattering information.

In short: We formulate an approach according to which each theory (in particular the nuclear ones) provides not only its results (numbers) but also probabilities that these numbers appear in nature; For instance - what is the probability that the results for ^{132}Sn obtained with the Hamiltonian optimized for ^{208}Pb , will in fact hold true? [when the experiments will be finally done!]. From this posing of the problem it becomes clear that we will present stochastic analysis of the parameter adjustments of theories [why do we find in the literature over 120 various parametrisations of the Skyrme-HF Hamiltonian - and yet, predictions for exotic nuclei obtained with them are so very different???] and general hints on: What to do? - but first of all - What NOT to do? with a given theory, if one does not want to lose the predictive power from the start?

Authors: SZPAK, Bartek (IFJ, Polish Academy of Sciences, Cracow, Poland); DUDEK, Jerzy (IPHC and Univerite de Strasbourg)

Orateur: DUDEK, Jerzy (IPHC and Univerite de Strasbourg)

ID de Contribution: 19

Type: Non spécifié

Nuclear response to two-neutron transfer via the (18O,16O) reaction

mardi 15 novembre 2011 16:50 (30 minutes)

A study of the structure of different nuclei was pursued at the Catania INFN-LNS laboratory by the (18O,16O) two-neutron transfer reaction at 84 MeV incident energy. The experiments were performed using several solid targets from light (9Be, 11B, 12,13C, 16O, 28Si) to heavier ones (58,64Ni, 120Sn, 208Pb). The 16O ejectiles were detected at forward angles by the MAGNEX magnetic spectrometer. Thanks to an innovative technique the ejectiles were identified without the need of time of flight measurements. Exploiting the large momentum acceptance (20%) and solid angle (50 msr) of the spectrometer, energy spectra were obtained with a relevant yield up to about 20 MeV excitation energy. The application of the powerful trajectory reconstruction technique did allow to get energy spectra with energy resolution of about 100 keV and angular distributions with angular resolution better than 0.3°. In the energy spectra several known low lying and resonant states of the product nuclei have been observed. A common feature observed with light nuclei is the appearance of unknown resonant structures at for example 10.5 and 13.6 MeV in 15C and 16 MeV in 14C. The strong population of these latter together with the measured width can reveal the excitation of a collective mode connected with the transfer of a pair. Considerations based on kinematical matching conditions and on the shell configuration of the explored nuclei explain why such a mode is so excited in such reactions. In addition the measured angular distributions seems to indicate a transfer of a correlated neutron pair in $L = 0$ configuration, compatible with the Giant Pairing Vibration mode. Theoretical calculations have been performed in order to estimate the contribution of the break-up both of two correlated neutrons and of two independent ones.

Author: Dr CAVALLARO, Manuela (INFN - LNS)

Orateur: Dr CAVALLARO, Manuela (INFN - LNS)

ID de Contribution: 20

Type: **Non spécifié**

Anomalous light flashes: from astronauts to protontherapy

lundi 14 novembre 2011 17:50 (20 minutes)

Heavy ion interactions in the eye can induce light flashes (phosphenes). Both Neil Armstrong and Edwin Aldrin reported unexpected visual perceptions during their Moon landing mission, due to cosmic rays interactions. We have undertaken a clinical ground study of proton-induced phosphenes. Sixty percent of the patients treated for choroidal melanomas at the Institut Curie –Centre de Protontherapie in Orsay using 73 MeV protons, report anomalous phosphenes. The present study may be the first indication of phosphenes triggered by protons of few tens of MeV.

Author: KHAN, Elias (IPN Orsay)**Orateur:** KHAN, Elias (IPN Orsay)

ID de Contribution: **21**

Type: **Non spécifié**

Status report of SPIRAL2

lundi 14 novembre 2011 14:10 (20 minutes)

Orateur: Dr LEWITOWICZ, Marek (GANIL)

ID de Contribution: 22

Type: **Non spécifié**

Determination of local energy density functionals from Brueckner-Hartree-Fock calculations.

lundi 14 novembre 2011 16:15 (30 minutes)

Orateur: Dr NGUYEN, van Giai (IPNO)

ID de Contribution: **23**

Type: **Non spécifié**

Protontherapy at LNS

lundi 14 novembre 2011 18:10 (20 minutes)

Orateurs: Dr AGODI, Clementina (LNS-INFN); Prof. CUTTONE, Giacomo (LNS)

ID de Contribution: 24

Type: **Non spécifié**

Low-lying dipole response within the second RPA in $^{40,48}\text{Ca}$ nuclei

lundi 14 novembre 2011 17:15 (30 minutes)

Orateur: Dr GAMBACURTA, Danilo (GANIL)

ID de Contribution: 25

Type: **Non spécifié**

(no title)

ID de Contribution: 26

Type: **Non spécifié**

Recent measurements on neutron transfer reactions at deep sub-barrier energies with PRISMA

mardi 15 novembre 2011 09:00 (30 minutes)

Recent measurements performed in inverse kinematics at deep sub-barrier energies in the $^{96}\text{Zr}+^{40}\text{Ca}$ system will be presented.

Target-like recoils have been fully identified with the large solid angle magnetic spectrometer PRISMA.

The experimental data for one and two neutron transfer channels have been compared with semi-classical microscopic calculations.

For the two neutron transfer channels it is found that the transition to the 0^+ state at ~ 6 MeV, whose wavefunction is dominated by the two neutrons in the $2p_{3/2}$ shell, is much larger than the ground state one. The comparison with the inclusive data reveals that transitions to states with high multipolarity and non-natural parity are important, suggesting that more complex two-particle correlations have to be taken into account.

Author: Dr CORRADI, lorenzo (INFN - Laboratori Nazionali di Legnaro)

Orateur: Dr CORRADI, lorenzo (INFN - Laboratori Nazionali di Legnaro)

ID de Contribution: 27

Type: **Non spécifié**

PARIS detector status

mercredi 16 novembre 2011 09:20 (20 minutes)

Orateur: Dr STEZOWSKI, Olivier (IPN Lyon)

ID de Contribution: **28**

Type: **Non spécifié**

FAZIA status

mercredi 16 novembre 2011 11:20 (20 minutes)

Orateur: Dr GRAMEGNA, Fabiana

ID de Contribution: **29**

Type: **Non spécifié**

(no title)

ID de Contribution: **30**

Type: **Non spécifié**

Latest news from ALTO

lundi 14 novembre 2011 15:00 (30 minutes)

Orateur: Dr IBRAHIM, Fadi (IPNO)

ID de Contribution: **31**

Type: **Non spécifié**

Recent news from the S3 spectrometer at SPIRAL2

mercredi 16 novembre 2011 12:10 (20 minutes)

Orateur: Dr CLÉMENT, Emmanuel (GANIL)

ID de Contribution: 32

Type: **Non spécifié**

Words from Director of GANIL

lundi 14 novembre 2011 13:50 (10 minutes)

Orateur: Dr GALES, Sydney

ID de Contribution: 33

Type: **Non spécifié**

Response of strip detectors

mercredi 16 novembre 2011 11:00 (20 minutes)

Orateur: Dr FISICHELLA, Maria (LNL)

ID de Contribution: 34

Type: **Non spécifié**

Welcome introduction

Orateurs: Dr GRAMEGNA, Fabiana (LNL); Dr SCARPACI, Jean-Antoine (IPNO)

ID de Contribution: 35

Type: **Non spécifié**

Words from Director of LNL

lundi 14 novembre 2011 14:00 (10 minutes)

Orateur: Prof. FIORENTINI, Gianni (LNL)

ID de Contribution: 36

Type: **Non spécifié**

Welcome address from IPNO Director

lundi 14 novembre 2011 13:40 (10 minutes)

Orateur: Dr AZAIEZ, Faiçal (director IPNO)

ID de Contribution: 37

Type: **Non spécifié**

The FRIB-LNS facility and the transfer and purification line for the radioactive ion beam at SPES

mercredi 16 novembre 2011 11:40 (30 minutes)

At the end of the 2010, the optic of the extraction beam line of the superconducting cyclotron of LNS was strongly modified. The installation of new quadrupoles and of a cooled target allows to increase the production and collection efficiency of the radioactive ion beam produced in flight.

A short description of the new optic and of the achieved performances will be presented.

A new device, called high energy chopper, will be installed next year along the extraction line.

This new device will remove a large part of the spurious beam, reducing the amount of the of a factor 10, with a clear upgrade of the FRIB-LNS performances.

The new design of the transfer beam line of the radioactive ion beam for the project SPES will be also presented. The main feature of the High resolution mass separator and of the preliminary study on the RFQ beam cooler will be also presented.

Author: Dr CALABRETTA, Luciano (INFN-LNS)

Co-auteurs: RIFUGGIATO, Danilo (INFN-LNS); COSENTINO, Luigi (INFN-LNS); MAGGIORE, Mario (INFN-LNL); PORCELLATO, annamaria (INFN-LNL); CARUSO, antonio (INFN-LNS); DAINELLI, antonio (INFN-LNL); MAUGERI, antonio (INFN-LNS); ZAPPALÀ, emilio (INFN-LNS); MOISIO, francesca (INFN-LNL); COMUNIAN, michele (INFN-LNL); PASSARELLO, santi (INFN-LNS)

Orateur: Dr CALABRETTA, Luciano (INFN-LNS)

ID de Contribution: **39**

Type: **Non spécifié**

NEDA detector status

mercredi 16 novembre 2011 10:20 (20 minutes)

Orateur: Dr NYBERG, Johan (Uppsala University)

ID de Contribution: **40**

Type: **Non spécifié**

(no title)

ID de Contribution: 41

Type: **Non spécifié**

Status of GALILEO

GALILEO is a project aiming to the building of a 4pi high-resolution gamma-ray array by using GASP tapered detectors and capsules of the EUROBALL Cluster detectors. The array will be located at the National Laboratories of Legnaro where stable beams are provided by the Tandem-ALPI-PIAVE accelerator complex and, in the future, exotic radioactive ion beams will be delivered by SPES. One of the most innovative activities of the project is the transformation of the original EUROBALL 7-capsules cluster detectors in triple cluster detectors. The triple cluster detectors will be placed on a ring at 90 degrees with respect to the beam axis while the GASP tapered detectors will cover symmetrically the forward and backward angles. R&D activities are conducted for the development of a complete cryostat for the triple cluster detector and for building new anti-Compton shields optimized for the shape of the new triple cluster detector. The development of the front-end, digital sampling, preprocessing and readout electronics is advancing in synergy with the recent developments made or ongoing for the AGATA project. The talk will make an overview of the status and perspectives of the GALILEO project within the general context of the gamma-ray spectroscopy activity at the National Laboratories of Legnaro.

Author: Dr UR, Calin Alexandru (INFN Sezione di Padova)

Orateur: Dr UR, Calin Alexandru (INFN Sezione di Padova)

ID de Contribution: **42**

Type: **Non spécifié**

ORGAM 2

mardi 15 novembre 2011 18:20 (30 minutes)

Orateurs: Dr AZAIEZ, Faïçal (IPN Orsay); Dr MATTEA, Iolanda (IPN Orsay)

ID de Contribution: **43**

Type: **Non spécifié**

(no title)

ID de Contribution: 44

Type: **Non spécifié**

A Recoil-Distance Doppler-Shift lifetime experiment on neutron-rich Zn isotopes with AGATA

mercredi 16 novembre 2011 12:30 (20 minutes)

Orateur: Dr LOUCHART, Corinne (CEA)

ID de Contribution: 45

Type: **Non spécifié**

LaBr3 detectors, phototubes, electronics and signal processing: hands-on report about on-going research

mercredi 16 novembre 2011 09:40 (20 minutes)

Since a couple of years, the Milano group (INFN and University) is interested in LaBr3 scintillators as detectors for nuclear physics experiments. Ten detectors (3.5" x 8" size, by Saint Gobain) are available and some of them have already been used (Legnaro, GSI, Riken, etc.) with satisfactory results.

A few physical and technical factors practically limiting the intrinsic performance of LaBr3 detectors have been highlighted and at least partially overcome by proper photo-tube selection, design of a dedicated active voltage divider network, acquisition analysis of signals in shape and subsequent digital algorithms.

Design of a dedicated multi-channel digital acquisition system for LaBr3 detectors is in progress.

Author: RIBOLDI, Stefano (Universita' degli Studi di Milano, INFN)

Co-auteurs: GIAZ, Agnese (Universita' degli Studi di Milano, INFN); Dr MILLION, Benedicte (INFN); BOIANO, Ciro (INFN); Dr CRESPI, Fabio (INFN); CAMERA, Franco (Universita' degli Studi di Milano, INFN); PELLEGRINI, Luna (Universita' degli Studi di Milano, INFN); Dr BLASI, Nives (INFN); Dr WIELAND, Oliver (INFN); NICOLINI, Roberto (Universita' degli Studi di Milano, INFN); BRAMBILLA, Sergio (INFN)

Orateur: RIBOLDI, Stefano (Universita' degli Studi di Milano, INFN)

ID de Contribution: 46

Type: **Non spécifié**

Studies for Aluminum photo-ionization in hot cavity for the SPES Project

mercredi 16 novembre 2011 12:50 (20 minutes)

Beside the Uranium Carbide (UCx) target that will be used in SPES (Selective Production of Exotic Species) project, a SiC target will be first to be used to deliver some p-rich beams and validate the function of the SPES facility. Hitting SiC target with protons, one of the elements coming out from nuclear reaction is Aluminum with its isotopes. In order to obtain an Aluminum ion beam, among various type of ionization techniques, main effectiveness is expected from laser photo-ionization. In the past off line studies on laser photoionization have been made in Pavia Spectroscopy Laboratory, testing the selective laser photoionization on Aluminum Hollow Cathode Lamp as atomic source. At Laboratori Nazionali di Legnaro, recently a XeCl excimer laser was installed in order to provide laser ionization in the hot cavity and start first measurements. Results are promising to justify further studies with this technique, aiming a better characterization of the SPES ion extraction capability under laser photoionization. Furthermore with the incoming Wien filter will be possible to monitor the selectivity of the process.

Author: Dr SCARPA, Daniele (I.N.F.N. - L.N.L.)

Orateur: Dr SCARPA, Daniele (I.N.F.N. - L.N.L.)

ID de Contribution: 47

Type: **Non spécifié**

Study of the $^{60}\text{Fe}(n,\gamma)^{61}\text{Fe}$ reaction of astrophysical interest via (d,p) transfer reaction

mardi 15 novembre 2011 17:50 (30 minutes)

Résumé

^{60}Fe is of special interest in nuclear astrophysics. Indeed, the recent detection of ^{60}Fe characteristic gamma-ray lines by RHESSI and INTEGRAL spacecrafts is a strong evidence that ^{60}Fe nucleosynthesis is still going on in the Galaxy. Moreover, the detection of an ^{60}Ni (daughter-nuclei of ^{60}Fe) excess in presolar grains give us information about the conditions of formation of the early solar system.

However, several reactions involved in ^{60}Fe nucleosynthesis suffer from large uncertainties, which imply uncertainties on the ^{60}Fe yields predicted by stellar models. It is the case of ^{60}Fe destruction reaction, $^{60}\text{Fe}(n,\gamma)^{61}\text{Fe}$.

We studied the direct capture part of the $^{60}\text{Fe}(n,\gamma)^{61}\text{Fe}$ reaction via the $d(^{60}\text{Fe},p)$ transfer reaction at LISE/GANIL. The experimental setup consists in two beam tracking detectors, four MUST2 telescopes, and four EXOGAM clovers. The method used to extract the energy of ^{61}Fe excited states populated via the transfer method as well as the measured angular distributions will be explained. DWBA analysis of the measured differential cross-sections, which allowed us to extract angular momentum and spectroscopic factors of the populated states below the neutron threshold will be presented. Experimental results will finally be compared to shell-model calculations.

Orateur: Mlle GIRON, Sandra (IPNO)