

Soft Physics at RHIC

Michal Šumbera^a (for the STAR and PHENIX Collaborations)

Nuclear Physics Institute ASCR, 25068 Řež, Czech Republic

Abstract. Recent soft physics results from collisions of ultra-relativistic nuclei at Relativistic Heavy Ion Collider (RHIC) operating at Brookhaven National Laboratory (BNL) are reviewed. Topics discussed cover the Beam Energy Scan program with some emphasis on anisotropic particle flow.

1 Introduction

At sufficiently high temperature T or baryon chemical potential μ_B QCD predicts a phase transition from hadrons to the plasma of its fundamental constituents – quarks and gluons. Search for and understanding of the nature of this transition has been a long-standing challenge to high-energy nuclear and particle physics community. In 2005, just five years after start up of RHIC, first convincing arguments on the existence of de-confined partonic matter were published [1]. Exciting discoveries made by four experiments BRAHMS, PHOBOS, PHENIX and STAR on perfect quark-gluon liquid [1], constituent number scaling of particle flow [2, 3], jet quenching [4] and heavy-quark suppression [5] were recently complemented by the first detection of anti-strange nucleus [6] and by the observation of the heaviest anti-nucleus – ${}^4\overline{He}$ [7]. The medium produced in collisions of ultra-relativistic nuclei at RHIC, having a highly non-trivial properties of strongly interacting quark-gluon plasma (sQGP), is definitely worth to study over much broader energy range. A central goal now is to map out as much as possible of the QCD phase diagram in T, μ_B plane trying to understand various ways in which the hadron-to-QGP transition may occur.

While the soft physics results from the high energy frontier, the Large Hadron Collider (LHC) at CERN, are covered by P. Kuijjer's contribution to this workshop [8], the low energy frontier of RHIC is presented in this talk. For the topics not included or not sufficiently covered in depth in this minireview I refer interested reader to consult PHENIX and STAR contributions in recently published proceedings of the *Quark Matter 2011* conference [9].

2 Beam Energy Scan Program

During eleven years of its operation the RHIC machine has delivered a variety of nuclear beams (Au, Cu, d). The most frequently used Au+Au collisions were until recently studied at c.m.s. energies per nucleon-nucleon pair $\sqrt{s_{NN}} = 200$ and 62 GeV. The last few years have witnessed, quite naturally, a shift of experimental activity from the top to lower energies. After few small-statistic exploration Au+Au runs at $\sqrt{s_{NN}} = 22$ GeV in 2005 and at $\sqrt{s_{NN}} = 9.2$ GeV in 2008, two remaining running experiments, PHENIX and STAR, collected large-statistics data sets at $\sqrt{s_{NN}} = 7.7, 11.5$ and 39 GeV in 2010, and 19.6 and 27 GeV in 2011. It is noteworthy that $\sqrt{s_{NN}} = 7.7$ GeV, which is much below the RHIC design injection energy of 19.6 GeV, is also the lowest ever achieved energy of hadron collider. The goal of this Beam Energy Scan (BES) program [11] is to search for the QCD critical point, onset of signature of QGP, and softening of the equation of state. The QCD critical point is a distinct singular

^a e-mail: sumbera@ujf.cas.cz

feature of the phase diagram in a T, μ_B plane, where the nature of the transition changes from a discontinuous (first-order) transition to an analytic crossover. Latter, according to lattice calculations, occurs when $\mu_B \approx 0$ and drives the de-confining phase transition at the top RHIC energy and above.




Fig. 1. Energy-dependence of μ_B (left), T_{ch} and T_{kin} (middle) and $\langle\beta\rangle$. The red stars are new STAR results at midrapidity from 5% most central Au+Au collisions at BES energies and the solid lines are the model calculations [10]. The black symbols are previous results from the RHIC, SPS and AGS experiments [11].

Statistical hadronization model fit to mid-rapidity particle ratios (π^-/π^+ , K^-/K^+ , \bar{p}/p , K^-/π^- and \bar{p}/π^-) from 5% most central Au+Au collisions was used by STAR to extract the chemical freeze-out (vanishing inelastic collisions) conditions [10]. Fig.1 shows that the BES program has extended the μ_B range at the RHIC from around 20MeV to about 400 MeV. π^- , K^- and \bar{p} yields were fit with a blast wave model to extract the kinetic freeze-out (vanishing elastic collisions) conditions [10]. The kinetic freeze-out temperature (T_{kin}) is observed to slightly decrease whereas the collective radial flow velocity $\langle\beta\rangle$ increases with decreasing $\sqrt{s_{NN}}$. The large μ_B values at midrapidity indicate the formation of high net-baryon density matter, which is expected to reach a maximum value around 8 GeV [11].

2.1 Anisotropic flow

Study of the conversion of coordinate space anisotropies into momentum space anisotropies plays a central role in ongoing efforts to characterize the transport properties of sQGP. The azimuthal anisotropic flow strength is usually parametrized via Fourier coefficients $v_n \equiv \langle \cos [n(\phi - \Psi_n)] \rangle$, where ϕ is the azimuthal angle of the particle, Ψ_n is the azimuthal angle of the initial-state spatial plane of symmetry (the reaction plane) and n is the order of the harmonic. The event planes from the higher moments at various rapidities are defined with various reaction plane detectors (e.g. Reaction Plane and Muon Piston Calorimeter detectors at $|\eta|=1.0-2.8$ and $|\eta|=3.1-3.7$, respectively, in PHENIX).

The big surprise at RHIC came from the measurement of the v_2 coefficient, integrated elliptic flow, which brings information on the pressure and stiffness of the equation of state during the earliest collision stages. It was found that v_2 increases by 70% from the top SPS energy $\sqrt{s_{NN}}=17.2$ GeV to the top RHIC energy $\sqrt{s_{NN}}=200$ GeV [1]. The large value of v_2 observed at RHIC and recently also at LHC [8],[12] is one of the cornerstones of the perfect liquid bulk matter dynamics. Moreover, the differential $v_2(p_T)$, characterizing in detail the hydrodynamic response to the initial geometry, seems to be unchanged between the top RHIC energy and LHC energy of $\sqrt{s_{NN}}=2.76$ TeV [8,12]. Hence, both at RHIC and at the LHC created matter behaves as the strongly coupled nearly perfect fluid. The latest results on $v_2(p_T)$ shown on the left panel of Fig.2 allow us to conclude that the interval over which the elliptic flow saturates now extends almost two orders of magnitude: from 2.76 TeV to 39 GeV. Since $v_2(p_T)$ at the top SPS energy is much below the saturation curve it would be interesting to see what happens at already collected $\sqrt{s_{NN}}=27$ and 19.6 GeV BES energies.

At midrapidity smooth distribution of the matter in the overlapping region of two equal-mass incoming nuclei implies vanishing of all odd harmonic. The central panel of Fig.1 shows that, due to fluctuations in the initial matter distribution, this assumption is ill-founded. Moreover, for $39 \text{ GeV} \leq \sqrt{s_{NN}} \leq 200 \text{ GeV}$ the data on $v_3(p_T)$ seems to saturate and so the 'lumpiness' of initial geometry over this energy interval remains the same. Excitation function of $v_4(p_T)$, which could provide additional constraints on initial geometries and transport coefficients, plotted on the right panel of Fig.1, shows the similar saturation. It is noteworthy that the initial state fluctuations also show up in two-particle correlation function $\Delta\eta$ and $\Delta\phi$ for particles with $2 < p_T < 5 \text{ GeV}/c$ from 1% most central Au+Au collisions [14].




Fig. 2. Azimuthal asymmetry coefficients $v_{2,3,4}(p_T)$ of charged hadrons at mid-rapidity ($|\eta| < 0.35$) from centrality 10–20% Au+Au collisions at $\sqrt{s_{NN}}=39, 62$ and 200 GeV [13]. Black lines indicate systematic errors on v_2 . The systematic errors for v_3 and v_4 are indicated by bands on data points.

2.2 Elliptic flow of identified particles

Interestingly, the flow patterns are also reflected in the constituent quark number (ncq) scaling of particle identified data. Plotting v_2/ncq versus $(m_T - m_0)/ncq$ for various particle species, where ncq is the number of constituent quarks of a hadron with mass m_0 and $m_T - m_0$ is its transverse kinetic energy, one finds the data to collapse onto a single universal curve. Suggested by parton coalescence and recombination models [3], the universal scaling of light flavor mesons and baryons [2], including multi-strange baryons and ϕ -meson [3] first observed at the top RHIC energy is now considered as an evidence of partonic collectivity of nuclear matter. For hadrons containing the heavy quarks the situation is less clear. Contrary to substantial elliptic flow of mesons containing the heavy quarks found recently by PHENIX [17] the latest STAR measurements of J/ψ [15] are consistent with $v_2 \approx 0$ disfavoring thus the coalescence scenario of J/ψ production from thermalized charm quarks.




Fig. 3. v_2/ncq as a function of $(m_T - m_0)/ncq$ for various particles and light nuclei produced in 0-80% central Au+Au collisions at $\sqrt{s_{NN}}=7.7$ GeV (left), $\sqrt{s_{NN}}=11.5$ GeV (middle) and $\sqrt{s_{NN}}=39$ GeV (right) [16].

Recent PHENIX measurements [13] of elliptic flow of π^\pm , K^\pm , p and \bar{p} from Au+Au collisions confirm that at $\sqrt{s_{NN}} = 62$ and 39 GeV the ncq -scaling still holds, with some deviations in the $(m_T - m_0)/ncq$ range of 0.2 – 0.5 GeV/c^2 especially for (anti)protons and more prominent for 39 GeV. This observation is confirmed and further extended by the new STAR measurements of the elliptic flow of

particles at $\sqrt{s_{NN}} = 39, 11.5$ and 7.7 GeV [11]. Differences are observed in the v_2 of particles and anti-particles, which increase as $\sqrt{s_{NN}}$ decreases suggesting that the ncq -scaling for all particle species (including nuclei) as observed at top RHIC energies [3] is no longer valid at these lower energies.

Fig.3 shows the STAR data on elliptic flow of various particles produced in 0-80% central Au+Au collisions at $\sqrt{s_{NN}} = 7.7, 11.5$ and 39 GeV [16]. Most of the particle species follow the ncq -scaling, except for the ϕ -mesons, which have v_2 at 11.5 GeV systematically lower than the other hadrons. This may provide an evidence for a change in the degrees of freedom around $\sqrt{s_{NN}} \approx 10$ GeV. If in addition, a hierarchy of the violation of the ncq -scaling could be established when going from p to Λ, Ξ and Ω it could provide further insights into the relative importance of hadronic and partonic phase in the early stage of the reaction.

3 Summary

Recent soft physics results from RHIC have substantially extended our knowledge of hot and dense de-confined QCD matter. The BES program covering a large part of the conjectured QCD phase diagram revealed significant differences in particle and anti-particle v_2 coming from the high net-baryon density at midrapidity. Small v_2 of ϕ -meson indicates that hadronic interactions start to dominate over partonic interactions around 11.5 GeV. Saturation of differential elliptic flow $v_2(p_T)$ from $\sqrt{s_{NN}} = 2.76$ TeV down to $\sqrt{s_{NN}} = 39$ GeV extends substantially the region where the sQGP can be created and studied under controlled laboratory conditions. A non-negligible contribution to azimuthal anisotropy of produced particles comes from the fluctuations in the initial matter distribution of colliding nuclei.

Acknowledgements

This work was supported in part by grants LC07048 and LA09013 of the Ministry of Education of the Czech Republic.

References

1. BRAHMS Collaborations, Nuclear Physics **A757**, (2005) 1; PHOBOS Collaboration, *ibid*, 28; STAR Collaboration, *ibid*, 102; PHENIX Collaboration, *ibid*, 184.
2. S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **91**, (2003) 072301, J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **95**, (2005) 12230.
3. B.I. Abelev *et al.* [STAR Collaboration], Phys. Rev. Lett. **99**, (2007) 112301.
4. K. Adcox *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **88**, (2002) 022301, C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **90**, (2003) 082302.
5. A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **98**, (2007) 172301, B.I. Abelev *et al.* [STAR Collaboration], Phys. Rev. Lett. **98**, (2007) 192301.
6. B.I. Abelev *et al.* [STAR Collaboration], Science **328**, (2010) 58.
7. H. Agakishiev *et al.* [STAR Collaboration], Nature **473**, (2011) 353.
8. P. Kuijter, these proceedings.
9. Proc. 22nd Int. Conf. on Ultra-Relativistic Nucleus-Nucleus Collisions (Annecy, France, 23-28 May 2011), Y. Schutz and U. A. Wiedemann eds., J. Phys. G **38**, (2011) 120301.
10. L. Kumar [STAR Collaboration], *ibid*, 124145.
11. B. Mohanty [STAR Collaboration], *ibid*, 124023.
12. J. Velkovska [CMS collaboration], *ibid*, 124011; J. Jia [ATLAS Collaboration], *ibid*, 124012; R. Snellings [ALICE Collaboration], *ibid*, 124013.
13. X. Gong [PHENIX Collaboration], *ibid*, 124146.
14. P. Sorensen [STAR Collaboration], *ibid*, 124029.
15. H. Masui [STAR Collaboration], *ibid*, 124002 .
16. A. Schmah [STAR Collaboration], *ibid*, 124049.
17. A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. C **84** (2011) 044905.