

Inverse Seesaw

・ ロ マ チ 全部 マ キ 中 マ マ マ

## Lepton Flavour Violation in the Supersymmetric Inverse Seesaw Model

#### Cédric Weiland

in collaboration with Asmâa Abada, Debottam Das and Avelino Vicente

Laboratoire de Physique Théorique d'Orsay, Université Paris-Sud 11, France

Rencontres de Physique des Particules Montpellier, May 15th, 2012

Plan		















- - Neutrino oscillations = Neutral lepton flavour violation. Why not have charged lepton flavour violation (cLFV)?
  - cLFV arises from higher order processes ⇒ negligible in the Standard Model
  - If observed:
    - Clear evidence of physics at a higher scale
    - Probe the origin of lepton mixing
    - Probe the origin of new physics
  - Complementary to other New Physics searches
    - High energy: LHC
    - High intensity:
      - B factories: Rare decays, etc
      - Neutrino dedicated experiments: U<sub>PMNS</sub> non-unitarity, etc

・ ロ マ チ 全部 マ キ 中 マ マ マ

• Other low energy experiments:  $(g-2)_{\mu}$ , EDM, etc

・ ロ ト ・ 雪 ト ・ ヨ ト ・ ヨ ト

## **Motivations**

- BSM to generate  $m_{\nu} \neq 0$ 
  - Radiative models
  - Extra dimensions
  - R-parity violation in supersymmetry
  - Seesaw mechanism → BAU through leptogenesis ?
- The SM doesn't only lack neutrino masses ⇒ The hierarchy problem
  - Strongly coupled theories : Technicolor, Composite Higgs
  - Extra-dimensions : Randall-Sundrum, Large extra dimension
  - Extending the SM field content/gauge group : 2HDM, Little Higgs, 4th generation, etc
  - Supersymmetric extensions : MSSM, NMSSM → Gauge coupling unification, DM candidate, graviton in local SUSY

・ ロ ト ・ 雪 ト ・ 国 ト ・ 日 ト

э

## The Seesaw Mechanisms

- $m_{\nu} \neq 0 \Rightarrow$  New physics at a high scale (> SM)
- Seesaw mechanism: Consider new fields at this scale ( $\sim M_R$ ) and Majorana mass terms  $\Rightarrow$  Generate  $m_{\nu}$  in a renormalizable way
- Example: Type I seesaw  $\mathcal{L}_{mass}^{\text{leptons}} = -Y^{\ell} \bar{L} \Phi \ell_R Y_{\nu} \bar{L} \tilde{\Phi} \nu_R \frac{1}{2} M_R \overline{\nu_R^C} \nu_R + \text{h.c.}$  $\Rightarrow$  After EW symmetry breaking, a neutrino mass matrix appears  $M_{6\times 6}^{\nu}$

$$\begin{split} M^{\nu} &= \begin{pmatrix} 0 & m_D \\ m_D^{\top} & m_R \end{pmatrix} & m_D = vY_{\nu} \text{ Dirac mass matrix} \\ &\Rightarrow \text{ Seesaw limit } M_R \gg m_D \\ &m_{\nu}^{\text{light}} \approx -m_D M_R^{-1} m_D^{\top} & \nu^{\text{light}} \approx \nu_L + \nu_L^C \\ &m_{\nu}^{\text{heavy}} \approx M_R & \nu^{\text{heavy}} \approx \nu_R + \nu_R^C \end{split}$$

•  $M^{\nu}$  symmetric (Majorana  $\nu$ )  $\Rightarrow M^{\nu} = ZD_{\nu}Z^{\dagger}$  with Z unitary matrix  $Z = \begin{pmatrix} V & Y \\ X & W \end{pmatrix}$ 

The same goes for  $M^{\ell}$  the charged leptons mass matrix  $\Rightarrow M^{\ell} = A_R D_{\ell} A_L^{\dagger}$  with  $A_{R,L}$  unitary matrices

 $\Rightarrow U_{PMNS} = A_L^{\dagger} V$  leptonic mixing matrix (similar to  $V_{CKM}$ )

A D > A P > A D > A D >

## Effective approach to seesaw mechanisms

- Notice that lepton number conservation is accidental in the SM (from the gauge group, field content and renormalizability)
- Need to violate L conservation to generate m<sub>ν</sub> ⇒ Effective non-renormalizable operators
- Unique dimension 5 operator for all seesaw mechanisms
   → Violates lepton number L ⇒ Majorana neutrinos

$$\delta \mathcal{L}^{d=5} = \frac{1}{2} c_{ij} \frac{(H \cdot L_i)^{\dagger} (H \cdot L_j)}{\Lambda} + \text{h.c.}$$

- To distinguish the several seesaw mechanisms, either
  - Directly produce the heavy states (LHC, ILC)
  - Look for dimension  $\ge 6$  operators effects  $\rightarrow LFV$

## The Inverse Seesaw Mechanism

• Type I seesaw:  $M_R \simeq 10^{14}$ GeV with natural Yukawa  $Y_{\nu} \sim \mathcal{O}(1)$  or  $M_R \sim 1$ TeV with Yukawa  $Y_{\nu} \sim \mathcal{O}(10^{-6})$ 

 $\Rightarrow$  cLFV is suppressed ( dimension 6 operator  $\propto \frac{Y_{\nu}^{\vee}Y_{\nu}}{|M_{R}|^{2}}$  )

- Inverse seesaw:  $M_R \simeq 1$  TeV with natural Yukawa  $Y_{\nu} \sim \mathcal{O}(1)$  $\Rightarrow$  cLFV is much less suppressed
  - → Might be testable at the LHC and future B factories (SuperB)
- Inverse seesaw  $\Rightarrow$  Consider fermionic gauge singlets  $N_i$  (L = -1, i = 1, 2, 3) and  $X_i$  (L = +1, i = 1, 2, 3) [Mohapatra and Valle, 1986]

$$\mathcal{L}_{inverse} = Y_{\nu ij}H \cdot L_i N_j - (M_R)_{ij}N_iX_j - \frac{1}{2}(\mu_X)_{ij}X_iX_j + \text{h.c.}$$

$$m_{\nu} \approx \frac{m_D^2 \mu_X}{m_D^2 + M_R^2}$$
  
$$m_{1,2} \approx \mp \sqrt{m_D^2 + M_R^2} + \frac{M_R^2 \mu_X}{2(m_D^2 + M_R^2)}$$

With  $m_D = Y_{\mu}v$ 



・ ロ ト ・ 雪 ト ・ ヨ ト ・ ヨ ト

## The Minimal Supersymmetric Model

- Same gauge group  $SU(3)_c \times SU(2)_L \times U(1)_Y$
- Field content = SM fields and their SUSY partners
   ⇒ Except for the Higgs sector → Up- and down-type Higgs
- More than a 100 free parameters, most of them from soft SUSY breaking terms

 $\Rightarrow$  Work in constrained frameworks (or find a SUSY breaking mechanism)

- mSUGRA: 4 free parameters  $m_{1/2}$ ,  $m_0$ ,  $A_0$  and  $sign(\mu) \rightarrow$  Nearly entirely excluded
- Constrained MSSM: 5 free parameters  $m_{1/2}$ ,  $m_0$ ,  $A_0$ ,  $\tan(\beta)$  and  $\operatorname{sign}(\mu) \rightarrow \operatorname{Very}$  restrictive boundary conditions

・ロット (雪) (日) (日)

## Supersymmetric Seesaw Models

- No  $\nu_R$  in the MSSM  $\Rightarrow m_{\nu} = 0$  $\rightarrow$  Implement a seesaw mechanism
- Non diagonal neutrino Yukawa couplings
   ⇒ LFV in the slepton mass matrices (radiatively induced)
   ⇒ LFV at low energies through RGE
- Amount of cLFV proportional to the Yukawa couplings
   ⇒ In the usual seesaw (type I), large scale to accommodate natural Yukawa couplings
   ⇒ Impossible to directly produce ν<sub>R</sub>
- Embed the inverse seesaw in the MSSM
   ⇒ Natural Yukawa couplings with a TeV new Physics scale



・ロット (雪) (日) (日)

## The Supersymmetric Inverse Seesaw Model

- MSSM extended by singlet chiral superfields  $\hat{N}_i$  and  $\hat{X}_i$  (i = 1, 2, 3) with respectively L = -1 and L = +1
- Defined by the superpotential:

$$\mathcal{W} = \varepsilon_{ab} \left[ Y_d^{ij} \hat{H}_d^a \hat{Q}_i^b \hat{D}_j + Y_u^{ij} \hat{Q}_i^a \hat{H}_u^b \hat{U}_j + Y_e^{ij} \hat{H}_d^a \hat{L}_i^b \hat{E}_j + Y_\nu^{ij} \hat{L}_i^a \hat{H}_u^b \hat{N}_j \right. \\ \left. - \mu \hat{H}_d^a \hat{H}_u^b \right] + M_{R_{ij}} \hat{N}_i \hat{X}_j + \frac{1}{2} \mu_{X_{ij}} \hat{X}_i \hat{X}_j$$

• Derive one of the new couplings:

$$Y_{\nu}^{ij}\varepsilon_{ab}L_{i}^{a}\widetilde{N}_{j}\widetilde{H}_{u}^{b}+\text{h.c.}\in-\mathcal{L}$$

- Work with a flavour-blind mechanism for SUSY breaking
- Derive the right-handed sneutrino mass:

$$M_{\tilde{N}}^2 = m_{\tilde{N}}^2 + M_R^2 + Y_{\nu}^{ji*} Y_{\nu}^{ij} v_u^2 \sim M_{\text{SUSY}}^2 \sim (1 \text{TeV})^2$$



## cLFV in Supersymmetric Seesaw Models

- Typically in SUSY, cLFV appears at the one-loop level through RGE-induced slepton mixing  $(\Delta m_{\tilde{L}}^2)_{ij}$ [Borzumati and Masiero, 1986, Hisano et al., 1996, Hisano and Nomura, 1999]  $\Rightarrow (\Delta m_{\tilde{L}}^2)_{ij} \propto (Y_{\nu}^{\dagger} Y_{\nu})_{ij} \ln \frac{M_{GUT}}{M_R}$
- Contribute to all cLFV observables
   → Dominant in most of the SUSY seesaw models
- Type I seesaw ( $Y_{\nu} \sim 1, M_R \sim 10^{14} \text{GeV}$ )  $\rightarrow (\Delta m_{\tilde{L}}^2)_{ij} \propto 5$
- Inverse seesaw ( $Y_{\nu} \sim 1$ ,  $M_R \sim 1$ TeV)  $\rightarrow (\Delta m_{\tilde{L}}^2)_{ij} \propto 30$  $\rightarrow \tilde{N}$ -mediated processes are no longer suppressed

## Z-mediated cLFV

- Photon and Higgs-mediated contributions usually dominate in the MSSM
  - In the SUSY inverse seesaw, 2 orders of magnitude enhancement of Higgs-mediated observables [A. Abada, D. Das and C. W., JHEP 1203 (2012) 100
- Z-mediated contributions are suppressed in the MSSM through cancellations at leading order [Hirsch et al., 2012]
  - No longer true in the SUSY inverse seesaw due to new contributions from the right-handed sneutrino



Relative contributions to BR( $\mu \rightarrow 3e$ )  $\Rightarrow \langle a \rangle \langle a \rangle \langle a \rangle$ 

## Z-mediated cLFV

- Why is there a cancellation in the MSSM ?

   → Neglect chargino mixing: Masses cancel out in the combination of loop functions from different diagram
   ⇒ cLFV∞(Z<sup>†</sup><sub>v</sub>Z<sub>v</sub>)<sub>ij</sub> = 0 [Hirsch et al., 2012]
- What happens in the SUSY inverse seesaw ?



- Diagrams with right-handed sneutrinos are no longer suppressed  $\Rightarrow$  Spoils the cancellation cLFV $\propto \sum_{i} Z_{V}^{ik} Z_{V}^{ij*} Y_{\nu}^{ik*} Y_{\nu}^{ij}$
- Dominant contribution: Z-penguins scale like  $m_Z^{-2}$ while  $\gamma$ -penguins scale like  $m_{SUSY}^{-2}$

• □ > • (□) • • □ > • □ >

#### $\mu - e$ Conversion



 $\mu - e$  conversion rate in Ti(48,22) as a function of  $M = M_R^2/\mu_X$ 

- Different values of  $M_R$ : little impact on observables (blue:  $M_R = 100$  GeV, red:  $M_R = 1$  TeV, black:  $M_R = 10$  TeV)
- Slightly dependent on CMSSM parameters: points from random values of  $m_0$  and  $M_{1/2}$  in the range [0, 3] TeV
- Current experimental limits  $4.3 \times 10^{-12}$  (SINDRUM II)  $\Rightarrow$  $(Y_{\nu}^{\dagger}Y_{\nu})_{12} < 2.7 \times 10^{-5}$



э

## $\mu \rightarrow 3e \text{ VS } \mu \rightarrow e\gamma$



- Z-dominated observable:  $\mu \rightarrow 3e$
- Slightly less constraining than  $\mu e$  conversion in Gold
- Small influence of *M<sub>R</sub>* and SUSY parameters



- Observable without any Z contribution: μ → eγ
- Below  $\mu \rightarrow 3e$  $\Rightarrow$  Not constraining
- Strong influence of *M<sub>R</sub>* and SUSY parameters



## Other Observables and Comments

- Brs of observables like τ → 3μ or τ → μη are dominated by Z-penguins and of the same order than Br(μ → 3e)
   ⇒ No chance to be observed at future B factories without specific textures of Y<sub>ν</sub>
- Collider observables like \$\tilde{\chi}\_2^0 → \$\tilde{\chi}\_1^0 \ell\_i \ell\_j\$ or Δm<sub>\tilde{\ell}\$</sub> are suppressed when compared to MSSM + type I seesaw due to small Y<sub>ν</sub>
- Non-degenerate singlets don't change the behaviour of observables dominated by Z-mediated contributions
- Higgs mass around 125 GeV can be accomodated with large  $A_0$ and moderately large tan  $\beta$  and singlets don't contribute to Higgs mass because of the smallness of  $Y_{\nu}$

・ ロ ト ・ 雪 ト ・ ヨ ト ・ ヨ ト

## Conclusion

- $cLFV \Rightarrow Clear signal of new physics$
- Enhancement from the inverse seesaw ⇒ Put constraints on Yukawa couplings
- Most constraining observable:  $\mu e$  conversion
- If nothing is detected ⇒ Strong constraints on the SUSY inverse seesaw, maybe exclusion if coupled with LHC (absence of) results on SUSY
- If cLFV is detected in the predicted range ⇒ Interplay of cLFV with other observables will help to disentangle the type of neutrino mass generation mechanism and shed light on the new physics



э

▲日 ▶ ▲圖 ▶ ▲ 臣 ▶ ▲ 臣 ▶ ...

æ

#### This is a work in progress but we have very good collaborators.



Thank you

## Three Seesaw Mechanisms

- Three classes of seesaw models at tree level ⇒ Three kinds of heavy fields
  - type I: RH neutrinos, SM gauge singlets
  - type II: scalar triplets
  - type III: fermionic triplets



▲ロト ▲御 ト ▲ 陸 ト ▲ 陸 ト 一 陸 …

・ロト ・ 戸 ・ ・ ヨ ・ ・ 日 ・

## Approche effective des mécanismes de seesaw

- Conservation accidentelle du nombre leptonique dans le MS (provient du groupe de jauge, du contenu en champ et de la renormalisabilité)
- Nécessité de violer L pour générer des masses de neutrino ⇒ Opérateurs effectifs non-renormalisables
- Unique opérateur de dimension 5 pour tous les seesaw
   → Viole le nombre leptonique ⇒ neutrinos de Majorana

$$\delta \mathcal{L}^{d=5} = \frac{1}{2} c_{ij} \frac{(H \cdot L_i)^{\dagger} (H \cdot L_j)}{\Lambda} + \text{h.c.}$$

- Pour distinguer les différents mécanismes de seesaw
  - Production directe des états lourds (LHC, ILC)
  - Effets des opérateurs de dimension 6  $\rightarrow$  LFV

# Higgs-mediated cLFV contribution through slepton mixing











Motivations Inverse Seesaw Supersymmetry Charged LFV Results

• Soft SUSY breaking lagrangian :

$$\begin{aligned} -\mathcal{L}_{\text{soft}} &= -\mathcal{L}_{\text{soft}}^{\text{MSSM}} + m_{\widetilde{N}}^2 \widetilde{N}_i^{\dagger} \widetilde{N}_i + m_X^2 \widetilde{X}_i^{\dagger} \widetilde{X}_i + \left( A_{\nu} Y_{\nu}^{ij} \varepsilon_{ab} \widetilde{L}_i^a \widetilde{N}_j H_u^b \right. \\ &+ B_{M_{R_i}} \widetilde{N}_i \widetilde{X}_i + \frac{1}{2} B_{\mu_{X_i}} \widetilde{X}_i \widetilde{X}_i + \text{h.c.} \right) \end{aligned}$$

RGE corrections to the left-handed slepton soft-breaking masses
 :

$$(\Delta m_{\tilde{L}}^2)_{ij} \simeq -\frac{1}{8\pi^2} (3m_0^2 + A_0^2) (Y_{\nu}^{\dagger} L Y_{\nu})_{ij}, \quad L = \ln \frac{M_{GUT}}{M_R}$$
  
=  $\xi (Y_{\nu}^{\dagger} Y_{\nu})_{ij}$ 

• LFV coefficient :

$$\kappa_{ij}^{E} = \frac{\epsilon_{2ij}^{\text{tot}}(Y_{\nu}^{\dagger}Y_{\nu})_{ij}}{\left[1 + \left(\epsilon_{1} + \epsilon_{2ii}^{\text{tot}}(Y_{\nu}^{\dagger}Y_{\nu})_{ii}\right)\tan\beta\right]^{2}}$$

Motivations

Supersymmetry

Results

• Branching ratios:

$$\mathsf{Br}(\tau \to 3\mu) \approx \frac{G_F^2 m_\mu^2 m_\tau^7 \tau_\tau}{768 \, \pi^3 M_A^4} |\kappa_{\tau\mu}^E|^2 \tan^6 \beta$$

$$\mathsf{Br}(B_s \to \ell_i \ell_j) = \frac{G_F^4 M_W^4}{8 \, \pi^5} \left| V_{tb}^* V_{ts} \right|^2 M_{B_s}^5 f_{B_s}^2 \, \tau_{B_s} \left( \frac{m_b}{m_b + m_s} \right)^2$$

$$imes ~ \sqrt{ \left[ 1 - rac{(m_{\ell_i} + m_{\ell_j})^2}{M_{B_s}^2} 
ight] \left[ 1 - rac{(m_{\ell_i} - m_{\ell_j})^2}{M_{B_s}^2} 
ight] }$$

$$\times \left\{ \left( 1 - \frac{(m_{\ell_i} + m_{\ell_j})^2}{M_{B_s}^2} \right) |c_S^{ij}|^2 + \left( 1 - \frac{(m_{\ell_i} - m_{\ell_j})^2}{M_{B_s}^2} \right) |c_P^{ij}|^2 \right\}$$

$$c_S^{\mu\tau} = c_P^{\mu\tau} \approx \frac{8\pi^2 m_\tau m_t^2}{M_W^2} \frac{\epsilon_Y \kappa_{\tau\mu}^E \tan^4 \beta}{\left[1 + (\epsilon_0 + \epsilon_Y Y_t^2) \tan \beta\right] \left[1 + \epsilon_0 \tan \beta\right]} \frac{1}{M_A^2} \bigvee$$

$$\begin{aligned} \frac{\mathsf{Br}(\tau \to \mu\eta)}{\mathsf{Br}(\tau \to 3\mu)} &\simeq & 36\,\pi^2 \left(\frac{f_\eta^8\,m_\eta^2}{m_\mu\,m_\tau^2}\right)^2 (1-x_\eta)^2 \left[\xi_s + \frac{\xi_b}{3} \left(1+\sqrt{2}\frac{f_\eta^0}{f_\eta^8}\right)\right]^2 \\ \frac{\mathsf{Br}(\tau \to \mu\eta')}{\mathsf{Br}(\tau \to \mu\eta)} &\simeq & \frac{2}{9} \left(\frac{f_{\eta'}^0}{f_\eta^8}\right)^2 \frac{m_{\eta'}^4}{m_\eta^4} \left(\frac{1-x_{\eta'}}{1-x_\eta}\right)^2 \left[\frac{1+\frac{3}{\sqrt{2}}\frac{f_{\eta'}^8}{f_{\eta'}^6}\left(\frac{\xi_s}{\xi_b} + \frac{1}{3}\right)}{\frac{\xi_s}{\xi_b} + \frac{1}{3} + \frac{\sqrt{2}}{3}\frac{f_{\eta}^0}{f_\eta^8}}\right]^2 \\ \frac{\mathsf{Br}(\tau \to \mu\pi)}{\mathsf{Br}(\tau \to \mu\eta)} &\simeq & \frac{4}{3} \left(\frac{f_\pi}{f_\eta^8}\right)^2 \frac{m_\pi^4}{m_\eta^4} (1-x_\eta)^{-2} \left[\frac{\frac{\xi_d}{\xi_b}\frac{1}{1+z} + \frac{1}{2}(1+\frac{\xi_s}{\xi_b})\frac{1-z}{1+z}}{\frac{\xi_s}{\xi_b} + \frac{1}{3} + \frac{\sqrt{2}}{3}\frac{f_\eta^0}{f_\eta^8}}\right]^2 \end{aligned}$$

$$\mathsf{Br}(H_k \to \mu \tau) = \tan^2 \beta \; (|\kappa_{\tau \mu}^E|^2) \; C_\Phi \; \mathsf{Br}(H_k \to \tau \tau)$$

$$C_{h} = \left[\frac{\cos(\beta - \alpha)}{\sin\alpha}\right]^{2}, \quad C_{H} = \left[\frac{\sin(\beta - \alpha)}{\cos\alpha}\right]^{2}, \quad C_{A} = 1$$

|--|

10,10,10,12,12, 2, 3000