


B_s Oscillations and Rare Decays at the Tevatron




GDR SUSY Workshop
28. April 2008 Strasbourg

Ralf Bernhard
University of Freiburg



Outline

- ✗ Motivation
- ✗ Measurement of B_s mixing: Δm_s
- ✗ Measurement of $\Delta\Gamma$
- ✗ CP Violation in the B_s system
- ✗ Rare leptonic decays
- ✗ $b \rightarrow l \bar{l} X$ transitions
- ✗ Summary





Motivation

Why huge matter-antimatter asymmetry in the universe?


Why B Physics? - It's got it all!

- × Electroweak symmetry breaking → determines flavor structure: CKM matrix, CP violation, FCNC's
- × QCD Modeling: production, spectroscopy, masses, lifetimes, decays → Challenges lattice gauge, Heavy Quark Effective Theory, strong symmetries
- × Search for new physics → rare decays and


Why at the Tevatron?



→ Complementary to $\Upsilon(4S)$ B factories




Tevatron




Tevatron continues to perform well

- Over 3.9fb^{-1} delivered and 3.4fb^{-1} recorded by each experiment, 2.8fb^{-1} analysed
- Peak luminosities of $\sim 3 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$
→ up to 10 interactions




Detectors



Relevant for B physics:

DØ Tracker: excellent coverage
& vertexing

- ✗ Silicon & scintillating fiber
- ✗ Small radii, but extending to $|\eta| < 2$
- ✗ New Layer 0 silicon on beam pipe in 2006, improving impact para. resol.
- ✗ Triggered muon coverage: $|\eta| < 2$
- ✗ E.g. triggers: dimuons, single muons, track displacement @ L2



CDF Tracker: excellent mass resolution
& vertexing

- ✗ Silicon, Layer 00
- ✗ Large radii drift chamber, many hits, excellent momentum resolution
- ✗ dE/dx (and TOF): particle id
- ✗ Triggered muon coverage: $|\eta| < 1$
- ✗ E.g. triggers: dimuons, lepton + displ. track, two displaced tracks

Mixing and Oscillations

Weak Eigenstates propagate according to Schrodinger:

$$i \frac{d}{dt} \begin{pmatrix} B^0 \\ \bar{B}^0 \end{pmatrix} = \begin{pmatrix} M - \frac{i\Gamma}{2} & M_{12} - \frac{i\Gamma_{12}}{2} \\ M_{12}^* - \frac{i\Gamma_{12}^*}{2} & M - \frac{i\Gamma}{2} \end{pmatrix} \begin{pmatrix} B^0 \\ \bar{B}^0 \end{pmatrix}$$

Diagonalize


CP Eigenstates: $|B^{\text{odd}}\rangle = |B^0\rangle + |\bar{B}^0\rangle$ $|B^{\text{even}}\rangle = |B^0\rangle - |\bar{B}^0\rangle$

Mass Eigenstates: $|B^H\rangle = p|B^0\rangle + q|\bar{B}^0\rangle$ $|B^L\rangle = p|B^0\rangle - q|\bar{B}^0\rangle$

Heavy *Light*

Mass Difference:

$$\Delta m = M_H - M_L \sim 2|M_{12}|$$



Conversion of matter to anti-matter

Mixing and Oscillations

Weak Eigenstates propagate according to Schrodinger:

$$i \frac{d}{dt} \begin{pmatrix} B^0 \\ \bar{B}^0 \end{pmatrix} = \begin{pmatrix} M - \frac{i\Gamma}{2} & M_{12} - \frac{i\Gamma_{12}}{2} \\ M_{12}^* - \frac{i\Gamma_{12}^*}{2} & M - \frac{i\Gamma}{2} \end{pmatrix} \begin{pmatrix} B^0 \\ \bar{B}^0 \end{pmatrix}$$

Diagonalize

CP Eigenstates: $|B^{\text{odd}}\rangle = |B^0\rangle + |\bar{B}^0\rangle$ $|B^{\text{even}}\rangle = |B^0\rangle - |\bar{B}^0\rangle$

Mass Eigenstates: $|B^H\rangle = p|B^0\rangle + q|\bar{B}^0\rangle$ $|B^L\rangle = p|B^0\rangle - q|\bar{B}^0\rangle$

Heavy *Light*

For the B_s^0 meson:

$$\Delta m_s = M_H - M_L \sim 2|M_{12}|$$

$$\Delta\Gamma_s^{CP} = \Gamma_{\text{even}} - \Gamma_{\text{odd}} \sim 2|\Gamma_{12}|$$

$$\Delta\Gamma_s = \Gamma_L - \Gamma_H \sim 2|\Gamma_{12}| \cos\phi_s$$

$$\Gamma_s = \frac{\Gamma_L + \Gamma_H}{2} ; \quad \bar{\tau}_s = \frac{1}{\Gamma_s} \quad \phi_s^{\text{SM}} = \arg \left[-\frac{M_{12}}{\Gamma_{12}} \right] \sim 0.004 \text{ in SM}$$

Tiny for B_d^0 meson, but
not for B_s^0 ! eigenstates propagate
with different lifetimes!

Mixing and Oscillations

Weak Eigenstates propagate according to Schrodinger:

All different!

$$i \frac{d}{dt} \begin{pmatrix} B^0 \\ \bar{B}^0 \end{pmatrix} = \begin{pmatrix} M - \frac{i\Gamma}{2} & M_{12} - \frac{i\Gamma_{12}}{2} \\ M_{12}^* - \frac{i\Gamma_{12}^*}{2} & M - \frac{i\Gamma}{2} \end{pmatrix} \begin{pmatrix} B^0 \\ \bar{B}^0 \end{pmatrix}$$

Diagonalize Probe entire matrix

CP Eigenstates: $|B^{\text{odd}}\rangle = |B^0\rangle + |\bar{B}^0\rangle$ $|B^{\text{even}}\rangle = |B^0\rangle - |\bar{B}^0\rangle$

Mass Eigenstates: $|B^{\text{H}}\rangle = p|B^0\rangle + q|\bar{B}^0\rangle$ $|B^{\text{L}}\rangle = p|B^0\rangle - q|\bar{B}^0\rangle$

Heavy *Light*

For the B_s^0 meson: Whole new window for New physics

$$\Delta m_s = M_H - M_L \sim 2|M_{12}|$$
Sensitive for NP

$$\Delta \Gamma_s^{CP} = \Gamma_{\text{even}} - \Gamma_{\text{odd}} \sim 2|\Gamma_{12}|$$
Not sensitive for NP

$$\Delta \Gamma_s = \Gamma_L - \Gamma_H \sim 2|\Gamma_{12}| \cos \phi_s$$
Very sensitive for NP


$$\Gamma_s = \frac{\Gamma_L + \Gamma_H}{2}; \quad \bar{\tau}_s = \frac{1}{\Gamma_s} \quad \phi_s^{\text{SM}} = \arg \left[-\frac{M_{12}}{\Gamma_{12}} \right] \sim 0.004 \text{ in SM}$$

Frequency of Oscillations

Decays: $B_s^0 \rightarrow \pi^+ D_s^- (\phi\pi^-)$
 $B_s^0 \rightarrow l D_s X, \dots$


measure B_s decay length

$$\bar{B}_s^0 = \left| \begin{array}{c} b \\ \bar{s} \end{array} \right\rangle$$
$$B_s^0 = \left| \begin{array}{c} \bar{b} \\ s \end{array} \right\rangle$$




Decay channels

hadronic



semileptonic



- all decay particles reconstructable
→ better time resolution
- low event rate
- higher combinatoric


more sensitive at higher Δm_s

- high event rate
- ν_μ momentum not measurable
→ sensitivity proper time limited by momentum measurement


more sensitive at lower Δm_s

CDF Signal Selection

$B_s \rightarrow \mu^+ D_s^- (\phi\pi) X$





$B_s \rightarrow \pi^+ D_s^- (\phi\pi)$




DØ Signal Selection

$$B_s \rightarrow \pi^+ D_s^- (\phi\pi) X$$



Only 250 reconstructed and tagged hadronic events (CDF ~500)



Decay channels

CDF (data sample size: $\int L dt = 1 \text{ fb}^{-1}$):


channel	candidates
$\bar{B}_s \rightarrow l D_s X$	61500
$\bar{B}_s \rightarrow \pi^- D_s^+ (\phi \pi^+)$	2000
$\bar{B}_s \rightarrow \pi^- D_s^+ (K^*(892)^0 K^+)$	1400
$\bar{B}_s \rightarrow \pi^- D_s^+ (\pi^+ \pi^- \pi^+)$	700
$\bar{B}_s \rightarrow \pi^- \pi^+ \pi^- D_s^+ (\phi \pi^+)$	700
$\bar{B}_s \rightarrow \pi^- \pi^+ \pi^- D_s^+ (K^*(892)^0 K^+)$	600
$\bar{B}_s \rightarrow \pi^- \pi^+ \pi^- D_s^+ (\pi^+ \pi^- \pi^+)$	200
partially reconstructed	3100

DØ (bigger dataset also includes resolution improvement through Layer0):

channel	candidates	improvements
$B_s \rightarrow \mu^+ D_s^- (\phi \pi) X$	44777 ± 415	data: $1.3 \text{ fb}^{-1} \rightarrow 2.4 \text{ fb}^{-1}$
$B_s \rightarrow e^+ D_s^- (\phi \pi) X$	1663 ± 102	data: $1.3 \text{ fb}^{-1} \rightarrow 2.4 \text{ fb}^{-1}$
$B_s \rightarrow \pi^+ D_s^- (\phi \pi) X$	249 ± 17	new channel
$B_s \rightarrow \mu^+ D_s^- (K^{*0} K^-) X$	18098 ± 903	data: $1.3 \text{ fb}^{-1} \rightarrow 2.4 \text{ fb}^{-1}$

Amplitude scan


$$P(\Delta m_s) \approx 1 \pm \cos \Delta m_s t \Rightarrow P(A) \approx 1 \pm A \cos \Delta m_s t$$




Motivation is to
simplify combining
results with other
experiments

$$\text{Prob}^{unmix,mix}(x) = \frac{K}{2c\tau_{B_s}} \exp\left(-\frac{Kx}{c\tau_{B_s}}\right) \left(1 \pm (1 - 2\eta) \frac{A}{\text{Amplitude}} \cos(\Delta m_s Kx/c)\right)$$

B_s : Δm_s , Frequency of Osci.



$$\Delta m_s = 18.56 \pm 0.87 \text{ ps}^{-1}$$



Power of hadronic $B_s^0 \rightarrow D_s \pi(X)$ decay mode
& two-displace track trigger

CDF, $> 5\sigma$ significance

$$\Delta m_s = 17.77 \pm 0.12 \text{ ps}^{-1}$$

CP Violation

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

In the SM CP violation occurs in only one place:
complex phases in unitary CKM matrix; NP, plenty of places!!!


e.g. 43 in MSSM

CP Violation

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

In the SM CP violation occurs in only one place:
complex phases in unitary CKM matrix; NP, plenty of places!!!

B_d unitarity condition $V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0$





CP Violation

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

In the SM CP violation occurs in only one place:
complex phases in unitary CKM matrix; NP, plenty of places!!!


B_d unitarity condition

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0$$



B_s : Δm_s , Frequency of Osci

$$\frac{\Delta m_s}{\Delta m_d} = V_{td}$$



Not much room for New Physics!

Measurement of $\Delta\Gamma$

First assume no CP violation
in B_s mixing, $\Phi_s=0$

- $B_s \rightarrow D_s^{(*)} D_s^{(*)}$

CP and mass eigenstates are the same

- Three channels

- [$D_s D_s$ (PP), $D_s^* D_s$ (VP), $D_s^* D_s^*$ (VV)]

- Heavy quark limit + factorization

- $B_s^{\text{odd}} \rightarrow D_s^* D_s$ is forbidden

- $D_s^* D_s^*$ in S-wave

- $\Rightarrow Ds^{(*)} Ds^{(*)}$ pure CP even

$$BF(B_s \rightarrow D_s^{(*)} D_s^{(*)}) = \left(\frac{\Delta\Gamma_{CP}}{2\Gamma} \right) \left(1 + O\left(\frac{\Delta\Gamma}{\Gamma}\right) \right)$$

- Flavor specific B_s lifetime

- Flavor specific decays carry equal amounts of B_H and B_L


- Get flavor specific lifetime if FS data with is fit w/ single exponential

$$e^{-t/\tau_{FS}} = \frac{1}{2} \cdot \left(e^{-t/\tau_H} + e^{-t/\tau_L} \right)$$



- $B_s \rightarrow J/\psi \phi: P \rightarrow VV$

- Even and odd paths distinguishable with angular analysis of final state particles

$\Delta\Gamma$ from $B_s \rightarrow D_s^{(*)} D_s^{(*)}$




$$N(D_s^{(*)} D_s^{(*)}) = 13.4 \begin{array}{l} +6.6 \\ -6.0 \end{array}$$




- Trigger on muon from semileptonic D_s decay
- Ignore any photons
- Look for correlated production of $D_s \rightarrow \phi\pi$ and $D_s \rightarrow \phi\mu$

$\Delta\Gamma$ from $B_s \rightarrow D_s^{(*)} D_s^{(*)}$



$$N(D_s^{(*)} D_s^{(*)}) = 13.4 \begin{array}{l} +6.6 \\ -6.0 \end{array}$$



$$BF(B_s \rightarrow D_s^{(*)} D_s^{(*)}) = 0.039 \begin{array}{l} +0.019 \\ -0.017 \end{array} \begin{array}{l} +0.016 \\ -0.015 \end{array}$$

$$\frac{\Delta\Gamma_{CP}}{\Gamma} = 0.079 \begin{array}{l} +0.038 \\ -0.035 \end{array} \begin{array}{l} +0.031 \\ -0.030 \end{array}$$

B_s Flavor Specific Lifetime



Know flavor at time of decay from charge of decay product

hep-ph/0201071

$$|B_s \rightarrow D_s^- \mu^+ \nu\rangle = \frac{1}{\sqrt{2}} \left(|B_H\rangle + |B_L\rangle \right)$$

50% CP-even, 50% CP-odd at time t_0



$$\tau_{FS} = \frac{1}{\bar{\Gamma}_s} \frac{1+y^2}{1-y^2} \text{ with } y = \frac{\Delta\Gamma}{2\Gamma}$$



$$\tau_{FS}(B_s) = 1.381 \pm 0.055^{+0.052}_{-0.046} \text{ ps}$$

B_s Flavor Specific Lifetime


$$|B_s \rightarrow D_s^- \pi^+(\pi^0)\rangle = \frac{1}{\sqrt{2}} \left(|B_H\rangle + |B_L\rangle \right)$$



$B_s \rightarrow J/\psi \phi$


- Heavy (H , CP-odd) and Light (L , CP-even) B_s states

$$\Delta\Gamma_s = \Gamma_L - \Gamma_H; \quad \Gamma_s = (\Gamma_L + \Gamma_H)/2; \quad \bar{\tau}_s = \frac{1}{\Gamma_s}$$




Not “flavor specific”,
predicted to be more
CP even than odd

- Decays into two vector mesons that are either CP-odd ($L=1$) or CP-even ($L=0,2$)
- Time-dependent angular distributions allow separation of components
- Simultaneous fit to lifetime and three angles




$B_s \rightarrow J/\psi \phi$


arXiv:0802.2255




even /
light



arXiv:0712.2242




odd /
heavy



$\Delta\Gamma$ and Γ_s

First assume no CP violation
in B_s mixing, $\Phi_s=0$




$$\Delta\Gamma_s = 0.076^{+0.059}_{-0.063} \pm 0.006 \text{ ps}^{-1}$$

$$\bar{\tau}_s = 1.52 \pm 0.04 \pm 0.02 \text{ ps}$$

$$\bar{\tau}_s = \frac{1}{\Gamma_s} = \frac{2}{\Gamma_H + \Gamma_L}$$

c.f. $\Delta\Gamma_s^{SM,pred} = 0.088 \pm 0.017 \text{ ps}^{-1}$ (hep-ph/0612167)



$$\Delta\Gamma_s = 0.14 \pm 0.07 \text{ ps}^{-1}$$



$$\bar{\tau}_s = 1.53 \pm 0.05 \pm 0.01 \text{ ps}$$

CP Violation

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

In the SM CP violation occurs in only one place:
complex phases in unitary CKM matrix; NP, plenty of places!!!

B_d unitarity condition $V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0$



CP Violation in B_s System

Explore new part of matrix


$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

In the SM CP violation occurs in only one place:
complex phases in unitary CKM matrix; NP, plenty of places!!!

B_s unitarity
condition

$$V_{us}V_{ub}^* + V_{cs}V_{cb}^* + V_{ts}V_{tb}^* = 0$$

"Squashed"
Triangle



$$\begin{aligned} \beta_s^{SM} &= \arg[-V_{ts}V_{tb}^*/V_{cs}V_{cb}^*] \\ &\approx 0.02 \\ &\text{Tiny!} \end{aligned}$$

CP Violation in B_s System

Explore new part of matrix


$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

In the SM CP violation occurs in only one place:
complex phases in unitary CKM matrix; NP, plenty of places!!!


B_s unitarity
condition

$$V_{us}V_{ub}^* + V_{cs}V_{cb}^* + V_{ts}V_{tb}^* = 0$$

Golden mode,
Tevatron



"Squashed"
Triangle



CP violation through
interference of diagrams
with and w/o mixing

CP Violation in B_s System

- How could New Physics affect these phases

$$2\beta_s^{SM} = 2 \arg[-V_{ts}V_{tb}^*/V_{cs}V_{cb}^*] \rightarrow 2\beta_s^{SM} - \phi_s^{NP}$$

~0.04

$$\phi_s^{SM} = \arg[-M_{12}/\Gamma_{12}] \rightarrow \phi_s^{SM} + \phi_s^{NP}$$

~0.004

Subtracts from one,
adds to other

- Both CDF and DØ measure/observe the phase responsible for CP violation in $B_s \rightarrow J/\psi \phi$ decays


$$\phi_s = -2\beta_s \approx \phi_s^{NP}$$

DØ CDF If large

- Use flavor tagging to identify initial flavor of B_s or Anti B_s in $J/\psi \phi$ decays (and know value of Δm_s)

CP Violation in $B_s \rightarrow J/\psi \phi$

- Even without initial state flavor tagging, have sensitivity to ϕ_s



DØ

PRL 98 , 121801 (2007)


$$\phi_s = -0.79 \pm 0.56^{+0.14}_{-0.01}$$

For one of the ambiguities

- But 4-fold ambiguity, reduce to 2-fold with flavor tagging...

CP Violation in $B_s \rightarrow J/\psi \phi$

- Now use initial state flavor tagging



- Ambiguities


$$2\beta_s^{J/\psi\phi} \rightarrow \pi - 2\beta_s^{J/\psi\phi} \quad \Delta\Gamma_s \rightarrow -\Delta\Gamma_s$$

Strong phases
(relative phases between polarization amplitudes)


$$\delta_{||} \rightarrow 2\pi - \delta_{||} \quad \delta_{\perp} \rightarrow \pi - \delta_{\perp}$$

CP Violation in $B_s \rightarrow J/\psi \phi$

- Now using initial state flavor tagging, constrain strong phases



Confidence regions underestimated using $2\Delta\log L$



Standard Model Probability = 6.6%, ~1.8 σ

- Ambiguities

$$2\beta_s^{J/\psi\phi} \rightarrow \pi - 2\beta_s^{J/\psi\phi} \quad \Delta\Gamma_s \rightarrow -\Delta\Gamma_s$$

Constrain based on B_d^0 observations


$$\delta_{||} \rightarrow 2\pi - \delta_{||}$$

$$\delta_{\perp} \rightarrow \pi - \delta_{\perp}$$

ϕ_s Measurement

In Standard Model: $\phi_s \approx \arg(-V_{ts}) \approx 0.004$ rad.

Observables: **Semileptonic asymmetries**, interference in decays to CP eigenstates



$$\frac{N(D_s\mu^+) - N(D_s\mu^-)}{N(D_s\mu^+) + N(D_s\mu^-)} = A_{SL}(\text{untagged}) \approx \frac{\Delta\Gamma}{\Delta m} \tan\phi$$

$$\frac{N(\mu^+\mu^+) - N(\mu^-\mu^-)}{N(\mu^+\mu^+) + N(\mu^-\mu^-)} = A_{SL}(\text>tagged) = 2A_{SL}(\text{untagged})$$

Same sign Dimuons

$$\frac{N(\mu^+ \mu^+) - N(\mu^- \mu^-)}{N(\mu^+ \mu^+) + N(\mu^- \mu^-)} = A_{SL}(\text{tagged}) = 2A_{SL}(\text{un>tagged})$$

$N(\text{same sign}) \approx 310\text{K}$

$$A_{SL} = -0.0092 \pm 0.0044 \pm 0.0032$$

$\sim 60/40$ mix of B_d and B_s

$Z \sim 2\chi$

$$A_{SL} = A_{SL}(B_d) + \frac{f_s Z_s}{f_d Z_d} A_{SL}(B_s)$$









$A_{SL}(B_d) = -0.0047 \pm 0.0046$ (HFAG, B-factories)

$$A_{SL}(B_s, \mu\mu) = -0.0064 \pm 0.0101$$

Regular flipping of polarity of solenoid (tracking) and toroid (muons) magnets essential for controlling systematic uncertainties

Exclusive $B_s \rightarrow D_s^\pm \mu \nu$ Results

Exclusive $D_s \mu$



CDF: 1.6 fb^{-1} , CDF Note 9015

$$\frac{N(D_s \mu^+) - N(D_s \mu^-)}{N(D_s \mu^+) + N(D_s \mu^-)} = A_{SL}(\text{untagged}) \approx \frac{\Delta \Gamma}{\Delta m} \tan \phi$$

$$A_{SL}(B_s, D_s \mu) = \\ 0.0245 \pm 0.0193 \pm 0.0035$$

DØ Combined:

$$A_{SL}(B_s, \mu\mu + D_s \mu) = \\ 0.0001 \pm 0.0090$$


Using Δm_s from CDF:

$$\Delta \Gamma_s \cdot \tan \phi_s = 0.02 \pm 0.16 \text{ ps}^{-1}$$

New

$$a_{SL}^s = 0.020 \pm 0.021 \pm 0.018$$


ϕ_s Results



Combined with older DØ analysis before flavor tagging

CP Violation in B_s : Combination






- In $B_s \rightarrow J/\psi \phi$ flavor-tagged analyses, in $(\Delta\Gamma_s, \phi_s)$ space CDF has $\sim 1.5\sigma$ deviation from SM, DØ $\sim 1.8\sigma$ deviation, consistent with each other
- Need to be careful, non-parabolic $\log(L)$, multiple correlations
(best is simply more data!!)



UTfit results

UTfit group, arXiv:0803.0659:

Δm_s , A_{SL}^s , $A_{SL}^{\mu\mu}$, τ_{fs}



> 3 σ deviation from SM

- Intriguing, results from DØ, CDF with more data coming soon; CDF+DØ+HFAG also working on combin.

Next talk from Marcella Bona

Direct CP violation in b Hadrons

- Direct (not through mixing) CP violation expected to be large in some b hadron decays, including B mesons and b Baryons
- Measure asymmetry: $f = \text{final state}$

$$A_{CP} = \frac{N(\bar{B} \rightarrow \bar{f}) - N(B \rightarrow f)}{N(\bar{B} \rightarrow \bar{f}) + N(B \rightarrow f)}$$


- CDF: Br 's and asymmetries of two-body charmless states, $B \rightarrow hh'$

CDF Note 9092

$$A_{CP}(\Lambda_b^0 \rightarrow p\pi^-) = 0.03 \pm 0.17 \pm 0.05$$


$$A_{CP}(\Lambda_b^0 \rightarrow pK^-) = 0.37 \pm 0.17 \pm 0.03$$

- Expectations, asymmetry $\sim 30\%$
- First CP asymmetry measurement in b baryon decays



Direct CP violation in b Hadrons

- ✗ DØ: Small (~1%) CP asymmetry expected in SM for $B^+ \rightarrow J/\psi K^+$
- ✗ Again, frequent solenoid and toroid polarity reversals essential to control charge asymmetry systematic uncertainties
- ✗ Correct for K^+/K^- asymmetry
- ✗ <1% precision factor ~2 better than current world average



Accepted by Phys. Rev. Lett.
arXiv:0802.3299


$$A_{CP}(B^+ \rightarrow J/\psi K^+) = +0.0075 \pm 0.0061 \pm 0.0027$$

Purely leptonic B decay

- $B \rightarrow l^+ l^-$ decay is helicity suppressed FCNC
- SM: $\text{BR}(B_s \rightarrow \mu^+ \mu^-) \sim 3.4 \times 10^{-9}$
- depends only on one SM operator in effective Hamiltonian, hadronic uncertainties small
- B_d relative to B_s suppressed by $|V_{td}/V_{ts}|^2 \sim 0.04$ if no additional sources of flavor violation
- reaching SM sensitivity: present limit for $B_s \rightarrow \mu^+ \mu^-$ comes closest to SM value

SM expectations:

	$\text{Br}(B_d \rightarrow l^+ l^-)$	$\text{Br}(B_s \rightarrow l^+ l^-)$
$l = e$	3.4×10^{-15}	8.0×10^{-14}
$l = \mu$	1.0×10^{-10}	3.4×10^{-9}
$l = \tau$	3.1×10^{-8}	7.4×10^{-7}




Current published limits at 95%CL:

	$\text{Br}(B_d \rightarrow l^+ l^-)$	$\text{Br}(B_s \rightarrow l^+ l^-)$
$l = e$	$< 6.1 \cdot 10^{-8}$	$< 5.4 \cdot 10^{-5}$
$l = \mu$	$< 1.8 \cdot 10^{-8}$	$< 5.8 \times 10^{-8}$
$l = \tau$	$< 2.5\%$	$< 5.0\%$



Purely leptonic B decay

- ✖ excellent probe for many new physics models
- ✖ particularly sensitive to models w/ extended Higgs sector
 - ✖ BR grows $\sim \tan^6 \beta$ in MSSM
 - ✖ 2HDM models $\sim \tan^4 \beta$
 - ✖ mSUGRA: BR enhancement correlated with shift of $(g-2)_\mu$
- ✖ also, testing ground for
 - ✖ minimal $SO(10)$ GUT models
 - ✖ R_p violating models, contributions at tree level
 - ✖ (neutralino) dark matter ...

Two-Higgs Doublet models:






R_p violating:



Search Strategy

- ✖ Preselection of Di Muon events
- ✖ Normalization channel $B^+ \rightarrow J/\psi K^+$
- ✖ Background estimation using sidebands
- ✖ Background reduction using a LHR ($D\emptyset$) or NN (CDF)



Limits

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) < \frac{N_{UL}}{N_{B^+}} \cdot \frac{\varepsilon_{\mu^+ \mu^- K}^{B^+}}{\varepsilon_{\mu^+ \mu^-}^{B_s^0}} \cdot \frac{\mathcal{B}(B^\pm \rightarrow J/\psi(\mu^+ \mu^-) K^\pm)}{\frac{f_{b \rightarrow B_s}}{f_{b \rightarrow B_{u,d}}} + R \cdot \frac{\varepsilon_d^{B_d^0}}{\varepsilon_{\mu^+ \mu^-}^{B_s^0}}}$$

Relative Normalization

$\varepsilon_{B^+}/\varepsilon_{B_s}$ relative efficiency of normalization to signal channel

f_s/f_u fragmentation ratio - use world average (3.71) with 15% uncertainty

$\varepsilon_{B_d}/\varepsilon_{B_s}$ relative efficiency for $B_d \rightarrow \mu^+ \mu^-$ versus $B_s \rightarrow \mu^+ \mu^-$ events in B_s search channel (~ 0.95) $R = BR(B_d)/BR(B_s)$ is small due to $|V_{td}/V_{ts}|^2$



at 90% CL

$Br(B_s \rightarrow \mu\mu)$	2 fb^{-1}	7.3×10^{-8}	Prelim. DØ
$Br(B_s \rightarrow \mu\mu)$	2 fb^{-1}	4.7×10^{-7}	Prelim. CDF
$Br(B_s \rightarrow \mu\mu)$	combined	3.6×10^{-8}	HFAG

DØ Note 5344

PRL 100,101802 (2008)

Rare decays constraining NP




Study of $b \rightarrow s l^+ l^-$


- × long-term goal: investigate $b \rightarrow s l^+ l^-$ FCNC transitions in B^+ , B_d and B_s mesons
- × $B^+ \rightarrow l^+ l^- K^+$ and $B_d \rightarrow l^+ l^- K^*$ established at B factories
- × $B_s \rightarrow l^+ l^- \phi$ only accessible at the Tevatron
 - × SM prediction:
 - × short distance BR: $\sim 1.6 \times 10^{-6}$
 - × About 30% uncertainty due to $B \rightarrow \phi$ form factor
- × 2HDM: enhancement possible, depending on parameters for $\tan\beta$ and M_{H^+}

$b \rightarrow s l^+ l^-$ Theory

$\mu\mu$ Mass spectrum



Sensitive to “new Physics” contribution




The forward backward asymmetry is defined as the asymmetry between forward and backward emitted positive leptons in the dilepton rest frame where the direction is relative to the direction of the B meson.


B- $\rightarrow\mu\mu h$ @ CDF

arXiv:0804.3908


CDF Run II Preliminary



CDF Run II Preliminary




CDF Run II Preliminary



Mode	$B^+ \rightarrow \mu^+\mu^-K^+$	$B_d^0 \rightarrow \mu^+\mu^-K^{*0}$	$B_s^0 \rightarrow \mu^+\mu^-\phi$
N_S	90	35	11
N_{BG}	45.3 ± 5.8	16.5 ± 3.6	3.5 ± 1.5
Gaussian significance	4.5	2.9	2.4
$N_{J/\psi h}$	6246 ± 79	2346 ± 48	421 ± 21
$\epsilon_{\mu^+\mu^-h}/\epsilon_{J/\psi h}$	0.71 ± 0.01	0.74 ± 0.02	0.84 ± 0.02
Rel $\mathcal{B} \pm stat \pm syst \times 10^{-3}$	$0.59 \pm 0.15 \pm 0.03$	$0.62 \pm 0.23 \pm 0.07$	$1.24 \pm 0.60 \pm 0.15$
Abs $\mathcal{B} \pm stat \pm syst \times 10^{-6}$	$0.60 \pm 0.15 \pm 0.04$	$0.82 \pm 0.31 \pm 0.10$	$1.16 \pm 0.56 \pm 0.42$
Rel \mathcal{B} 95%CL limit $\times 10^{-3}$	—	—	2.61
Rel \mathcal{B} 90%CL limit $\times 10^{-3}$	—	—	2.30

Branching ratio overview



Need more statistics to measure charge asymmetry
vs. invariant di-lepton mass and to add Bs channel!

Summary

- Δm_s established and well measured at the Tevatron
- B_s system and CP studies opening a powerful new window: possibly already providing hints of new phenomena?
- Limit on rare decay $B_s \rightarrow \mu^+ \mu^-$ is getting more and more stringent, help constrain physics beyond the SM
- Other FCNC ($b \rightarrow s l l$) decays test the SM
- Tevatron doing very well, expect to at least *double* our data-set by the end of running