


Results of Dark Matter searches with Baksan

M.M.Boliev, S.V.Demidov, O.V.Suvorova, S.P.Mikheev,
INR RAS


Exotic Physics with Neutrino Telescopes 2013
3 April 2013



Outline


- I Baksan Underground Scintillator Telescope
- II Event selection
- III Signal simulation
- IV Sun survey by BUST
- V Results
- VI Future plans and conclusions

Baksan Neutrino Observatory



Baksan Underground Scintillator Telescope


General view



- depth: 850 hg/cm^2
- size: $17 \text{ m} \times 17 \text{ m} \times 11 \text{ m}$
- 3150 tanks of size $70 \text{ cm} \times 70 \text{ cm} \times 30 \text{ cm}$
- angular resolution: about 1.5°
- time resolution: 5 ns
- general trigger rate: 17 Hz
- muon fluxes upward/downward ratio: $\sim 10^{-7}$

In operation since 18 December 1978

Baksan Underground Scintillator Telescope



Baksan Underground Scintillator Telescope

Baksan Underground Scintillator Telescope



Baksan Underground Scintillator Telescope

Time-of-flight method and event selection

- ▶ time resolution is about 5 ns (Yu. Andreyev et al., 1979, S.P.Mikheev, 1984)
- ▶ probability of imitation of “wrong” direction is considerably diminished if more than two planes involved
- ▶ two special triggers for upward muons: **T1** - for zenith angle range $95^\circ \div 180^\circ$, **T2** - for almost horizontal events: $80^\circ \div 100^\circ$

Trigger T1

- ▶ ≥ 3 scintillator planes
- ▶ ≥ 2 negative Δt
- ▶ ≤ 3 external scintillator planes

Trigger T2

- ▶ = 2 vertical scintillator planes
- ▶ = 0 horizontal scintillator planes
- ▶ $\Delta t \geq 30$ ns ($\text{pathlength} \geq 10$ m)

trigger rate 0.02 Hz (1800 events per day)

Event selection: additional cuts

Cuts Level 1

- ▶ Only one reconstructed track with $\beta < 0$
- ▶ Enter point should be below exit point
- ▶ For T2: exclude events with $0 < \phi < 180$ with respect to least shallow depth

Cuts Level 2



- ▶ Only through going tracks (no stopping muons or neutrino interactions inside)
- ▶ Geometrical cuts to exclude events close to plane edge (1.5 m)
- ▶ Muon range inside detector $> 500 \text{ g/cm}^2$ (excluded muons with $E_\mu < 1 \text{ GeV}$)
- ▶ $-1.3 < 1/\beta < -0.7$ (from MC: 95% of upward-going events)

December 1978 – November 2009; livetime 24.12 yrs;

1700 muons after Cuts Level 1; 1255 muons after Cuts Level 2

MC simulation and reconstruction

O.Suvorova, M.Boliev, S.Mikheev et al., 1996



Muon energy threshold $E_{\mu} > 1$ GeV

Efficiency of registration upward-going muon with $E > E_{th}$ is about 0.3

Signal simulation

- ▶ (Anti)Neutrinos are produced in the result of DM annihilations produced in the center of the Sun
- ▶ Propagation of neutrinos in the Sun and Earth
- ▶ Expected muon flux from dark matter annihilation in the Sun

$$\Phi_\mu = \frac{\Gamma_A}{4\pi R^2} \times \sum_{\nu_j, \bar{\nu}_j} \int_{E_{th}}^{m_{DM}} dE_{\nu_j} P(E_{\nu_j}, E_{th}) \frac{dN_{\nu_j}}{dE_{\nu_j}}$$

$P(E_{\nu_j}, E_{th})$ - probability of neutrino-muon conversion,





- ▶ $\frac{dN_{\nu_j}}{dE_{\nu_j}}$ - spectra of neutrino at production point - depend on annihilation channel: $\chi\bar{\chi} \rightarrow \dots$
- ▶ Benchmark channels: $b\bar{b}$ (soft spectrum), W^+W^- and $\tau^+\tau^-$ (hard spectrum)

Signal simulation: overview and parameters


- ▶ We use our C program; compare results with WIMPsim
(M.Blennow, J.Edsjo, T.Ohansson, 2008)
- ▶ Initial neutrino spectra at the center of the Sun (M.Cirelli,
N.Fornengo et al., Nucl.Phys. B727 (2005) 99)
- ▶ Annihilation point near the center of the Sun
- ▶ Neutrino oscillations, 3×3 scheme ($\Delta m_{21} = 7.63 \cdot 10^{-5}$ eV 2 ,
 $|\Delta m_{31}| = 2.55 \cdot 10^{-3}$ eV 2 , $\delta_{CP} = 0$, $\sin^2 \theta_{12} = 0.32$, $\sin^2 \theta_{23} = 0.49$,
 $\sin^2 \theta_{13} = 0.026$, D.V. Forero, M. Tortola, J.W.F. Valle, arXiv:1205.4018)
- ▶ Matter effects: solar model, J.N.Bahcall, A.M.Serenelli,, S.Basu
(2005)
- ▶ NC and CC interactions (including τ -mass effects) in the Sun and
the Earth: change in neutrino fluxes and spectra
- ▶ ν_τ regeneration: $\nu_\tau \rightarrow \tau^- + \dots$, $\tau^- \rightarrow \nu_\tau, \bar{\nu}_e, \bar{\nu}_\mu + \dots$ - secondary
neutrinos

Comparison with WIMPSim: ν_μ spectra at 1 a.u.


For the same initial neutrino spectra



Muon flux calculation and cone half-angle




$b\bar{b}, m_{DM} = 90$ GeV



$\tau^+\tau^-, m_{DM} = 90$ GeV

- ▶ Muons are produced in CC interactions
- ▶ Mean muon energy losses in rock (D.E.Groom, N.V.Mokhov, S.I.Striganov, 2001)
$$\langle \frac{dE}{dx} \rangle = -(\alpha(E) + \beta(E)E)\rho$$
- ▶ Multiple Coulomb scattering
- ▶ 90% of signal events inside cone half-angle for all annihilation channels; were chosen before looking at data.




$b\bar{b}$
 $\tau^+\tau^-$
 W^+W^-


Upward going muons:

December 1978 - November 2009; livetime 24.12 yrs

Event rate



Muon distribution with respect to position of the Sun




About 50 events per year

Direction to the Sun corresponds to $\cos \Psi_{\mu-\text{Sun}} = 1$


Data and expected background

December 1978 - November 2009; livetime 24.12 yrs

Sun below horizon




Number of signal and background events inside cone half-angle $\Psi_{\mu-\text{Sun}}$



Background – from data with shifted position of the Sun

Upper limits on muon fluxes from DM annihilations

$$\Phi_{\mu}^{lim} = \frac{N^{90}}{0.9 \times S_{eff} \times T}; \quad S_{eff}(E_{th}) = \frac{\int dE d\theta S(E, \theta) \times \epsilon(E_{th}, E, \theta) \times \Phi_{\mu}(E, \theta)}{\int dE d\theta \Phi(E, \theta)}$$



Recalculation to upper limits on SD

G. Wikstrom, J. Edsjo, 2009


- ▶ Firstly, we recalculate $\Phi_\mu \rightarrow \Gamma_A$
- ▶ In equilibrium between capture and annihilation processes:
 $\Gamma_A = C_{DM}/2$
- ▶ Capture rate is determined by the SI and SD elastic cross section of DM particles on nucleons
- ▶ Recalculation $\Gamma_A \rightarrow \sigma_p^{SD}, \sigma_p^{SI}$ (Olga Suvorova, S.D., 2010)

$$\Gamma_A = \Gamma_A^{SD} + \Gamma_A^{SI},$$

$$\frac{\sigma_p^{SD}}{\Gamma_A^{SD}} \cdot \Gamma_A^{Upp.Lim.} = \sigma_p^{SD,Upp.Lim.}, \quad \frac{\sigma_p^{SI}}{\Gamma_A^{SI}} \cdot \Gamma_A^{Upp.Lim.} = \sigma_p^{SI,Upp.Lim.}$$

- ▶ Upper limits on SD cross sections are strong - a lot of hydrogen in the Sun

Upper limits on SD elastic cross section



Future plans: optimization of analysis

- ▶ present analysis uses cone half-angle Ψ which contains 90% of signal events
- ▶ we can optimize our search with respect to value of cone half-angle (Hill, Rawlins, 2003)

$$\text{sensitivity} = \frac{\bar{N}^{90}(\Psi)}{x(\Psi) \times S_{\text{eff}}(x) \times T},$$

where $x(\Psi)$ is a fraction of event inside cone half-angle Ψ ,
 \bar{N}^{90} - mean expected upper limit

- ▶ still in progress

Summary

- ▶ Analysis of upward-going muon data collected for 24.11 years of livetime by neutrino experiment at Baksan Underground Scintillator Telescope has been performed
- ▶ No significant excess was found in search for muon signal from dark matter annihilations in the Sun
- ▶ New limits on muon flux, annihilation rate, SD and SI elastic cross sections
- ▶ Mild changes can be expected from the results of optimization

Conclusions

Thank you!