

Generalized parton distributions of ^3He and the **neutron** orbital structure

Matteo Rinaldi & Sergio Scopetta

Dipartimento di Fisica dell'Università di Perugia
and INFN, Sezione di Perugia, Italy.



December, 7th 2012



Outline


- Hard exclusive processes and **nuclear** Generalized Parton Distributions (GPDs)
- GPDs of ^3He :
 - * GPDs in Impulse Approximation (IA)
(S.S., PRC 70 (2004) 015205; PRC 79 (2009) 025207)
 - * **neutron** dominance for one ^3He observable
(M. Rinaldi, S.S. PRC 85, 062201(R) (2012))
- Extracting the **neutron** information from ^3He data
(M. Rinaldi, S.S. arXiv:1208.2831 [nucl-th])
- Conclusions

GPDs - why?


- Initially, the interest in GPDs was a consequence of the “Spin crisis”(EMC, '88): most of the **proton spin NOT** carried by the **quark helicities** Σ

- Spin Sum Rule:
$$\Sigma + L_q + J_g = \frac{1}{2}$$

- OAM** (L_q) accessed through non forward processes:




DVCS \rightarrow GPDs



SiDIS \rightarrow TMDs

GPDs: Definition (X. Ji PRL 78 (97) 610)

For a $J = \frac{1}{2}$ target,
in a hard-exclusive process,
($Q^2, \nu \rightarrow \infty$)
such as (coherent) DVCS:




the GPDs $H_q(x, \xi, \Delta^2)$ and $E_q(x, \xi, \Delta^2)$ are introduced:

$$\int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle P' | \bar{\psi}_q(-\lambda n/2) \gamma^\mu \psi_q(\lambda n/2) | P \rangle = H_q(x, \xi, \Delta^2) \bar{U}(P') \gamma^\mu U(P) + E_q(x, \xi, \Delta^2) \bar{U}(P') \frac{i\sigma^{\mu\nu} \Delta_\nu}{2M} U(P) + \dots$$

- $\Delta = P' - P$, $q^\mu = (q_0, \vec{q})$, and $\bar{P} = (P + P')^\mu / 2$
- $x = k^+ / P^+$; $\xi = \text{"skewness"} = -\Delta^+ / (2\bar{P}^+)$
- $x \leq -\xi \longrightarrow$ GPDs describe *antiquarks*;
 $-\xi \leq x \leq \xi \longrightarrow$ GPDs describe $q\bar{q}$ pairs; $x \geq \xi \longrightarrow$ GPDs describe *quarks*

GPDs: limits

- when $P' = P$, i.e., $\Delta^2 = \xi = 0$, one recovers the usual PDFs:



$$H_q(x, \xi, \Delta^2) \Rightarrow H_q(x, 0, 0) = q(x); \quad E_q(x, 0, 0) \text{ unknown}$$

- the x -integration yields the q -contribution to the Form Factors (ffs)

$$\int dx \int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle P' | \bar{\psi}_q(-\lambda n/2) \gamma^\mu \psi_q(\lambda n/2) | P \rangle =$$

$$\int dx H_q(x, \xi, \Delta^2) \bar{U}(P') \gamma^\mu U(P) + \int dx E_q(x, \xi, \Delta^2) \bar{U}(P') \frac{\sigma^{\mu\nu} \Delta_\nu}{2M} U(P) + \dots$$

$$\Rightarrow \int dx H_q(x, \xi, \Delta^2) = F_1^q(\Delta^2) \quad \int dx E_q(x, \xi, \Delta^2) = F_2^q(\Delta^2)$$

$$\Rightarrow \text{Defining } \boxed{\tilde{G}_M^q = H_q + E_q} \quad \text{one has } \int dx \tilde{G}_M^q(x, \xi, \Delta^2) = G_M^q(\Delta^2)$$

GPDs: A unique tool...

- to explore the **3-dimensional structure** of hadrons at **parton level** and for many other aspects...
- ...the most important here: access to the parton **orbital angular momentum (OAM)**, solution (?) of the “**Spin Crisis**”: **Ji Sum Rule**:

$$\langle J_q \rangle = \langle \Sigma_q \rangle + \langle L_q \rangle = \int_{-1}^1 dx x \tilde{G}_M^q(x, 0, 0)$$

... but also an experimental challenge:

- Hard exclusive processes \longrightarrow small X-sections;
- Difficult extraction:

$$T_{DVCS} \propto \int_{-1}^1 dx \frac{H_q(x, \xi, \Delta^2)}{x - \xi + i\epsilon} + \dots$$

- Competing **BH** process! Interference (σ -differences) measured.

Why nuclei? - (many reasons...)

I will show results for ^3He , in IA, for the coherent channel... But there is much more:

- for the deuteron:
Berger, Cano, Diehl, Pire, PRL 87 (2001) 142302; Cano & Pire, EPJA 19 (2004) 423; Kirchner & Müller, EPJC 32 (2003) 347.
- beyond the coherent channel:
Liuti & Taneja PRC 72 032201 (R) 2005: spin-0 nuclei, applied to ^4He ;
Guzey, PRC 78 (2008) 025211: constraining the neutron information from incoherent DVCS off nuclei at large t (spin-0 nuclei);
- beyond IA (Shadowing: low x_{Bj} , large distances):
Freund & Strikman, PRC 69 (2004) 015203; Goeke et al., EPJ. A36 (2008) 49-60;
- for finite-heavy nuclei:
Guzey & Strikman, PRC 68 (2003) 015204; Kirchner & Müller, EPJC 32, 347 (2003)
- discussing other issues: Color Transparency phenomena, Liuti & Taneja, PRD 70, 074019 (2004); Energy-momentum tensor in nuclei: Polyakov PLB 555 (2003) 57; Guzey & Siddikov J. Phys. G 32, 251 (2006); Guzey, Thomas, Tsushima PRC 79 (2009) 055205 ...

Why nuclei?

Several relevant issues can be investigated...:

- the nuclear short range structure, at quark level, can be accessed and the reaction mechanism of DIS off nuclei, e.g. the validity of I.A. and the relevance of effects beyond it (non nucleonic degrees of freedom, nucleon modifications...) can be investigated... origin of the EMC effect...
- very important here: the neutron, always from nuclear targets

... with some effort: measurements are difficult:


- In principle, need for a recoil detector to be sure that the nucleus did not break (despite of this, some data are already available!);
- Few data already available:

Airapetian et al. (Hermes) NPB 829, 1 (2010); PRC 81, 035202 (2010)
 $D(\vec{e}, e'\gamma)X$, $\vec{D}(\vec{e}, e'\gamma)X$, $Ne(\vec{e}, e'\gamma)X$; and then $He, N, Ne, Kr, Xe...$
Little A -dependence found

Mazouz et al. (JLab Hall A) PRL 99.242501 (2007); experiment E08-025 (2010)
 $D(\vec{e}, e'\gamma)X = d(\vec{e}, e'\gamma)d + n(\vec{e}, e'\gamma)X + p(\vec{e}, e'\gamma)X$; ³He target?

GPDs for ^3He : why?

- ^3He is **theoretically well known**. Even a **relativistic treatment** may be implemented.
- ^3He has been used extensively as an **effective neutron target**, especially to unveil the **spin content** of the **free neutron**, due to its peculiar spin structure:



In S -wave
 $^3\vec{H}e = \vec{n}!$

^3He always **promising** when the **neutron angular momentum properties** have to be studied. To what extent for **OAM** and \tilde{G}_M^q ? The answer here.

- To this aim, ^3He is a **unique** target:

- * in isoscalar systems, such as ^2H and ^4He , the contribution of the neutron E_q is basically cancelled by that of the proton one ($\kappa_p \simeq -\kappa_n$); **very difficult to extract the neutron E_q , crucial to access OAM, in coherent experiments;**
- * **heavier targets do not allow refined theoretical treatments.**

GPDs of ^3He in IA

coherent DVCS in I.A.


(^3He does not break-up, $\Delta^2 \ll M^2, \xi^2 \ll 1$,):

In a symmetric frame ($\bar{p} = (p + p')/2$) :

$$\begin{aligned} k^+ &= (x + \xi)\bar{P}^+ = (x' + \xi')\bar{p}^+ , \\ (k + \Delta)^+ &= (x - \xi)\bar{P}^+ = (x' - \xi')\bar{p}^+ , \end{aligned}$$

one has, for a given GPD, H_q or \tilde{G}_M^q ,


$$GPD_q(x, \xi, \Delta^2) \simeq \int \frac{dz^-}{4\pi} e^{ix\bar{P}^+ z^-} {}_A \langle P' S' | \hat{O}_q^\mu | PS \rangle_A \big|_{z^+=0, z_\perp=0} .$$



GPDs of ^3He in IA

coherent DVCS in I.A.

(^3He does not break-up, $\Delta^2 \ll M^2, \xi^2 \ll 1$,):



In a symmetric frame ($\bar{p} = (p + p')/2$) :

$$\begin{aligned} k^+ &= (x + \xi)\bar{P}^+ = (x' + \xi')\bar{p}^+ , \\ (k + \Delta)^+ &= (x - \xi)\bar{P}^+ = (x' - \xi')\bar{p}^+ , \end{aligned}$$

one has, for a given GPD, H_q or \tilde{G}_M^q ,

$$GPD_q(x, \xi, \Delta^2) \simeq \int \frac{dz^-}{4\pi} e^{ix\bar{P}^+ z^-} {}_A \langle P' S' | \hat{O}_q^\mu | PS \rangle_A |_{z^+=0, z_\perp=0} .$$


By properly inserting **complete sets of states** for the interacting nucleon and the recoiling system :

GPDs of ^3He in IA

coherent DVCS in I.A.

(^3He does not break-up, $\Delta^2 \ll M^2, \xi^2 \ll 1$,):

In a symmetric frame ($\bar{p} = (p + p')/2$) :



$$\begin{aligned} k^+ &= (x + \xi)\bar{P}^+ = (x' + \xi')\bar{p}^+ , \\ (k + \Delta)^+ &= (x - \xi)\bar{P}^+ = (x' - \xi')\bar{p}^+ , \end{aligned}$$

one has, for a given GPD, H_q or \tilde{G}_M^q ,

$$\begin{aligned} GPD_q(x, \xi, \Delta^2) &= \int \frac{dz^-}{4\pi} e^{ix'\bar{p}^+z^-} \langle P'S' | \sum_{\vec{P}'_R, S'_R, \vec{p}', s'} \{ |P'_R S'_R\rangle | p' s' \} \} \langle P'_R S'_R | \\ &\quad \langle p' s' | \hat{O}_q^\mu \sum_{\vec{P}_R, S_R, \vec{p}, s} \{ |P_R S_R\rangle | p s \} \} \{ \langle P_R S_R | \langle p s | \} | PS \rangle , \end{aligned}$$

and, since $\{ \langle P_R S_R | \langle p s | \} | PS \rangle = \langle P_R S_R, p s | PS \rangle (2\pi)^3 \delta^3(\vec{P} - \vec{P}_R - \vec{p}) \delta_{S, S_R} \delta_{s, s}$,

GPDs of ^3He in IA

H_q^A can be obtained in terms of H_q^N (S.S. PRC 70, 015205 (2004), PRC 79, 025207 (2009)):

$$H_q^A(x, \xi, \Delta^2) = \sum_N \int dE \int d\vec{p} \sum_S \sum_s P_{SS,ss}^N(\vec{p}, \vec{p}', E) \frac{\xi'}{\xi} H_q^N(x', \Delta^2, \xi'),$$

and $\tilde{G}_M^{3,q}$ in terms of $\tilde{G}_M^{N,q}$ (M. Rinaldi, S.S. PRC 85, 062201(R) (2012); arXiv:1208.2831 [nucl-th]):

$$\tilde{G}_M^{3,q}(x, \Delta^2, \xi) = \sum_N \int dE \int d\vec{p} [P_{+-,+ -}^N - P_{+-,- +}^N](\vec{p}, \vec{p}', E) \frac{\xi'}{\xi} \tilde{G}_M^{N,q}(x', \Delta^2, \xi'),$$

where $P_{SS,ss}^N(\vec{p}, \vec{p}', E)$ is the one-body, spin-dependent, off-diagonal spectral function for the nucleon N in the nucleus,


$$P_{SS',ss'}^N(\vec{p}, \vec{p}', E) = \frac{1}{(2\pi)^6} \frac{M\sqrt{ME}}{2} \int d\Omega_t \sum_{s_t} \langle \vec{P}' S' | \vec{p}' s', \vec{t}_{st} \rangle_N \langle \vec{p} s, \vec{t}_{st} | \vec{P} S \rangle_N,$$

evaluated by means of a **realistic** treatment based on **Av18 wave functions** (w.f. from A. Kievsky *et al* NPA 577, 511 (1994), overlaps from A. Kievsky *et. al*, PRC 56, 64 (1997)).

Nucleon GPDs: initially, a simple model by Radyushkin&Musatov (PRD 61, 074027 (2000))

A few words about $P_N^3(\vec{p}, \vec{p} + \vec{\Delta}, E)$:

$$P_N^3(\vec{p}, \vec{p} + \vec{\Delta}, E) = \frac{1}{(2\pi)^3} \frac{1}{2} \sum_M \sum_{f,s} \langle \vec{P}' M | (\vec{P} - \vec{p}) S_f, (\vec{p} + \vec{\Delta}) s \rangle \\ \times \langle (\vec{P} - \vec{p}) S_f, \vec{p} s | \vec{P} M \rangle \delta(E - E_{min} - E_f^*).$$



- the two-body recoiling system can be either the deuteron or a scattering state;
- when a deeply bound nucleon, with high removal energy $E = E_{min} + E_f^*$, leaves the nucleus, the recoiling system is left with high excitation energy E_f^* ;
- the three-body bound state and the two-body bound or scattering state are evaluated within the same (Av18) interaction: the extension of the treatment to heavier nuclei would be extremely difficult

The $\tilde{G}_M^{3,q}$ calculation has the correct limits:

For H_q^3 , correct forward limit and x -integral (S.S. PRC 70, (2004), PRC 79, (2009));



For \tilde{G}_M^3 (M. Rinaldi, S.S. PRC 85, 062201(R) (2012); arXiv:1208.2831 [nucl-th]):

1 - Forward limit: no control on $E_q^3(x, 0, 0)$
no possible check;

2 - Magnetic F.F.:

$$\sum_q \int dx \tilde{G}_M^{3,q}(x, \xi, \Delta^2) = G_M^3(\Delta^2)$$

- in perfect agreement with previous IA, Av18 calculations (L.E. Marcucci et al. PRC 58 (1998))
- in good agreement with data in the region relevant to the coherent process, $-\Delta^2 \ll 0.15 \text{ GeV}^2$
- To have agreement at higher Δ^2 , effects beyond IA are necessary: not important for the coherent channel!




$\tilde{G}_M^{3,q}$: proton and neutron contributions

1 - Forward limit, $\Delta^2 = 0$, $\xi = 0$:

As we hoped, the **neutron** contribution to ${}^3\text{He}$ largely dominates!

($x_3 = (M_A/M)x \simeq 3x$):

The **proton** contribution to ${}^3\text{He}$ is almost negligible!




2 - Non-forward, $\Delta^2 = -0.1 \text{ GeV}^2$, $\xi = 0.1$:

The **neutron** contribution to ${}^3\text{He}$ still dominates

The **proton** contribution to ${}^3\text{He}$ gets sizable

How to get the
neutron information?



$\tilde{G}_M^{3,q}$: Flavor separation

For the u flavor, the neutron contribution (dashed) to ${}^3\text{He}$ (full) is less important than for the d flavor:

Understandable, sketching the formula:


$$\tilde{G}_M^{3,q} \approx P_p^3 \otimes \tilde{G}_M^{p,q} + P_n^3 \otimes \tilde{G}_M^{n,q},$$

where $P_{p(n)}^3$ describes the proton (neutron) dynamics in ${}^3\text{He}$.

As already explained, due to the spin structure of ${}^3\text{He}$, $P_n^3 \gg P_p^3 \longrightarrow$ neutron dominates in the forward limit.

With increasing Δ^2 , for the u flavor, $\tilde{G}_M^{p,u} \gg \tilde{G}_M^{n,u} \longrightarrow$ the proton contribution grows. Not for d !

Besides, 1/2 of the d content of ${}^3\text{He}$ comes from the neutron, only 1/5 of the u one comes from it.




Extracting the neutron - I:

The convolution formula can be written as

$$\tilde{G}_M^{3,q}(x_3, \Delta^2, \xi) = \sum_N \int_{x_3}^{\frac{M_A}{M}} \frac{dz}{z} g_N^3(z, \Delta^2, \xi) \tilde{G}_M^{N,q} \left(\frac{x_3}{z}, \Delta^2, \frac{\xi}{z}, \right),$$

where $g_N^3(z, \Delta^2, \xi)$ is a “light cone off-forward momentum distribution” and, since close to the forward limit it is strongly peaked around $z = 1$

$$g_N^3(z, \Delta^2, \xi) = \int dE \int d\vec{p} \tilde{P}_N^3(\vec{p}, \vec{p} + \vec{\Delta}, E) \delta \left(z + \xi - \frac{M_A}{M} \frac{p^+}{P^+} \right)$$



Extracting the neutron - I:

The convolution formula can be written as

$$\tilde{G}_M^{3,q}(x_3, \Delta^2, \xi) = \sum_N \int_{x_3}^{\frac{M_A}{M}} \frac{dz}{z} g_N^3(z, \Delta^2, \xi) \tilde{G}_M^{N,q} \left(\frac{x_3}{z}, \Delta^2, \frac{\xi}{z} \right),$$

where $g_N^3(z, \Delta^2, \xi)$ is a “light cone off-forward momentum distribution” and, since close to the forward limit it is strongly peaked around $z = 1$

$$\begin{aligned} \tilde{G}_M^{3,q}(x_3, \Delta^2, \xi) &\simeq \text{low } \Delta^2 \simeq \sum_N \tilde{G}_M^{N,q}(x_3, \Delta^2, \xi) \int_0^{\frac{M_A}{M}} dz g_N^3(z, \Delta^2, \xi) \\ &= G_M^{3,p,point}(\Delta^2) \tilde{G}_M^p(x_3, \Delta^2, \xi) + G_M^{3,n,point}(\Delta^2) \tilde{G}_M^n(x_3, \Delta^2, \xi). \end{aligned}$$

where, at $x_3 < 0.7$, the **magnetic point like ff** has been introduced


$$G_M^{3,N,point}(\Delta^2) = \int dE \int d\vec{p} \tilde{P}_N^3(\vec{p}, \vec{p} + \vec{\Delta}, E) = \int_0^{\frac{M_A}{M}} dz g_N^3(z, \Delta^2, \xi).$$

Extracting the neutron - II:

Validity of the approximated formula:

full: IA calculation, $\tilde{G}_M^3(x, \Delta^2, \xi)$ and
 proton and neutron contributions to it,
 at $\Delta^2 = -0.1 \text{ GeV}^2$, $\xi = 0.1$;

dashed: same quantities, with the
 approximated formula:



$$\begin{aligned} \tilde{G}_M^{3,q}(x, \Delta^2, \xi) &\simeq G_M^{3,p,point}(\Delta^2) \tilde{G}_M^p(x, \Delta^2, \xi) \\ &+ G_M^{3,n,point}(\Delta^2) \tilde{G}_M^n(x, \Delta^2, \xi) \end{aligned}$$

Impressive agreement! The **only Nuclear Physics ingredient** in the approximated formula is the **magnetic point like ff**, which is under good theoretical control:

Δ^2 [GeV ²]	$G_M^{3,p,point}$ Av18	$G_M^{3,p,point}$ Av14	$G_M^{3,n,point}$ Av18	$G_M^{3,n,point}$ Av14
0	-0.044	-0.049	0.879	0.874
-0.1	0.040	0.038	0.305	0.297
-0.2	0.036	0.035	0.125	0.119

Extracting the neutron - III:

The approximated relation can now be solved to extract the neutron contribution:


$$\tilde{G}_M^{n,extr}(x, \Delta^2, \xi) \simeq \frac{1}{G_M^{3,n,point}(\Delta^2)} \left\{ \tilde{G}_M^3(x, \Delta^2, \xi) - G_M^{3,p,point}(\Delta^2) \tilde{G}_M^p(x, \Delta^2, \xi) \right\},$$

from data for $\tilde{G}_M^3(x, \Delta^2, \xi)$ and $\tilde{G}_M^p(x, \Delta^2, \xi)$, using as theoretical ingredients the **magnetic point like ffs** only.

The procedure works nicely!

full : the neutron model for $\tilde{G}_M^n(x, \Delta^2, \xi)$ and the different flavor contributions to it used in the IA calculation,
at $\Delta^2 = -0.1 \text{ GeV}^2$, $\xi = 0.1$;

dashed: the neutron extracted using the IA calculation for $\tilde{G}_M^3(x, \Delta^2, \xi)$ and the model used in it for $\tilde{G}_M^p(x, \Delta^2, \xi)$ together with the **magnetic point like ffs**.



Extracting the neutron - IV:


The validity of the extraction procedure is emphasized showing the following ratio, which would be one if the procedure were perfect:

$$r_n(x, \Delta^2, \xi) = \frac{\tilde{G}_M^{n,extr}(x, \Delta^2, \xi)}{\tilde{G}_M^n(x, \Delta^2, \xi)}$$

full: forward limit

dashed: $\Delta^2 = -0.1 \text{ GeV}^2, \xi = 0$;

dot-dashed: $\Delta^2 = -0.1 \text{ GeV}^2, \xi = 0.1$;



at $x < 0.7$, in all the kinematical range relevant for coherent DVCS at JLab, the error in the extraction is a few percents.

Extracting the neutron - V:


The validity of the extraction procedure is emphasized showing the same ratio, evaluated using different models for the GPDs as input in the IA calculation:

$$r_n(x, \Delta^2, \xi) = \frac{\tilde{G}_M^{n,extr}(x, \Delta^2, \xi)}{\tilde{G}_M^n(x, \Delta^2, \xi)}$$

dashed: the model of Radyushkin, at $\Delta^2 = -0.1 \text{ GeV}^2, \xi = 0$;

full: a very different model based on a constituent quark scenario (S.S, V. Vento EPJA 16, 527 (2003)) at $\Delta^2 = -0.1 \text{ GeV}^2, \xi = 0$;

crosses: another very different model, the MIT bag model (X.-D. Ji et al., PRD 56 (1997) 5511) at $\Delta^2 = -0.1 \text{ GeV}^2, \xi = 0$.



at $x < 0.7$, in all the kinematical range relevant for coherent DVCS at JLab, the error in the extraction due to the use of a different nucleonic model is a few percents at most.

Conclusions

What has been done:

- * An instant form, I.A. calculation of $\tilde{G}_M^3(x, \xi, \Delta^2)$, within AV18 wave functions;
- * the neutron contribution dominates at low Δ^2 ;
- * an extraction procedure of the neutron information, able to take into account all the nuclear effects encoded in an IA analysis, weakly dependent on both the nuclear potential and the nucleonic model used in the calculation, has been proposed;

What is being done: to estimate X-sections (DVCS, BH, Int....)



→ a proposal of coherent DVCS off ^3He at JLab@12 GeV?

What has to be done, in case experiments are performed at higher Δ^2 :



- * To implement a RELATIVISTIC TREATMENT
(S.S., Del Dotto, Pace, Salmè et al., Nuovo Cim. C035N2 (2012) 101-106)
- * and/or to go beyond IA, including many body currents into the scheme.
- * Incoherent? Nice, but FSI have to be evaluated

What is expected: To get the OAM parton structure of the neutron!
(and “imaging” (IPD-PDs), “spatial distribution of nuclear forces”...)

coherent vs. incoherent DVCS:



Coherent



Incoherent


The calculation has the correct limits:

1 - Forward limit: the ratio:

$$R_q(x, 0, 0) = \frac{H_q^3(x, 0, 0)}{2H_q^p(x, 0, 0) + H_q^n(x, 0, 0)}$$

$$= \frac{q^3(x)}{2q^p(x) + q^n(x)}$$


shows an EMC-like behavior;



2 - Charge F.F.:

$$\int dx H_q^3(x, \xi, \Delta^2) = F_q^3(\Delta^2)$$



in good agreement with data in the region relevant to the coherent process, $\Delta^2 \ll 0.25 \text{ GeV}^2$.



Nuclear effects - general features



Nuclear effects grow with ξ at fixed Δ^2 , and with Δ^2 at fixed ξ :



$$R_q^{(0)}(x, \xi, \Delta^2) = \frac{H_q^3(x, \xi, \Delta^2)}{2H_q^{3,p}(x, \xi, \Delta^2) + H_q^{3,n}(x, \xi, \Delta^2)}$$



$$H_q^{3,N}(x, \xi, \Delta^2) = \tilde{H}_q^N(x, \xi) F_q^3(\Delta^2)$$

$R_q^{(0)}(x, \xi, \Delta^2)$ would be one if there were no nuclear effects;
as it is found also for the deuteron, there is **no factorization** into terms
dependent separately on Δ^2 and x, ξ (the factorization hypotheses has been
used to estimate nuclear GPDs).

Nuclear effects - flavor dependence



Nuclear effects are bigger for the d flavor rather than for the u flavor:



$$R_q^{(0)}(x, \xi, \Delta^2) = \frac{H_q^3(x, \xi, \Delta^2)}{2H_q^{3,p}(x, \xi, \Delta^2) + H_q^{3,n}(x, \xi, \Delta^2)}$$

$$H_q^{3,N}(x, \xi, \Delta^2) = \tilde{H}_q^N(x, \xi) F_q^3(\Delta^2)$$

$R_q^{(0)}(x, \xi, \Delta^2)$ would be one if there were no nuclear effects;







This is a typical **conventional, IA** effect (spectral functions are different for p and n in ${}^3\text{He}$, not isoscalar!); if (not) found, clear indication on the reaction mechanism of **DIS off nuclei**.

Nuclear effects - flavor dependence




The **d** and **u** distributions follow the pattern of the **neutron** and **proton** light-cone momentum distributions, respectively:



Nuclear effects @ $x = \xi$



Nuclear effects are large even in the important region $x = \xi$:



Nuclear effects - the binding

General IA formula: $H_q^A(x, \xi, \Delta^2) \simeq \sum_N \int_x^1 \frac{dz}{z} h_N^A(z, \xi, \Delta^2) H_q^N\left(\frac{x}{z}, \frac{\xi}{z}, \Delta^2\right)$

where



$$h_N^A(z, \xi, \Delta^2) = \int dE d\vec{p} P_N^A(\vec{p}, \vec{p} + \vec{\Delta}, E) \delta\left(z + \xi - \frac{p^+}{P^+}\right)$$

$$P_N^3(\vec{p}, \vec{p} + \vec{\Delta}, E) = \bar{\sum}_M \sum_{s,f} \langle \vec{P}' M | \vec{P}_f, (\vec{p} + \vec{\Delta}) s \rangle \\ \times \langle \vec{P}_f, \vec{p} s | \vec{P} M \rangle \delta(E - E_{min} - E_f^*)$$

using the Closure Approximation, $E_f^* = \bar{E}$:

$$P_N^3(\vec{p}, \vec{p} + \vec{\Delta}, E) \simeq \bar{\sum}_M \sum_s \langle \vec{P}' M | a_{\vec{p}+\vec{\Delta},s} a_{\vec{p},s}^\dagger | \vec{P} M \rangle \\ \delta(E - E_{min} - \bar{E}) = \\ = n(\vec{p}, \vec{p} + \vec{\Delta}) \delta(E - E_{min} - \bar{E}),$$



Spectral function substituted by a Momentum distribution



Nuclear effects - the binding

Nuclear effects are bigger than in the forward case:
dependence on the binding

- In calculations using $n(\vec{p}, \vec{p} + \vec{\Delta})$ instead of $P_N^3(\vec{p}, \vec{p} + \vec{\Delta}, E)$, in addition to the **IA**, also the **Closure** approximation has been assumed;
- 5 % to 10 % **binding** effect between $x = 0.4$ and 0.7 - much **bigger** than in the forward case;
- for $A > 3$, the evaluation of $P_N^3(\vec{p}, \vec{p} + \vec{\Delta}, E)$ is **difficult** - such an effect is **not under control**: **Conventional** nuclear effects can be **mistaken for exotic** ones;
- for ${}^3\text{He}$ it is possible : this makes it a **unique** target, even among the **Few-Body** systems.



Dependence on the NN interaction

Nuclear effects are bigger than in the forward case: dependence on the potential



Forward case: Calculations using the **AV14** or **AV18** interactions are **indistinguishable**



Non-forward case: Calculations using the **AV14** and **AV18** interactions **do differ**:

