




<http://cern.ch/Gfitter>

EPJC 72, 2205 (2012), arXiv:1209.2716

After the Higgs: Status and Prospects of The ElectroWeak fit of the SM and Beyond






This presentation:

- Introduction to the Electroweak Fit
 - Inputs to the electroweak fit
- ✓ After the Higgs: predictions for key observables
- ✓ BSM: Modified Higgs couplings
- ✓ Prospects for LHC-300 and ILC/GigaZ
- Conclusion & Outlook




A **G**eneric **F**itter Project
for HEP Model Testing

- Gfitter = state-of-the-art HEP model testing tool for LHC era
- Latest results always available at: <http://cern.ch/Gfitter>
 - (Most) results of this presentation: EPJC 72, 2205 (2012)
 - LHC-300 and ILC/GigaZ prospects paper to appear on arXiv this week !
- Gfitter software and features:
 - Modular, object-oriented C++, relying on ROOT, XML, python, etc.
 - Core package with data-handling, fitting, and statistics tools
 - Independent “plug-in” physics libraries: SM, 2HDM, multiple BSM models, ...




- ✓ Observables receive quantum loop corrections from ‘unseen’ virtual effects.
- ✓ If system is over-constrained, one can fit for unknown parameters or test the model’s self-consistency.
- ✓ If precision is better than typical loop factor ($\alpha \approx 1/137$), test the model or try to obtain info on new physics in loops.
 - For example, in the past EW fits were used to predict the Higgs mass.

- Huge amount of pioneering work by many!
- Needed to understand importance of loop corrections
 - Important observables (now) known at least at two-loop order, sometimes more.
- High-precision Standard Model (SM) predictions and measurements required
 - First from LEP/SLC, then Tevatron, now LHC.



- Top mass predictions from loop effects available since ~1990.
 - Official LEPEW fit since 1993.
- The EW fits have always been able to predict the top mass correctly!

- EW fits performed by many groups in past and present.
 - D. Bardinet al. (ZFITTER), G. Passarino et al. (TOPAZ0), LEPEW WG (M. Grünewald, K. Mönig et al.), J. Erler (GAP), Bayesian fit (M. Ciuchini, L. Silvestrini et al.), etc ...
 - Important results obtained!
- Several groups pursuing global beyond-SM fits, especially SUSY.
- Global SM fits also used at lower energies [CKM-matrix].
- Fits of the different groups agree very well.
- Some differences in treatment of theory errors, which just start to matter.
 - E.g. Gfitter uses “R-fit prescription”: theoretical uncertainties included in χ^2 with flat likelihood in allowed ranges
 - I.e. theoretical and experimental errors added linearly (= conservative).




The predictive power of the SM

- As the Z boson couples to all fermions, it is ideal to measure & study both the electroweak and strong interactions.

- Tree level relations for $Z \rightarrow f\bar{f}$

- $$i\bar{f}\gamma^\mu (g_{V,f} - g_{A,f}\gamma_5) f Z_\mu$$





- Prediction EWSB at tree-level:

$$\frac{M_W^2}{M_Z^2 c_W^2} = 1$$

- The impact of loop corrections

- Absorbed into EW form factors: ρ , κ , Δr
- Effective couplings at the Z-pole
- Quadratically dependent on m_t ,
logarithmic dependence on M_H




$$g_{V,f} = \sqrt{\rho_Z^f} (I_3^f - 2Q^f \sin^2 \theta_{\text{eff}}^f)$$

$$g_{A,f} = \sqrt{\rho_Z^f} I_3^f$$

$$\sin^2 \theta_{\text{eff}}^f = \kappa_Z^f \sin^2 \theta_W$$


$$M_W^2 = \frac{M_Z^2}{2} \left(1 + \sqrt{1 - \frac{\sqrt{8}\pi\alpha(1 + \Delta r)}{G_F M_Z^2}} \right)$$

Gfitter group, EPJC 72, 2003 (2012)



- M_H was last missing input parameter of the electroweak fit
- Indirect determination from EW fit (2012): $M_H = 96^{+31}_{-24}$ GeV
 - With direct limits incorporated in the EW fit: $M_H = 120^{+12}_{-5}$ GeV

- Discovery of Higgs-like boson at LHC
 - Cross section, production rate time branching ratios, spin, parity sofar compatible with SM Higgs boson.
- This talk: assume boson is SM Higgs.
- Use in EW fit: $M_H = 125.7 \pm 0.4 \text{ GeV}$
 - ATLAS: $M_H = 126.0 \pm 0.4 \pm 0.4 \text{ GeV}$
 - CMS: $M_H = 125.3 \pm 0.4 \pm 0.5 \text{ GeV}$
[arXiv:1207.7214, arXiv:1207.7235]
- Change in average between fully uncorrelated and fully correlated systematic uncertainties is minor:
 $\delta M_H : 0.4 \rightarrow 0.5 \text{ GeV}$



Unique situation:

- *For first time SM is fully over-constrained.*
- *And for first time electroweak observables can be unambiguously predicted at loop level.*
- *Powerful predictions of key observables now possible, much better than w/o M_H*

Can now test for:

- Self-consistency of SM
 - Possible contributions from BSM models
-
- The focus of this talk ...

■ Total cross-section of $Z \rightarrow f\bar{f}$

- Expressed in terms of partial decay width of initial and final width:


$$\sigma_{f\bar{f}}^Z = \sigma_{f\bar{f}}^0 \frac{s\Gamma_Z^2}{(s - M_Z^2)^2 + s^2\Gamma_Z^2/M_Z^2} \frac{1}{R_{\text{QED}}} \quad \text{with} \quad \sigma_{f\bar{f}}^0 = \frac{12\pi}{M_Z^2} \frac{\Gamma_{ee}\Gamma_{f\bar{f}}}{\Gamma_Z^2}$$

Corrected for QED radiation

- Full width: $\Gamma_Z = \Gamma_{ee} + \Gamma_{\mu\mu} + \Gamma_{\tau\tau} + \Gamma_{\text{had}} + \Gamma_{\text{inv}}$
- (Correlated set of measurements.)

■ Set of input (width) parameters to EW fit:

- Z mass and width: M_Z , Γ_Z
- Hadronic pole cross section:
 $\sigma_{\text{had}}^0 = 12\pi/M_Z^2 \cdot \Gamma_{ee}\Gamma_{\text{had}}/\Gamma_Z^2$
- Three leptonic ratios (lepton univ.):
 $R_\ell^0 = R_e^0 = \Gamma_{\text{had}}/\Gamma_{ee} (= R_\mu^0 = R_\tau^0)$
- Hadronic-width ratios: R_b^0 , R_c^0



■ Definition of Asymmetry

- Distinguish vector and axial-vector couplings of the Z

$$A_f = \frac{g_{L,f}^2 - g_{R,f}^2}{g_{L,f}^2 + g_{R,f}^2} = \frac{2g_{V,f} g_{A,f}}{g_{V,f}^2 + g_{A,f}^2}$$

- Directly related to: $\sin^2 \theta_{\text{eff}}^{f\bar{f}} = \frac{1}{4Q_f} \left(1 + \mathcal{R}e \left(\frac{g_{V,f}}{g_{A,f}} \right) \right)$

■ Observables

- In case of no beam polarisation (LEP) use final state angular distribution to define *forward/backward asymmetry*:

$$A_{FB}^f = \frac{N_F^f - N_B^f}{N_F^f + N_B^f}$$

$$A_{FB}^{0,f} = \frac{3}{4} A_e A_f$$

- Polarised beams (SLC), define *left/right asymmetry*:


$$A_{LR}^f = \frac{N_L^f - N_R^f}{N_L^f + N_R^f} \frac{1}{\langle |P|_e \rangle} \quad A_{LR}^0 = A_e$$

- Measurements: $A_{FB}^{0,\ell}$, $A_{FB}^{0,c}$, $A_{FB}^{0,b}$, A_ℓ , A_c , A_b


Latest averages for M_W and m_{top}



Latest Tevatron result from: arXiv:1204.0042




Mass of the W Boson



March 2012

Tevatron result from: arXiv:1305.3929

Mass of the Top Quark



(LHC average: 173.29 ± 0.95 GeV/c²)

- The EW fit requires precise knowledge of $\alpha(M_Z)$ – better than 1% level
 - Enters various places: hadr. radiator functions, predictions of M_W and $\sin^2\theta_{\text{eff}}^f$
- Conventionally parametrized as ($\alpha(0)$ = fine structure constant) :

$$\alpha(s) = \frac{\alpha(0)}{1 - \Delta\alpha(s)}$$

- Evolution with renormalization scale:

$$\Delta\alpha(s) = \Delta\alpha_{\text{lep}}(s) + \Delta\alpha_{\text{had}}^{(5)}(s) + \Delta\alpha_{\text{top}}(s)$$

- The EW fit requires precise knowledge of $\alpha(M_Z)$ – better than 1% level
 - Enters various places: hadr. radiator functions, predictions of M_W and $\sin^2\theta_{\text{eff}}^f$
- Conventionally parametrized as ($\alpha(0)$ = fine structure constant) :

$$\alpha(s) = \frac{\alpha(0)}{1 - \Delta\alpha(s)}$$

- Evolution with renormalization scale:

$$\Delta\alpha(s) = \Delta\alpha_{\text{lep}}(s) + \Delta\alpha_{\text{had}}^{(5)}(s) + \Delta\alpha_{\text{top}}(s)$$

- Leptonic term known up to *four* loops (for $q^2 \gg m_l^2$)
- Top quark contribution known up to 2 loops, *small*: -0.7×10^{-4}

[C. Sturm, arXiv:
1305.0581]

[M. Steinhauser,
PLB 429, 158 (1998)]

- The EW fit requires precise knowledge of $\alpha(M_Z)$ – better than 1% level
 - Enters various places: hadr. radiator functions, predictions of M_W and $\sin^2\theta_{\text{eff}}^f$
- Conventionally parametrized as ($\alpha(0)$ = fine structure constant) :

$$\alpha(s) = \frac{\alpha(0)}{1 - \Delta\alpha(s)}$$

- Evolution with renormalization scale:

$$\Delta\alpha(s) = \Delta\alpha_{\text{lep}}(s) + \Delta\alpha_{\text{had}}^{(5)}(s) + \Delta\alpha_{\text{top}}(s)$$

- Hadronic contribution (from the 5 light quarks) completely dominates overall uncertainty on $\alpha(M_Z)$.
- Difficult to calculate, cannot be obtained from pQCD alone.
 - Analysis of low-energy e^+e^- data
 - Usage of pQCD if lack of data
- Similar analysis to evaluation of hadronic contribution to $(g-2)_\mu$

$$\Delta\alpha_{\text{had}}^{(5)}(M_Z) = (274.9 \pm 1.0) \cdot 10^{-4}$$

[M. Davier et al., Eur. Phys. J. C71, 1515 (2011)]

- Radiative corrections are important!

- E.g. consider tree-level EW unification relation:

$$M_W^2 \Big|_{\text{tree-level}} = \frac{M_Z^2}{2} \cdot \left(1 + \sqrt{1 - \frac{\sqrt{8}\pi\alpha}{G_F M_Z^2}} \right)$$

- This predicts: $M_W = (79.964 \pm 0.005) \text{ GeV}$
- Experiment: $M_W = (80.385 \pm 0.015) \text{ GeV}$

- Without loop corrections: shift of 400 MeV, 27σ discrepancy!

- Radiative corrections are important!
 - E.g. consider tree-level EW unification relation:

$$M_W^2 \Big|_{\text{tree-level}} = \frac{M_Z^2}{2} \cdot \left(1 + \sqrt{1 - \frac{\sqrt{8}\pi\alpha}{G_F M_Z^2}} \right)$$
 - This predicts: $M_W = (79.964 \pm 0.005) \text{ GeV}$
 - Experiment: $M_W = (80.385 \pm 0.015) \text{ GeV}$
- Without loop corrections: shift of 400 MeV, 27σ discrepancy!
- In EW fit with Gfitter we use state-of-the-art calculations:
 - M_W Mass of the W boson [M. Awramik et al., Phys. Rev. D69, 053006 (2004)]
 - $\sin^2\theta_{\text{eff}}^f$ Effective weak mixing angle [M. Awramik et al., JHEP 11, 048 (2006),
M. Awramik et al., Nucl.Phys.B813:174-187 (2009)]
 - Full two-loop + leading beyond-two-loop form factor corrections
 - Γ_{had} QCD Adler functions at N³LO [P. A. Baikov et al., PRL108, 222003 (2012)]
 - N³LO prediction of the hadronic cross section
 - R_b Partial width of Z→bb [Freitas et al., JHEP08, 050 (2012)] ← **Update!**
EW 2-loop calc.

- Radiative corrections are important!
 - E.g. consider tree-level EW unification relation:

$$M_W^2 \Big|_{\text{tree-level}} = \frac{M_Z^2}{2} \cdot \left(1 + \sqrt{1 - \frac{\sqrt{8}\pi\alpha}{G_F M_Z^2}} \right)$$
 - This predicts: $M_W = (79.964 \pm 0.005) \text{ GeV}$
 - Experiment: $M_W = (80.385 \pm 0.015) \text{ GeV}$
- Without loop corrections: shift of 400 MeV, 27σ discrepancy!
- In EW fit with Gfitter we use state-of-the-art calculations:
 - M_W Mass of the W boson [M. Awramik et al., Phys. Rev. D69, 053006 (2004)]
 - $\sin^2\theta_{\text{eff}}^f$ Effective weak mixing angle [M. Awramik et al., JHEP 11, 048 (2006),
M. Awramik et al., Nucl.Phys.B813:174-187 (2009)]
 - Full two-loop + leading beyond-two-loop form factor corrections
 - Γ_{had} QCD Adler functions at N³LO [P. A. Baikov et al., PRL108, 222003 (2012)]
 - N³LO prediction of the hadronic cross section
 - R_b Partial width of Z→bb [Freitas et al., JHEP08, 050 (2012)] ← Update!
EW 2-loop calc.
- Two nuisance parameters in EW fit for theoretical uncertainties:
 - $\delta M_W (4 \text{ MeV}), \delta \sin^2\theta_{\text{eff}}^f (4.7 \times 10^{-5})$

- The branching ratio R_b^0 = partial decay width of $Z \rightarrow b\bar{b}$ to $Z \rightarrow q\bar{q}$
- We use calculation with full EW 2-loop corrections of $Z \rightarrow b\bar{b}$
 - From A. Freitas et al, JHEP 1208 (2012) 050, Erratum. 1305 (2013) 074.

Recently a mistake was found in this calculation.

- **Old:** Two-loop corrections to R_b^0 comparable to experimental uncertainty (6.6×10^{-4})
 - Moved theoretical prediction by 1.5σ
 - Much more than the originally estimated theory uncertainty!
 - **New:** bug in calculation of R_b^0 has been corrected, resulting in a sizable reduction of the size of the two-loop correction.
- All results shown here and on Gfitter homepage use the corrected R_b^0 calculation.

■ Latest experimental inputs:

- **Z-pole observables:** from LEP / SLC
[ADLO+SLD, Phys. Rept. 427, 257 (2006)]
- **M_W and Γ_W** from LEP/Tevatron
[arXiv:1204.0042, arXiv:1302.3415]
- **m_{top}** latest avg from Tevatron
[arXiv:1305.3929]
- **m_c , m_b** world averages (PDG)
[PDG, J. Phys. G33,1 (2006)]
- **$\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)$** including α_S dependency
[Davier et al., EPJC 71, 1515 (2011)]
- **M_H** from LHC
[arXiv:1207.7214, arXiv:1207.7235]

■ 7 (+2) free fit parameters:

- M_H , M_Z , $\alpha_S(M_Z^2)$, $\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)$, m_t , m_c , m_b
- 2 theory nuisance parameters
 - δM_W (4 MeV), $\delta \sin^2 \theta_{\text{eff}}^l$ (4.7×10^{-5})

M_H [GeV] ^o	125.7 ± 0.4	LHC
M_W [GeV]	80.385 ± 0.015	Tevatron
Γ_W [GeV]	2.085 ± 0.042	
M_Z [GeV]	91.1875 ± 0.0021	LEP
Γ_Z [GeV]	2.4952 ± 0.0023	
σ_{had}^0 [nb]	41.540 ± 0.037	
R_ℓ^0	20.767 ± 0.025	
$A_{\text{FB}}^{0,\ell}$	0.0171 ± 0.0010	SLC
$A_\ell^{(*)}$	0.1499 ± 0.0018	
$\sin^2 \theta_{\text{eff}}^\ell(Q_{\text{FB}})$	0.2324 ± 0.0012	SLC
A_c	0.670 ± 0.027	
A_b	0.923 ± 0.020	LEP
$A_{\text{FB}}^{0,c}$	0.0707 ± 0.0035	
$A_{\text{FB}}^{0,b}$	0.0992 ± 0.0016	Tevatron
R_c^0	0.1721 ± 0.0030	
R_b^0	0.21629 ± 0.00066	
$\overline{m}_c$ [GeV]	$1.27^{+0.07}_{-0.11}$	Tevatron
$\overline{m}_b$ [GeV]	$4.20^{+0.17}_{-0.07}$	
m_t [GeV]	173.20 ± 0.87	
$\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)^{(\dagger\Delta)}$	2756 ± 10	

Electroweak Fit – SM Fit Results



■ From the Gfitter Group, EPJC 72, 2205 (2012)


■ Left: full fit incl. M_H

■ Middle: not incl. M_H


■ Right: fit incl M_H , not the row


Parameter	Input value	Free in fit	Fit Result	Fit without M_H measurements	Fit without exp. input in line
M_H [GeV] ^o	$125.7^{+0.4}_{-0.4}$	yes	$125.7^{+0.4}_{-0.4}$	94.7^{+25}_{-22}	94.7^{+25}_{-22}
M_W [GeV]	80.385 ± 0.015	–	$80.367^{+0.006}_{-0.007}$	$80.367^{+0.006}_{-0.007}$	80.360 ± 0.011
Γ_W [GeV]	2.085 ± 0.042	–	2.091 ± 0.001	2.091 ± 0.001	2.091 ± 0.001
M_Z [GeV]	91.1875 ± 0.0021	yes	91.1878 ± 0.0021	91.1878 ± 0.0021	91.1978 ± 0.0114
Γ_Z [GeV]	2.4952 ± 0.0023	–	2.4954 ± 0.0014	2.4954 ± 0.0014	2.4950 ± 0.0017
σ_{had}^0 [nb]	41.540 ± 0.037	–	41.479 ± 0.014	41.479 ± 0.014	41.471 ± 0.015
R_ℓ^0	20.767 ± 0.025	–	20.740 ± 0.017	20.740 ± 0.017	20.715 ± 0.026
$A_{\text{FB}}^{0,\ell}$	0.0171 ± 0.0010	–	$0.01626^{+0.0001}_{-0.0002}$	$0.01626^{+0.0001}_{-0.0002}$	0.01624 ± 0.0002
$A_\ell^{(*)}$	0.1499 ± 0.0018	–	0.1472 ± 0.0007	0.1472 ± 0.0007	–
$\sin^2\theta_{\text{eff}}^\ell(Q_{\text{FB}})$	0.2324 ± 0.0012	–	$0.23149^{+0.00010}_{-0.00008}$	$0.23149^{+0.00010}_{-0.00008}$	0.23150 ± 0.00009
A_c	0.670 ± 0.027	–	$0.6679^{+0.00034}_{-0.00028}$	$0.6679^{+0.00034}_{-0.00028}$	0.6680 ± 0.00031
A_b	0.923 ± 0.020	–	$0.93464^{+0.00005}_{-0.00007}$	$0.93464^{+0.00005}_{-0.00007}$	0.93463 ± 0.00006
$A_{\text{FB}}^{0,c}$	0.0707 ± 0.0035	–	0.0738 ± 0.0004	0.0738 ± 0.0004	0.0737 ± 0.0004
$A_{\text{FB}}^{0,b}$	0.0992 ± 0.0016	–	0.1032 ± 0.0005	0.1032 ± 0.0005	0.1034 ± 0.0003
R_c^0	0.1721 ± 0.0030	–	0.17223 ± 0.00006	0.17223 ± 0.00006	0.17223 ± 0.00006
R_b^0	0.21629 ± 0.00066	–	0.21548 ± 0.00005	0.21548 ± 0.00005	0.21547 ± 0.00005
$\overline{m}_c$ [GeV]	$1.27^{+0.07}_{-0.11}$	yes	$1.27^{+0.07}_{-0.11}$	$1.27^{+0.07}_{-0.11}$	–
$\overline{m}_b$ [GeV]	$4.20^{+0.17}_{-0.07}$	yes	$4.20^{+0.17}_{-0.07}$	$4.20^{+0.17}_{-0.07}$	–
m_t [GeV]	173.20 ± 0.87	yes	173.53 ± 0.82	173.53 ± 0.82	$176.11^{+2.88}_{-2.35}$
$\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)^{(\dagger\Delta)}$	2757 ± 10	yes	2755 ± 11	2755 ± 11	2718^{+49}_{-43}
$\alpha_s(M_Z^2)$	–	yes	$0.1190^{+0.0028}_{-0.0027}$	$0.1190^{+0.0028}_{-0.0027}$	0.1190 ± 0.0027
$\delta_{\text{th}} M_W$ [MeV]	$[-4, 4]_{\text{theo}}$	yes	4	4	–
$\delta_{\text{th}} \sin^2\theta_{\text{eff}}^\ell^{(\dagger)}$	$[-4.7, 4.7]_{\text{theo}}$	yes	–0.6	–0.5	–

- Results drawn as *pull values*:
→ deviations to the *indirect* determinations, divided by *total error*.
- Total error:
error of direct measurement plus error from indirect determination.
- Black: direct measurement (data)
- Orange: full fit
- Light-blue: fit excluding input from the row
- The prediction (light blue) is often more precise than the measurement!




- Results drawn as *pull values*:
→ deviations to the *indirect* determinations, divided by *total error*.
- Total error:
error of direct measurement plus error from indirect determination.
- Black: direct measurement (data)
- Orange: full fit
- Light-blue: fit excluding input from the row
- The prediction (light blue) is often more precise than the measurement!






Plot inspired by Eberhardt et al. [arXiv:1209.1101]

- No individual value exceeds 3σ
- Small pulls for M_H , M_Z , $\Delta \alpha_{\text{had}}^{(5)}(M_Z^2)$, $\overline{m}_c$, $\overline{m}_b$ indicate that input accuracies exceed fit requirements
- Largest deviations in b-sector: $A_{\text{FB}}^{0,b}$ with 2.5σ
 - $\rightarrow$ largest contribution to χ^2
- R_b^0 using one-loop calculation -0.8σ
 - R_b^0 has only little dependence on M_H
- Most affected when including M_H : M_W prediction:
 - Shift in predicted M_W value of ~ 13 MeV.




- Toy analysis: p-value for wrongly rejecting the SM = 18^{+2}_{-2} (theo) %
 - p-value is equivalent to 0.9σ
 - Evaluated with 20k pseudo experiments – follows χ^2 with 14 d.o.f.
 - For comparison: $\chi^2_{\text{min}} = 18.1 \rightarrow \text{Prob}(\chi^2_{\text{min}}, 14) = 20\%$
- Large value of χ^2_{min} *not* due to inclusion of M_H measurement.
 - Without M_H measurement: $\chi^2_{\text{min}} = 16.7 \rightarrow \text{Prob}(\chi^2_{\text{min}}, 13) = 21\%$

- Scan of $\Delta\chi^2$ profile versus M_H
 - Grey band: fit w/o M_H measurement
 - Blue line: full SM fit, with M_H meas.
 - Fit w/o M_H measurement gives:
 $M_H = 94^{+25}_{-22}$ GeV
 - Consistent at 1.3σ with LHC measurement.




- Bottom plot: impact of other most sensitive Higgs observables

- Determination of M_H removing all sensitive observables except the given one.
- Known tension (2.5σ) between $A_l(\text{SLD})$, $A_{\text{FB}}^{0,b}$, and M_W clearly visible.



- Scan of $\Delta\chi^2$ profile versus M_W
 - Also shown: SM fit with minimal inputs:
 M_Z , G_F , $\Delta\alpha_{\text{had}}^{(5)}(M_Z)$, $\alpha_s(M_Z)$, M_H , and fermion masses
 - Good consistency between total fit and SM w/ minimal inputs
- M_H measurement allows for precise constraint on M_W
 - Agreement at 1.4σ
- Fit result for indirect determination of M_W (full fit w/o M_W):



$$\begin{aligned}
 M_W &= 80.3593 \pm 0.0056_{m_t} \pm 0.0026_{M_Z} \pm 0.0018_{\Delta\alpha_{\text{had}}} \\
 &\quad \pm 0.0017_{\alpha_s} \pm 0.0002_{M_H} \pm 0.0040_{\text{theo}} \\
 &= 80.359 \pm 0.011_{\text{tot}} ,
 \end{aligned}$$

- More precise estimate of M_W than the direct measurements!
 - Uncertainty on world average measurement: 15 MeV

- Right: scan of $\Delta\chi^2$ profile versus $\sin^2\theta_{\text{eff}}^l$
 - All sensitive measurements removed from the SM fit.
 - Also shown: SM fit with minimal inputs


- M_H measurement allows for very precise constraint on $\sin^2\theta_{\text{eff}}^l$

- Fit result for indirect determination of $\sin^2\theta_{\text{eff}}^l$:


$$\begin{aligned}\sin^2\theta_{\text{eff}}^l &= 0.231496 \pm 0.000030_{m_t} \pm 0.000015_{M_Z} \pm 0.000035_{\Delta\alpha_{\text{had}}} \\ &\quad \pm 0.000010_{\alpha_S} \pm 0.000002_{M_H} \pm 0.000047_{\text{theo}}, \\ &= 0.23150 \pm 0.00010_{\text{tot}},\end{aligned}$$

- More precise than direct determination (from LEP/SLD) !

- Uncertainty on LEP/SLD average: 1.6×10^{-4}



Indirect determination of top mass




- Shown: scan of $\Delta\chi^2$ profile versus m_t (without m_t measurement)
 - M_H measurement allows for significant better constraint of m_t
 - Indirect determination consistent with direct measurements
 - Remember: fully obtained from loop corrections!

Indirect result: $m_t = 176.1^{+2.9}_{-2.4}$ GeV

Tevatron: 173.2 ± 0.9 GeV
LHC: 173.3 ± 1.0 GeV

State of the SM: W versus top mass

- Scan of M_W vs m_t , with the direct measurements excluded from the fit.
- Results from Higgs measurement significantly reduces allowed indirect parameter space → corners the SM!




- Observed agreement demonstrates impressive consistency of the SM!


State of the SM: W mass versus $\sin^2\theta_{\text{eff}}^l$



- Scan of M_W vs $\sin^2\theta_{\text{eff}}^l$, with direct measurements excluded from the fit.
- Again, significant reduction allowed indirect parameter space from Higgs mass measurement.




- M_W and $\sin^2\theta_{\text{eff}}^l$ have become *the* sensitive probes of new physics!
 - Reason: both are ‘tree-level’ SM predictions.





- If energy scale of NP is high, BSM physics appears dominantly through vacuum polarization corrections
 - Aka, “oblique corrections”
- Oblique corrections reabsorbed into electroweak form factors
 - $\Delta\rho$, $\Delta\kappa$, Δr parameters, appearing in: M_W^2 , $\sin^2\theta_{\text{eff}}$, G_F , α , etc.
- Electroweak fit sensitive to BSM physics through oblique corrections
 - Similar to sensitivity to top and Higgs loop corrections.



- Oblique corrections from New Physics described through STU parametrization
[Peskin and Takeuchi, Phys. Rev. D46, 1 (1991)]

$$O_{\text{meas}} = O_{\text{SM,REF}}(m_H, m_t) + c_S \textcolor{red}{S} + c_T \textcolor{red}{T} + c_U \textcolor{red}{U}$$

- $\textcolor{red}{S}$: New Physics contributions to neutral currents
- $\textcolor{red}{T}$: Difference between neutral and charged current processes – sensitive to weak isospin violation
- $\textcolor{red}{U}$: (+S) New Physics contributions to charged currents. U only sensitive to W mass and width, usually very small in NP models (often: $U=0$)
- Also implemented: extended parameters (VWX), correction to $Z \rightarrow b\bar{b}$ couplings.
[Burgess et al., Phys. Lett. B326, 276 (1994)]
[Burgess et al., Phys. Rev. D49, 6115 (1994)]

Fit results for S, T, U

- S, T, U obtained from fit to the EW observables
- SM: $M_H = 126 \text{ GeV}$, $m_t = 173 \text{ GeV}$
 - This defines $(S, T, U) = (0, 0, 0)$
- SM: S, T depend logarithmically on M_H

Fit result:



$$S = 0.03 \pm 0.10$$

$$T = 0.05 \pm 0.12$$

$$U = 0.03 \pm 0.10$$

	S	T	U
S	1	+0.89	-0.54
T		1	-0.80
U			1

- Stronger constraints from fit with $U=0$.
- Also available for $Z \rightarrow b\bar{b}$ correction.
- No indication for new physics.
- Can now use this to constrain 4th gen, Ex-Dim, T-C, *Higgs couplings*, etc.



- Study of potential deviations of Higgs couplings from SM.
- BSM modeled as extension of SM through effective Lagrangian.
 - Consider leading corrections only.

- Popular benchmark model:

- Scaling of Higgs-vector boson (κ_V) and Higgs-fermion couplings (κ_F)
- No additional loops in the production or decay of the Higgs, no invisible Higgs decays and undetectable width.


$$L_V = \frac{h}{v} \left(2\kappa_V m_W^2 W_\mu W^\mu + \kappa_V m_Z^2 Z_\mu Z^\mu \right)$$

$$L_F = -\frac{h}{v} \left(\kappa_F m_t \bar{t}t + \kappa_F m_b \bar{b}b + \kappa_F m_\tau \bar{\tau}\tau \right)$$

- Main effect on EWPO due to modified Higgs coupling to gauge bosons (κ_V)



- Involving the longitudinal d.o.f.



- Most BSM models: $\kappa_V < 1$


- Additional Higgses typically give *positive* contribution to M_W .

- Main effect on EWPO due to Higgs coupling to gauge bosons (κ_V).


$$S = \frac{1}{12\pi} (1 - \kappa_V^2) \log \left(\frac{\Lambda^2}{M_H^2} \right), \quad T = -\frac{3}{16\pi c_W^2} (1 - \kappa_V^2) \log \left(\frac{\Lambda^2}{M_H^2} \right), \quad \Lambda = \frac{\lambda}{\sqrt{|1 - \kappa_V^2|}}$$

- Formulas from: Espinosa et al [arXiv:1202.3697]


- Cut-off scale Λ represents mass scale of new states that unitarize longitudinal gauge-boson scattering.
- (As required in this model.)
- λ is varied between 1 and 10 TeV, nominally fixed to 3 TeV ($4\pi v$).



ATLAS arXiv:1307.1427




CMS PAS-HIG-13-005





- Input: Higgs production times decay rate measurements (μ 's)
- Interpret as κ_V and κ_F using LHC HXSWG formalism.
- [arXiv:1209.0040]


Reproduction of ATLAS and CMS results



arXiv:1307.1427



CMS-PAS-HIG-13-005




■ Decent reproduction of ATLAS and CMS results within limited public-info available.

■ Private LHC combination:

- $K_V = 1.00 \pm 0.06$
- $K_F = 0.89 \pm 0.13$
- Perfectly consistent with SM.

■ Result from stand-alone EW fit:


- $K_V = 1.032^{+0.036}_{-0.025} (\lambda = 1 \text{ TeV})$
- $K_V = 1.024^{+0.024}_{-0.018} (\lambda = 3 \text{ TeV})$
- $K_V = 1.019^{+0.019}_{-0.014} (\lambda = 10 \text{ TeV})$



■ Note: some dependency in central value and error on cut-off scale λ .


1. EW fit sofar more precise result for κ_V than current LHC experiments.
2. EW fit: positive deviation of κ_V from 1.0.
 - (Many BSM models: $\kappa_V < 1$)

- EW fit: positive deviation of κ_V from one driven by small tension in W mass prediction versus measurement.




- Private LHC combination:

- $\kappa_V = 1.00 \pm 0.06$
- $\kappa_F = 0.89 \pm 0.13$



- Above: dependency on λ
- (Will be interesting to see how these measurements develop.)



Prospects for the Standard Model fit

Prospects of EW fit tested for two scenarios:

1. LHC Run-2+3
2. ILC with GigaZ(*)

(*) *GigaZ*:

- Operation of ILC at lower energies like Z-pole or WW threshold.
 - Allows to perform precision measurements of EW sector of the SM.
- At Z-pole, several billion Z's can be studied within 1-2 months.
- Physics of LEP1 and SLC can be revisited with few days of data.

In following studies:

Central values of input measurements adjusted to $M_H = 126$ GeV.

- *(Except where indicated.)*

Future Linear Collider can improve precision of EWPO's tremendously.

- *WW threshold scan + kinematic reconstruction, to obtain M_W*
 - From threshold scan: $\delta M_W : 15 \rightarrow 5 \text{ MeV}$
- *ttbar threshold scan, to obtain m_t*
 - Obtain m_t indirectly from production cross section: $\delta m_t : 0.9 \rightarrow 0.1 \text{ GeV}$
- *Z pole measurements*
 - High statistics: 10^9 Z decays: $\delta R^0_{\text{lep}} : 2.5 \cdot 10^{-2} \rightarrow 4 \cdot 10^{-3}$
 - With polarized beams, uncertainty on $\delta A^{0,f}_{LR} : 10^{-3} \rightarrow 10^{-4}$,
which translates to $\delta \sin^2 \theta^{\text{eff}}_l : 1.6 \cdot 10^{-4} \rightarrow 1.3 \cdot 10^{-5}$
- *H $\rightarrow$ ZZ and H $\rightarrow$ WW couplings: measured at 1% precision.*

ILC prospects: from ILC TDR (Vol-2).

LHC Run-2+3 (300/fb)

- *W mass measurement* : $\delta M_W : 15 \rightarrow 8 \text{ MeV}$
- *Final top mass measurement* m_t : $\delta m_t : 0.9 \rightarrow 0.6 \text{ GeV}$
- *H $\rightarrow$ ZZ and H $\rightarrow$ WW couplings*: measured at 4.5% precision.


LHC prospects: possibly optimistic scenario, but not impossible.

LHC Run-2+3 (300/fb)



- *W mass measurement* : $\delta M_W : 15 \rightarrow 8 \text{ MeV}$
- *Final top mass measurement* m_t : $\delta m_t : 0.9 \rightarrow 0.6 \text{ GeV}$
- *H $\rightarrow$ ZZ and H $\rightarrow$ WW couplings*: measured at 4.5% precision.

For both LHC-300 and ILC:

- *Low-energy data results to improve $\Delta\alpha_{\text{had}}$* :
 - ISR-based (BABAR), KLOE-II, VEPP-2000 (at energy below cc resonance), and BESIII e^+e^- cross-section measurements, in particular around cc resonance.
 - Plus: improved α_s , improvements in theory: $\Delta\alpha_{\text{had}}: 10^{-4} \rightarrow 5 \cdot 10^{-5}$
- *Assuming ~25% of today's theoretical uncertainties on M_W and $\sin^2\theta_{\text{eff}}^l$*
 - Implies three-loop EW calculations!
 - δM_W ($4 \rightarrow 1 \text{ MeV}$), $\delta \sin^2\theta_{\text{eff}}^l$ ($4.7 \times 10^{-5} \rightarrow 1 \times 10^{-5}$)
 - (Theoretical uncertainty estimates from recent Snowmass report)



- Logarithmic dependency on $M_H \rightarrow$ cannot compete with direct M_H meas.
- Indirect prediction M_H dominated by theory uncertainties.
 - No theory uncertainties: $M_H = 126 \pm 7$ GeV
 - With theory errors (R-fit scheme): $M_H = 126^{+10}_{-9}$ GeV
 - Present day theory uncertainties: $M_H = 126^{+20}_{-17}$ GeV
- If EWP-data central values unchanged, i.e. keep favoring low value of Higgs mass (94 GeV), $\sim 5\sigma$ discrepancy with measured Higgs mass.



- Huge reduction of uncertainty on indirect determinations of m_t , m_W , and $\sin^2\theta_{\text{eff}}^l$, by a factor of 3 or more.
- Assuming central values of m_t and M_W do not change, (at ILC) a deviation between the SM prediction and the direct measurements would be prominently visible.

Confrontation of measurement and prediction





- Breakdown of individual contributions to errors of M_W and $\sin^2\theta_{\text{eff}}^l$
- Parametric uncertainties (not the full fit).


Parameter	Scenario	error due to uncertainty ($\pm 1\sigma$)							
		δ_{meas}	δ_{pred}	δ_{exp}	δM_Z	δm_t	$\delta \Delta\alpha_{\text{had}}$	$\delta \alpha_s$	δ_{theo}
M_W [MeV]	Present	15	10.4	6.4	2.6	5.2	1.8	1.7	4.0
	LHC	8	5.8	4.8	2.6	3.6	0.9	1.7	1.0
	ILC	5	3.8	2.8	2.6	0.6	0.9	0.4	1.0
$\sin^2\theta_{\text{eff}}^l$ ($^{\circ}$)	Present	16	9.5	4.8	1.5	2.8	3.5	1.0	4.7
	LHC	16	4.1	3.1	1.5	1.9	1.6	1.0	1.0
	ILC	1.3	3.2	2.2	1.5	0.3	1.6	0.2	1.0

($^{\circ}$) In units of 10^{-5} .

- M_W and $\sin^2\theta_{\text{eff}}^l$ are sensitive probes of new physics! For all scenarios.
- At ILC/GigaZ, precision of M_Z will become important again!



- For STU parameters, improvement of factor of >3 is possible at ILC.
- Again, at ILC a deviation between the SM predictions and direct measurements would be prominently visible.
- Competitive results between EW fit and Higgs coupling measurements!
 - (At level of 1%.)




- Including M_H measurement, precise predictions of EW observables at loop level are possible for the first time.
- Overall consistency of the SM fit is very good.
 - M_H consistent at 1.3σ with indirect prediction from EW fit.
 - p-Value of global electroweak fit of SM: $18^{+2} \%$ (pseudo-experiments)

- Paradigm shift for EW fit: from Higgs mass prediction to consistency tests of the Standard Model.
- Knowledge of M_H dramatically improves SM prediction of key observables
 - M_W ($28 \rightarrow 11$ MeV), $\sin^2\theta_{\text{eff}}^l$ ($2.3 \times 10^{-5} \rightarrow 1.0 \times 10^{-5}$), m_t ($6.2 \rightarrow 2.5$ GeV)
 - Only surpassed so far by top mass measurement.

■ δM_W (indirect) =

- Large contributions to δM_W (and $\delta \sin^2\theta_{\text{eff}}^l$) from top and unknown higher-order EW corrections.

■ δM_W (direct) = 15 MeV



■ Improved accuracies set benchmark for new direct measurements!

- M_W , $\sin^2\theta_{\text{eff}}^l$ (and Higgs couplings) sensitive probes of new physics.

- Next step is evident: further exploration of Higgs couplings in the EW fit.
 - (Several groups already doing this. Gfitter too.)
- Prospects: including new data electroweak fits remain very interesting in coming years!
 - In particular ILC provides excellent New Physics sensitivity.
- Latest results always available at: <http://cern.ch/Gfitter>
 - Results of this presentation: EPJC 72, 2205 (2012)
 - *LHC-300 and ILC/GigaZ prospects paper to appear on arXiv this week !*

Thanks!



A **G**eneric **Fitter** Project for HEP Model Testing

Backup

Summary of indirect predictions



Parameter	Experimental input [$\pm 1\sigma$]			Indirect determination [$\pm 1\sigma_{\text{exp}} \pm 1\sigma_{\text{theo}}$]		
	Present	LHC	ILC/GigaZ	Present	LHC	ILC/GigaZ
M_H [GeV]	0.4	< 0.1	< 0.1	$^{+24}_{-20} \text{ } ^{+1}_{-2}$	$^{+20}_{-18} \text{ } ^{+2}_{-3}$	$^{+6.9}_{-6.6} \text{ } ^{+2.8}_{-2.3}$
M_W [MeV]	15	8	5	6.4 ± 4.6	5.0 ± 1.1	1.8 ± 1.7
M_Z [MeV]	2.1	2.1	2.1	10.6 ± 1.0	7.0 ± 0.9	2.6 ± 1.1
m_t [GeV]	0.9	0.6	0.1	$^{+2.3}_{-2.3} \text{ } ^{+0.4}_{-0.1}$	$^{+1.0}_{-1.8} \text{ } ^{+0.0}_{-0.4}$	$^{+0.7}_{-0.2} \text{ } ^{+0.3}_{-0.2}$
$\sin^2 \theta_{\text{eff}}^\ell$ [$\cdot 10^{-5}$]	16	16	1.3	4.8 ± 4.7	2.7 ± 1.2	2.0 ± 1.2
$\Delta \alpha_{\text{had}}^5 M_Z^2$ [$\cdot 10^{-5}$]	10	4.7	4.7	$^{+42}_{-42} \text{ } ^{+7}_{-1}$	$^{+35}_{-35} \text{ } ^{+5}_{-4}$	6 ± 3
R_l^0 [$\cdot 10^{-3}$]	25	25	4	—	—	—
$\alpha_s(M_Z^2)$ [$\cdot 10^{-4}$]	—	—	—	27 ± 1	27 ± 1	4 ± 1
$S _{U=0}$	—	—	—	0.10	0.10	0.03
$T _{U=0}$	—	—	—	0.12	0.07	0.03

- M_W and $\sin^2 \theta_{\text{eff}}^\ell$ are (and will be) sensitive probes of new physics!

- From: arXiv:1308.6176

Table 9: Selected set of precision measurements at TLEP. The statistical errors have been determined with (i) a one-year scan of the Z resonance with 50% data at the peak, leading to 710^{+1} Z visible decays, with resonant depolarization of single bunches for energy calibration at O(20min) intervals; (ii) one year at the Z peak with 40% longitudinally-polarized beams and a luminosity reduced to 20% of the nominal luminosity; (iii) a one-year scan of the WW threshold (around 161 GeV), with resonant depolarization of single bunches for energy calibration at O(20min) intervals; and (iv) a five-years scan of the $t\bar{t}$ threshold (around 346 GeV). The systematic uncertainties indicated below are only a “first look” estimate and will be revisited in the course of the design study.


Quantity	Physics	Present precision		Statistical uncertainty	Systematic uncertainty	Key	Challenge
m_Z (keV)	Input	91187500 ± 2100	Z Line shape scan	5 keV	< 100 keV	E_{beam} calibration	QED corrections
Γ_Z (keV)	$\Delta\rho$ (not $\Delta\alpha_{\text{had}}$)	2495200 ± 2300	Z Line shape scan	8 keV	< 100 keV	E_{beam} calibration	QED corrections
R_t	α_s, δ_b	20.767 ± 0.025	Z Peak	0.0001	< 0.001	Statistics	QED corrections
N_ν	PMNS Unitarity, ...	2.984 ± 0.008	Z Peak	0.00008	< 0.004		Bhabha scat.
N_ν	... and sterile ν 's	2.92 ± 0.05	$Z\gamma, 161$ GeV	0.001	< 0.001	Statistics	
R_b	δ_b	0.21629 ± 0.00066	Z Peak	0.000003	< 0.000060	Statistics, small IP	Hemisphere correlations
A_{LR}	$\Delta\rho, e_3, \Delta\alpha_{\text{had}}$	0.1514 ± 0.0022	Z peak, polarized	0.000015	< 0.000015	4 bunch scheme, 2exp	Design experiment
m_W (MeV)	$\Delta\rho, e_3, e_2, \Delta\alpha_{\text{had}}$	80385 ± 15	WW threshold scan	0.3 MeV	< 0.5 MeV	E_{beam} , Statistics	QED corrections
$m_{t\bar{t}}$ (MeV)	Input	173200 ± 900	$t\bar{t}$ threshold scan	10 MeV	< 10 MeV	Statistics	Theory interpretation

- Uncertainty estimates used:

Parameter	Experimental input [$\pm 1\sigma$]		
	Present	LHC	ILC/GigaZ
M_H [GeV]	0.4	< 0.1	< 0.1
M_W [MeV]	15	8	5
M_Z [MeV]	2.1	2.1	2.1
m_t [GeV]	0.9	0.6	0.1
$\sin^2\theta_{\text{eff}}^\ell$ [$\cdot 10^{-5}$]	16	16	1.3
$\Delta\alpha_{\text{had}}^5 M_Z^2$ [$\cdot 10^{-5}$]	10	4.7	4.7
R_l^0 [$\cdot 10^{-3}$]	25	25	4
$\alpha_s(M_Z^2)$ [$\cdot 10^{-4}$]	—	—	—

- ILC prospects from: ILC TDR (Vol-2).
- Theoretical uncertainty estimates from recent Snowmass report
- Central values of input measurements adjusted to $M_H = 126$ GeV.

- Scan of $\Delta\chi^2$ versus α_s
 - Also shown: SM fit with minimal inputs: M_Z , G_F , $\Delta\alpha_{\text{had}}^{(5)}(M_Z)$, $\alpha_s(M_Z)$, M_H , and fermion masses
- Determination of α_s at $N^3\text{LO}$.
 - Most sensitive through total hadronic cross-section σ_{had}^0 and partial leptonic width R_l^0



$$\alpha_s(M_Z) = 0.1190^{+0.0028}_{-0.0027} (\text{exp.}) \pm 0.0001 (\text{theo.})$$

- Theory uncertainty at per-mille level (obtained by scale variation of Γ_{had}).
- *In good agreement with value from τ decays, also at $N^3\text{LO}$, and with WA.*
 - (Improvements in precision only expected with ILC/GigaZ. See later.)


- The branching ratio R_b^0 : partial decay width of $Z \rightarrow b\bar{b}$ to $Z \rightarrow q\bar{q}$
- Freitas et al: full EW 2-loop calculation of $Z \rightarrow b\bar{b}$
- Contribution of same terms as in the calculation of $\sin^2\theta_{\text{eff}}^{b\bar{b}}$
→ cross-check of two results found good agreement
- Two-loop EW corrections now much smaller than experimental uncertainty (6.6×10^{-4})

M_H [GeV]	1-loop EW and QCD correction to FSR $\mathcal{O}(\alpha) + \text{FSR}_{\alpha, \alpha_s, \alpha_s^2}$ [10^{-4}]	2-loop EW correction $\mathcal{O}(\alpha_{\text{ferm}}^2)$ [10^{-4}]	2-loop EW and 2+3-loop QCD correction to FSR $\mathcal{O}(\alpha_{\text{ferm}}^2) + \text{FSR}_{\alpha_s^3, \alpha\alpha_s, m_b^2\alpha_s, m_b^4}$ [10^{-4}]	1+2-loop QCD correction to gauge boson self-energies $\mathcal{O}(\alpha\alpha_s, \alpha\alpha_s^2)$ [10^{-4}]
100	−35.66	−0.856	−2.496	−0.407
200	−35.85	−0.851	−2.488	−0.407
400	−36.09	−0.846	−2.479	−0.406

- Partial widths are defined inclusively: contain both QCD and QED contributions.
- Corrections expressed as so-called radiator functions $R_{A,f}$ and $R_{V,f}$

$$\Gamma_{f\bar{f}} = N_c^f \frac{G_F M_Z^3}{6\sqrt{2}\pi} (|g_{A,f}|^2 R_{A,f} + |g_{V,f}|^2 R_{V,f})^2$$

- High sensitivity to the strong coupling α_s
- Recently, full four-loop calculation of QCD Adler function became available ($N^3\text{LO}$)
- Much-reduced scale dependence!
- Theoretical uncertainty of 0.1 MeV, compared with experimental uncertainty of 2.0 MeV.




[P. Baikov et al., Phys. Rev. Lett. 108, 222003 (2012)]


[P. Baikov et al. Phys. Rev. Lett. 104, 132004 (2010)]

- Full EW one- and two-loop calculation of fermionic and bosonic contributions.
- One- and two-loop QCD corrections and leading terms of higher order corrections.
- Results for Δr include terms of order $O(\alpha)$, $O(\alpha\alpha_s)$, $O(\alpha\alpha_s^2)$, $O(\alpha^2_{\text{ferm}})$, $O(\alpha^2_{\text{bos}})$, $O(\alpha^2\alpha_s m_t^4)$, $O(\alpha^3 m_t^6)$
- Uncertainty estimate:
 - Missing terms of order $O(\alpha^2\alpha_s)$: about 3 MeV (from $O(\alpha^2\alpha_s m_t^4)$)
 - Electroweak three-loop correction $O(\alpha^3)$: < 2 MeV
 - Three-loop QCD corrections $O(\alpha_s^3)$: < 2 MeV
- Total: $\delta M_W \approx 4 \text{ MeV}$

[M Awramik et al., Phys. Rev. D69, 053006 (2004)]
 [M Awramik et al., Phys. Rev. Lett. 89, 241801 (2002)]



[A Freitas et al., Phys. Lett. B495, 338 (2000)]



Calculation of $\sin^2(\theta_{\text{eff}}^l)$

[M Awramik et al, Phys. Rev. Lett. 93, 201805 (2004)]
[M Awramik et al., JHEP 11, 048 (2006)]



- Effective mixing angle:

$$\sin^2 \theta_{\text{eff}}^{\text{lept}} = (1 - M_W^2/M_Z^2) (1 + \Delta\kappa)$$


- Two-loop EW and QCD correction to $\Delta\kappa$ known, leading terms of higher order QCD corrections.

- Fermionic two-loop correction about 10^{-3} , whereas bosonic one 10^{-5} .

- Uncertainty estimate obtained with different methods, geometric progression, leading to total of:
 $\delta\sin^2(\theta_{\text{eff}}^l) = 4.7 \times 10^{-5}$



- Difficult to define a pole mass for heavy, unstable and colored particle.
 - Single top decays before hadronizing. To have colorless final states, additional quarks needed.
 - *Non-perturb.* color-reconnection effects in fragmentation → biases in simulation.
 - ‘Renormalon’ ambiguity in top mass definition.
 - For pole mass, not for $\overline{\text{MS}}$ scheme.
 - Impact of finite top width effects.
- **Result: $m_t^{\text{exp}} \not\equiv m_t^{\text{pole}}$, and event-dependent.**
- The top mass extracted in hadron collisions is not well defined below a precision of $O(\Gamma_t) \sim 1 \text{ GeV}$
- Hard to estimate additional theo. uncertainties. With 0.5 GeV on m_t :
 - $M_H = 90^{+34}_{-21} \text{ GeV}$, $M_W = 80.359 \pm 0.013 \text{ GeV}$, $\sin^2 \theta_{\text{eff}}^l = 0.23148 \pm 0.00010$.
 - Only small deterioration in precision.



- From the Gfitter Group, EPJC 72, 2205 (2012)
- Left: full fit incl. M_H

Parameter	Input value	Free in fit	Fit Result
M_H [GeV] ^o	$125.7^{+0.4}_{-0.4}$	yes	$125.7^{+0.4}_{-0.4}$
M_W [GeV]	80.385 ± 0.015	–	$80.367^{+0.006}_{-0.007}$
Γ_W [GeV]	2.085 ± 0.042	–	2.091 ± 0.001
M_Z [GeV]	91.1875 ± 0.0021	yes	91.1878 ± 0.0021
Γ_Z [GeV]	2.4952 ± 0.0023	–	2.4954 ± 0.0014
σ_{had}^0 [nb]	41.540 ± 0.037	–	41.479 ± 0.014
R_ℓ^0	20.767 ± 0.025	–	20.740 ± 0.017
$A_{\text{FB}}^{0,\ell}$	0.0171 ± 0.0010	–	$0.01626^{+0.0001}_{-0.0002}$
$A_\ell^{(*)}$	0.1499 ± 0.0018	–	0.1472 ± 0.0007
$\sin^2\theta_{\text{eff}}^\ell(Q_{\text{FB}})$	0.2324 ± 0.0012	–	$0.23149^{+0.00010}_{-0.00008}$
A_c	0.670 ± 0.027	–	$0.6679^{+0.00034}_{-0.00028}$
A_b	0.923 ± 0.020	–	$0.93464^{+0.00005}_{-0.00007}$
$A_{\text{FB}}^{0,c}$	0.0707 ± 0.0035	–	0.0738 ± 0.0004
$A_{\text{FB}}^{0,b}$	0.0992 ± 0.0016	–	0.1032 ± 0.0005
R_c^0	0.1721 ± 0.0030	–	0.17223 ± 0.00006
R_b^0	0.21629 ± 0.00066	–	0.21548 ± 0.00005
$\overline{m}_c$ [GeV]	$1.27^{+0.07}_{-0.11}$	yes	$1.27^{+0.07}_{-0.11}$
$\overline{m}_b$ [GeV]	$4.20^{+0.17}_{-0.07}$	yes	$4.20^{+0.17}_{-0.07}$
m_t [GeV]	173.20 ± 0.87	yes	173.53 ± 0.82
$\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)^{(\dagger\Delta)}$	2757 ± 10	yes	2755 ± 11
$\alpha_s(M_Z^2)$	–	yes	$0.1190^{+0.0028}_{-0.0027}$
$\delta_{\text{th}} M_W$ [MeV]	$[-4, 4]_{\text{theo}}$	yes	4
$\delta_{\text{th}} \sin^2\theta_{\text{eff}}^\ell^{(\dagger)}$	$[-4.7, 4.7]_{\text{theo}}$	yes	–0.6

- From the Gfitter Group, EPJC 72, 2205 (2012)

- Left: full fit incl. M_H

- Middle: not incl. M_H

Parameter	Input value	Free in fit	Fit Result	Fit without M_H measurements
M_H [GeV] ^o	$125.7^{+0.4}_{-0.4}$	yes	$125.7^{+0.4}_{-0.4}$	94.7^{+25}_{-22}
M_W [GeV]	80.385 ± 0.015	–	$80.367^{+0.006}_{-0.007}$	$80.367^{+0.006}_{-0.007}$
Γ_W [GeV]	2.085 ± 0.042	–	2.091 ± 0.001	2.091 ± 0.001
M_Z [GeV]	91.1875 ± 0.0021	yes	91.1878 ± 0.0021	91.1878 ± 0.0021
Γ_Z [GeV]	2.4952 ± 0.0023	–	2.4954 ± 0.0014	2.4954 ± 0.0014
σ_{had}^0 [nb]	41.540 ± 0.037	–	41.479 ± 0.014	41.479 ± 0.014
R_ℓ^0	20.767 ± 0.025	–	20.740 ± 0.017	20.740 ± 0.017
$A_{\text{FB}}^{0,\ell}$	0.0171 ± 0.0010	–	$0.01626^{+0.0001}_{-0.0002}$	$0.01626^{+0.0001}_{-0.0002}$
$A_\ell^{(*)}$	0.1499 ± 0.0018	–	0.1472 ± 0.0007	0.1472 ± 0.0007
$\sin^2\theta_{\text{eff}}^\ell(Q_{\text{FB}})$	0.2324 ± 0.0012	–	$0.23149^{+0.00010}_{-0.00008}$	$0.23149^{+0.00010}_{-0.00008}$
A_c	0.670 ± 0.027	–	$0.6679^{+0.00034}_{-0.00028}$	$0.6679^{+0.00034}_{-0.00028}$
A_b	0.923 ± 0.020	–	$0.93464^{+0.00005}_{-0.00007}$	$0.93464^{+0.00005}_{-0.00007}$
$A_{\text{FB}}^{0,c}$	0.0707 ± 0.0035	–	0.0738 ± 0.0004	0.0738 ± 0.0004
$A_{\text{FB}}^{0,b}$	0.0992 ± 0.0016	–	0.1032 ± 0.0005	0.1032 ± 0.0005
R_c^0	0.1721 ± 0.0030	–	0.17223 ± 0.00006	0.17223 ± 0.00006
R_b^0	0.21629 ± 0.00066	–	0.21548 ± 0.00005	0.21548 ± 0.00005
$\overline{m}_c$ [GeV]	$1.27^{+0.07}_{-0.11}$	yes	$1.27^{+0.07}_{-0.11}$	$1.27^{+0.07}_{-0.11}$
$\overline{m}_b$ [GeV]	$4.20^{+0.17}_{-0.07}$	yes	$4.20^{+0.17}_{-0.07}$	$4.20^{+0.17}_{-0.07}$
m_t [GeV]	173.20 ± 0.87	yes	173.53 ± 0.82	173.53 ± 0.82
$\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)^{(\dagger\Delta)}$	2757 ± 10	yes	2755 ± 11	2755 ± 11
$\alpha_s(M_Z^2)$	–	yes	$0.1190^{+0.0028}_{-0.0027}$	$0.1190^{+0.0028}_{-0.0027}$
$\delta_{\text{th}} M_W$ [MeV]	$[-4, 4]_{\text{theo}}$	yes	4	4
$\delta_{\text{th}} \sin^2\theta_{\text{eff}}^{(\dagger)}$	$[-4.7, 4.7]_{\text{theo}}$	yes	–0.6	–0.5

Electroweak Fit – SM Fit Results



- From the Gfitter Group, EPJC 72, 2205 (2012)

- Left: full fit incl. M_H

- Middle: not incl. M_H

- Right: fit incl M_H , not the row

Parameter	Input value	Free in fit	Fit Result	Fit without M_H measurements	Fit without exp. input in line
M_H [GeV] ^o	$125.7^{+0.4}_{-0.4}$	yes	$125.7^{+0.4}_{-0.4}$	94.7^{+25}_{-22}	94.7^{+25}_{-22}
M_W [GeV]	80.385 ± 0.015	–	$80.367^{+0.006}_{-0.007}$	$80.367^{+0.006}_{-0.007}$	80.360 ± 0.011
Γ_W [GeV]	2.085 ± 0.042	–	2.091 ± 0.001	2.091 ± 0.001	2.091 ± 0.001
M_Z [GeV]	91.1875 ± 0.0021	yes	91.1878 ± 0.0021	91.1878 ± 0.0021	91.1978 ± 0.0114
Γ_Z [GeV]	2.4952 ± 0.0023	–	2.4954 ± 0.0014	2.4954 ± 0.0014	2.4950 ± 0.0017
σ_{had}^0 [nb]	41.540 ± 0.037	–	41.479 ± 0.014	41.479 ± 0.014	41.471 ± 0.015
R_ℓ^0	20.767 ± 0.025	–	20.740 ± 0.017	20.740 ± 0.017	20.715 ± 0.026
$A_{\text{FB}}^{0,\ell}$	0.0171 ± 0.0010	–	$0.01626^{+0.0001}_{-0.0002}$	$0.01626^{+0.0001}_{-0.0002}$	0.01624 ± 0.0002
$A_\ell^{(*)}$	0.1499 ± 0.0018	–	0.1472 ± 0.0007	0.1472 ± 0.0007	–
$\sin^2\theta_{\text{eff}}^\ell(Q_{\text{FB}})$	0.2324 ± 0.0012	–	$0.23149^{+0.00010}_{-0.00008}$	$0.23149^{+0.00010}_{-0.00008}$	0.23150 ± 0.00009
A_c	0.670 ± 0.027	–	$0.6679^{+0.00034}_{-0.00028}$	$0.6679^{+0.00034}_{-0.00028}$	0.6680 ± 0.00031
A_b	0.923 ± 0.020	–	$0.93464^{+0.00005}_{-0.00007}$	$0.93464^{+0.00005}_{-0.00007}$	0.93463 ± 0.00006
$A_{\text{FB}}^{0,c}$	0.0707 ± 0.0035	–	0.0738 ± 0.0004	0.0738 ± 0.0004	0.0737 ± 0.0004
$A_{\text{FB}}^{0,b}$	0.0992 ± 0.0016	–	0.1032 ± 0.0005	0.1032 ± 0.0005	0.1034 ± 0.0003
R_c^0	0.1721 ± 0.0030	–	0.17223 ± 0.00006	0.17223 ± 0.00006	0.17223 ± 0.00006
R_b^0	0.21629 ± 0.00066	–	0.21548 ± 0.00005	0.21548 ± 0.00005	0.21547 ± 0.00005
$\overline{m}_c$ [GeV]	$1.27^{+0.07}_{-0.11}$	yes	$1.27^{+0.07}_{-0.11}$	$1.27^{+0.07}_{-0.11}$	–
$\overline{m}_b$ [GeV]	$4.20^{+0.17}_{-0.07}$	yes	$4.20^{+0.17}_{-0.07}$	$4.20^{+0.17}_{-0.07}$	–
m_t [GeV]	173.20 ± 0.87	yes	173.53 ± 0.82	173.53 ± 0.82	$176.11^{+2.88}_{-2.35}$
$\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)^{(\dagger\Delta)}$	2757 ± 10	yes	2755 ± 11	2755 ± 11	2718^{+49}_{-43}
$\alpha_s(M_Z^2)$	–	yes	$0.1190^{+0.0028}_{-0.0027}$	$0.1190^{+0.0028}_{-0.0027}$	0.1190 ± 0.0027
$\delta_{\text{th}} M_W$ [MeV]	$[-4, 4]_{\text{theo}}$	yes	4	4	–
$\delta_{\text{th}} \sin^2\theta_{\text{eff}}^{(\dagger)}$	$[-4.7, 4.7]_{\text{theo}}$	yes	–0.6	–0.5	–

Electroweak Fit – SM Fit Results



■ From the Gfitter Group, EPJC 72, 2205 (2012)

■ Left: full fit incl. M_H

■ Middle: not incl. M_H

■ Right: fit incl M_H , not the row

Parameter	Input value	Free in fit	Fit Result	Fit without M_H measurements	Fit without exp. input in line
M_H [GeV] ^o	$125.7^{+0.4}_{-0.4}$	yes	$125.7^{+0.4}_{-0.4}$	94.7^{+25}_{-22}	94.7^{+25}_{-22}
M_W [GeV]	80.385 ± 0.015	–	$80.367^{+0.006}_{-0.007}$	$80.367^{+0.006}_{-0.007}$	80.360 ± 0.011
Γ_W [GeV]	2.085 ± 0.042	–	2.091 ± 0.001	2.091 ± 0.001	2.091 ± 0.001
M_Z [GeV]	91.1875 ± 0.0021	yes	91.1878 ± 0.0021	91.1878 ± 0.0021	91.1978 ± 0.0114
Γ_Z [GeV]	2.4952 ± 0.0023	–	2.4954 ± 0.0014	2.4954 ± 0.0014	2.4950 ± 0.0017
σ_{had}^0 [nb]	41.540 ± 0.037	–	41.479 ± 0.014	41.479 ± 0.014	41.471 ± 0.015
R_ℓ^0	20.767 ± 0.025	–	20.740 ± 0.017	20.740 ± 0.017	20.715 ± 0.026
$A_{\text{FB}}^{0,\ell}$	0.0171 ± 0.0010	–	$0.01626^{+0.0001}_{-0.0002}$	$0.01626^{+0.0001}_{-0.0002}$	0.01624 ± 0.0002
$A_\ell^{(*)}$	0.1499 ± 0.0018	–	0.1472 ± 0.0007	0.1472 ± 0.0007	–
$\sin^2\theta_{\text{eff}}^\ell(Q_{\text{FB}})$	0.2324 ± 0.0012	–	$0.23149^{+0.00010}_{-0.00008}$	$0.23149^{+0.00010}_{-0.00008}$	0.23150 ± 0.00009
A_c	0.670 ± 0.027	–	$0.6679^{+0.00034}_{-0.00028}$	$0.6679^{+0.00034}_{-0.00028}$	0.6680 ± 0.00031
A_b	0.923 ± 0.020	–	$0.93464^{+0.00005}_{-0.00007}$	$0.93464^{+0.00005}_{-0.00007}$	0.93463 ± 0.00006
$A_{\text{FB}}^{0,c}$	0.0707 ± 0.0035	–	0.0738 ± 0.0004	0.0738 ± 0.0004	0.0737 ± 0.0004
$A_{\text{FB}}^{0,b}$	0.0992 ± 0.0016	–	0.1032 ± 0.0005	0.1032 ± 0.0005	0.1034 ± 0.0003
R_c^0	0.1721 ± 0.0030	–	0.17223 ± 0.00006	0.17223 ± 0.00006	0.17223 ± 0.00006
R_b^0	0.21629 ± 0.00066	–	0.21548 ± 0.00005	0.21548 ± 0.00005	0.21547 ± 0.00005
$\overline{m}_c$ [GeV]	$1.27^{+0.07}_{-0.11}$	yes	$1.27^{+0.07}_{-0.11}$	$1.27^{+0.07}_{-0.11}$	–
$\overline{m}_b$ [GeV]	$4.20^{+0.17}_{-0.07}$	yes	$4.20^{+0.17}_{-0.07}$	$4.20^{+0.17}_{-0.07}$	–
m_t [GeV]	173.20 ± 0.87	yes	173.53 ± 0.82	173.53 ± 0.82	$176.11^{+2.88}_{-2.35}$
$\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)^{(\dagger\Delta)}$	2757 ± 10	yes	2755 ± 11	2755 ± 11	2718^{+49}_{-43}
$\alpha_s(M_Z^2)$	–	yes	$0.1190^{+0.0028}_{-0.0027}$	$0.1190^{+0.0028}_{-0.0027}$	0.1190 ± 0.0027
$\delta_{\text{th}} M_W$ [MeV]	$[-4, 4]_{\text{theo}}$	yes	4	4	–
$\delta_{\text{th}} \sin^2\theta_{\text{eff}}^\ell^{(\dagger)}$	$[-4.7, 4.7]_{\text{theo}}$	yes	–0.6	–0.5	–

Moriond 2011: Prediction for Higgs mass

LEP + Tevatron (Fall 2010) :



- CL_{s+b}^{2s} central value $\pm 1\sigma$: $M_H = 120.2_{-5.2}^{+17.9}$ GeV
- 2σ interval:
 $-2\ln Q$: [115,152] GeV
 $CL_{s+b}^{2-sided}$: [114,155] GeV

LEP + Tevatron (Moriond 2011) :


- CL_{s+b}^{2s} central value $\pm 1\sigma$: $M_H = 120.2_{-4.7}^{+12.3}$ GeV
- 2σ interval:
 $-2\ln Q$: [115,138] GeV
 $CL_{s+b}^{2-sided}$: [114,149] $\cup$ [152,155] GeV

Fit with LEP + Tevatron + LHC (H $\rightarrow$ WW) searches (Moriond 2011) :

- Central value unchanged
- 2σ interval:
 $-2\ln Q$: [115,137] GeV
 $CL_{s+b}^{2-sided}$: [114,14?] GeV



- Low energy observables with interesting precision will soon become available.




- Input correlation coefficients between Z pole measurements

	M_Z	Γ_Z	σ_{had}^0	R_ℓ^0	$A_{\text{FB}}^{0,\ell}$		$A_{\text{FB}}^{0,c}$	$A_{\text{FB}}^{0,b}$	A_c	A_b	R_c^0	R_b^0
M_Z	1	-0.02	-0.05	0.03	0.06	$A_{\text{FB}}^{0,c}$	1	0.15	0.04	-0.02	-0.06	0.07
Γ_Z		1	-0.30	0.00	0.00	$A_{\text{FB}}^{0,b}$		1	0.01	0.06	0.04	-0.10
σ_{had}^0			1	0.18	0.01	A_c			1	0.11	-0.06	0.04
R_ℓ^0				1	-0.06	A_b				1	0.04	-0.08
$A_{\text{FB}}^{0,\ell}$					1	R_c^0					1	-0.18

Table 2: Correlation matrices for observables determined by the Z lineshape fit (left), and by heavy flavour analyses at the Z pole (right) [56].

Full EW 2-loop calculations

- Recent paper by A. Freitas, arXiv:1310.2256.
- Contains full two-loop fermionic EW corrections to the Z-boson width and production rate.
- Only small impact on EW fit results compared with 1-loop results.



Experimental inputs – Predicted uncertainties



Parameter	Experimental input [$\pm 1\sigma$]			
	Present	LHC	ILC/GigaZ	TLEP
M_H [GeV]	0.4 $\Rightarrow$	< 0.1	< 0.1	< 0.1
M_W [MeV]	15 $\Rightarrow$	8 $\Rightarrow$	5 $\Rightarrow$	1.3
M_Z [MeV]	2.1	2.1	2.1	$\Rightarrow$ 0.1
m_t [GeV]	0.9 $\Rightarrow$	0.6	0.1	0.08
Γ_Z [MeV]	2.3	2.3	$\Rightarrow$ 0.8	$\Rightarrow$ 0.1
$\sin^2\theta_{\text{eff}}^\ell$ [$\cdot 10^{-5}$]	16	16	$\Rightarrow$ 1.3	$\Rightarrow$ 0.3
R_l^0 [$\cdot 10^{-3}$]	25	25	$\Rightarrow$ 4	$\Rightarrow$ 1.3
$\Delta\alpha_{\text{had}}^5(M_Z^2)$ [$\cdot 10^{-5}$]	10 $\Rightarrow$	4.7	4.7	4.7
$\alpha_S(M_Z^2)$ [$\cdot 10^{-4}$]	—	—	—	—
$\delta_{\text{th}} M_W$ [MeV]	4 $\Rightarrow$	1	1	1
$\delta_{\text{th}} \sin^2\theta_{\text{eff}}^\ell$ [$\cdot 10^{-5}$]	4.7 $\Rightarrow$	1	1	1


TLEP scenario:

- *Preliminary estimates*
- Clearly not the same level of understanding as LHC or ILC.
- Uncertainties may turn out completely different.
 - From arXiv:1308.6176,
 - and Snowmass report.
 - Of these two, we take most conservative estimate.
- Note: top mass dominated by theoretical uncertainty.
- Higher statistics
- From beam energy precision: improved M_Z and Γ_Z


Prospects of the EW fit: Higgs mass (126 GeV)



Present / LHC / ILC



Future scenario




- Logarithmic dependency on $M_H \rightarrow$ cannot compete with direct M_H meas.
- Indirect prediction M_H dominated by theory uncertainties.
 - ILC with (without) theory errors: $M_H = 126^{+10}_{-9} (\pm 7)$ GeV
 - ILC with present-day theory uncertainties: $M_H = 126^{+20}_{-17}$ GeV
 - TLEP with (without) theory errors: $M_H = 126 \pm 5 (\pm 3)$ GeV


Prospects of the EW fit: Higgs mass (94 GeV)



Present / LHC / ILC



Future scenario




- If EWP-data central values are unchanged, i.e. they keep favoring low value of Higgs mass (94 GeV), $>5\sigma$ discrepancy with measured Higgs mass.
 - In both ILC and TLEP scenarios.




Prospects of the EW fit: W mass and $\sin^2\theta_{\text{eff}}^l$



Present / LHC / ILC




Future scenario




Prospects of the EW fit: W mass versus $\sin^2\theta_{\text{eff}}^l$



Present / LHC / ILC



Future scenario



- Huge reduction of uncertainty on indirect determinations of m_W , and $\sin^2\theta_{\text{eff}}^l$, by a factor of ≈ 3 (≈ 4 -5) at ILC (TLEP).
- Assuming central values of M_W and $\sin^2\theta_{\text{eff}}^l$ do not change, a deviation between the SM prediction and the direct measurements would be prominently visible, at both ILC and TLEP.
 - But also in LHC-300 scenario, from improved theory uncertainties.

Confrontation of measurement and prediction



- Breakdown of individual contributions to errors of M_W and $\sin^2\theta_{\text{eff}}^l$
- Parametric uncertainties (not the full fit).


Parameter	Scenario	error due to uncertainty ($\pm 1\sigma$)							
		δ_{meas}	δ_{pred}	δ_{exp}	δM_Z	δm_t	$\delta \Delta\alpha_{\text{had}}$	$\delta\alpha_s$	δ_{theo}
M_W [MeV]	Present	15	10.4	6.4	2.6	5.2	1.8	1.7	4.0
	LHC	8	5.8	4.8	2.6	3.6	0.9	1.7	1.0
	ILC	5	3.8	2.8	2.6	0.6	0.9	0.4	1.0
	Future	1.3	2.0	1.0	0.1	0.5	0.9	0.3	1.0
$\sin^2\theta_{\text{eff}}^l$ ($^{\circ}$)	Present	16	9.5	4.8	1.5	2.8	3.5	1.0	4.7
	LHC	16	4.1	3.1	1.5	1.9	1.6	1.0	1.0
	ILC	1.3	3.2	2.2	1.5	0.3	1.6	0.2	1.0
	Future	0.3	2.7	1.7	0.1	0.3	1.6	0.2	1.0

($^{\circ}$) In units of 10^{-5} .


- M_W and $\sin^2\theta_{\text{eff}}^l$ are sensitive probes of new physics! In all scenarios.
- At ILC/GigaZ, precision of M_Z will become important again.
- At TLEP ('Future'), limited by external inputs: theory errors and $\Delta\alpha_{\text{had}}$

Prospects of the EW fit: W versus top mass

Present / LHC / ILC




Future scenario




- Huge reduction of uncertainty on indirect determinations of m_t and m_W by a factor of $\gtrsim 3$ ($\gtrsim 5$) at ILC (TLEP).
- Assuming central values of m_t and M_W do not change, a deviation between the SM prediction and the direct measurements would be prominently visible.

Prospects of EW fit: S versus T

Present / LHC / ILC



Future scenario



- For STU parameters, improvement of factor of ≥ 4 (≥ 10) is possible at ILC (TLEP).
- Again, at both ILC and TLEP a deviation between the SM predictions and direct measurements would be prominently visible.

Predicted uncertainties from EW fit



Parameter	error due to uncertainty ($\pm 1\sigma$)									
	δ_{meas}	$\delta_{\text{fit}}^{\text{tot}}$	$\delta_{\text{fit}}^{\text{exp}}$	$\delta_{\text{fit}}^{\text{theo}}$	δM_W	δM_Z	δm_t	$\delta \sin^2 \theta_{\text{eff}}^{(\circ)}$	$\delta \Delta \alpha_{\text{had}}^{(\circ)}$	$\delta \alpha_S^{(\Delta)}$
ILC prospects										
M_H [GeV]	< 0.1	$^{+9.6}_{-9.0}$	$^{+6.9}_{-6.6}$	$^{+2.7}_{-2.4}$	$^{+4.2}_{-0.8}$	$^{+4.4}_{-4.0}$	$^{+0.9}_{-0.8}$	$^{+3.1}_{-3.3}$	$^{+4.2}_{-4.1}$	$^{+0.6}_{-0.6}$
M_W [MeV]	5	3.6	1.9	1.7	—	1.7	0.3	1.2	0.7	0.2
M_Z [MeV]	2.1	3.7	2.6	1.1	2.4	—	0.5	1.3	1.9	0.3
m_t [GeV]	0.1	1.0	0.7	$^{+0.3}_{-0.2}$	$^{+0.5}_{-0.6}$	0.5	—	$^{+0.3}_{-0.2}$	0.4	—
$\sin^2 \theta_{\text{eff}}^{(\circ)}$	1.3	3.2	2.0	1.2	1.7	1.2	0.2	—	1.5	0.1
$\Delta \alpha_{\text{had}}^{(\circ)}$	4.7	8.6	5.7	2.9	2.5	4.2	0.8	3.9	—	0.5
Future prospects										
M_H [GeV]	< 0.1	5.3	3.3	2.0	3.0	0.3	1.0	$^{+0.0}_{-1.2}$	3.2	0.6
M_W [MeV]	1.3	1.9	0.4	1.5	—	0.1	0.3	0.2	0.1	0.1
M_Z [MeV]	0.1	1.5	1.0	0.5	1.0	—	0.3	—	0.9	0.4
m_t [GeV]	0.08	0.38	0.24	0.14	0.24	0.03	—	0.01	0.22	0.02
$\sin^2 \theta_{\text{eff}}^{(\circ)}$	0.3	$^{+2.8}_{-2.4}$	1.4	$^{+1.5}_{-1.1}$	1.2	—	0.1	—	1.3	0.5
$\Delta \alpha_{\text{had}}^{(\circ)}$	4.7	0.4	0.1	0.3	—	—	0.1	0.1	—	—

$^{(\circ)}$ In units of 10^{-5} . $^{(\Delta)}$ In units of 10^{-4}

- Breakdown of uncertainties derived from EW fit. (Note: *correlated* errors.)
- Compared to parametric breakdown: reduced experimental, but increased theory errors. Slightly smaller total errors.